

POTENTIOMETRIC SURFACE MAP OF THE BEDROCK AQUIFERS OF HAMILTON COUNTY, INDIANA


Hamilton County, Indiana is located in the central portion of the state. The entire county is situated within the White and West Fork White River Basin. The Potentiometric Surface Map (PSM) of the Bedrock aquifers of Hamilton County was mapped by contouring the elevations of over 1400 static water-levels reported on well records received primarily over a 50 year period. These wells are completed in unconsolidated aquifers at various depths, and typically, under confined conditions (bounded by impermeable layers above and below the water bearing formation). However, some wells were completed under unconfined (not bounded by impermeable layers) settings. The potentiometric surface is a measure of the pressure on water in a water bearing formation. Water in an unconfined aquifer is at atmospheric pressure and will not rise in a well above the top of the water bearing formation, in contrast to water in a confined aquifer which is under hydrostatic pressure and will rise in a well above the top of the water bearing formation.

Static water-level measurements in individual wells used to construct county PSM's are indicative of the water-level at the time of well completion. The groundwater level within an aquifer constantly fluctuates in response to rainfall, evapotranspiration, groundwater movement, and pumping. Therefore, current site specific conditions may differ due to local or seasonal variations in measured static water levels. Because fluctuations in groundwater are typically small, static water-levels can be used to construct a generalized PSM. Groundwater flow is naturally from areas of recharge toward areas of discharge. As a general rule, but certainly not always, groundwater flow approximates the overlying topography and intersects the land surface at major streams. The contour type was determined based on the amount of data and the degree of change in water levels between wells in each mapped area. Portions of the county are lacking in data and/or are covered by deposits that have limited to non-existent aquifer potential. Therefore, potentiometric surface elevations contours have not been extended through these areas.


Universal Transverse Mercator (UTM) coordinates for the water wells were either physically obtained in the field, determined through address geocoding, or reported on water well records; however, the location of the majority of the water well records used to make the PSM were address geocoded. Elevation data were either obtained from topographic maps or a digital elevation model. Quality control/quality assurance procedures were utilized to refine or remove data where errors were readily apparent.

Bedrock potentiometric surface elevations in Hamilton County range from a high of 900 feet mean sea level (msl) in the northwest region of the county, to a low of 720 feet msl in the south-central portion. Groundwater flow direction within the White and West Fork White River Basin is generally towards the White River. In the far western portion of the county groundwater flows west towards Eagle Creek and Little Eagle Creek. Also, in the southeast corner groundwater flows towards Fall Creek.

The county PSM can be used to define the regional groundwater flow path and to identify significant areas of groundwater recharge and discharge. County PSM's represent overall regional characteristics and are not intended to be a substitute for site-specific studies.




Hillsbade Map of Hamilton County, Indiana



Vertical Exaggeration = 5x

EXPLANATION

- 800 Line of equal elevation, in feet above mean sea level
- Stream
- County Road
- State Road
- US Highway
- Interstate
- Basin Boundary
- Municipal Boundary
- Lake & River
- No Aquifer Material or Limited Data



Map Use and Disclaimer Statement

We request that the following agency be acknowledged in products derived from this map: Indiana Department of Natural Resources, Division of Water. This map was compiled by staff of the Indiana Department of Natural Resources, Division of Water using data believed to be reasonably accurate. However, a degree of error is inherent in all maps. This product is distributed "as is" without warranties of any kind, either expressed or implied. This map is intended for use only at the published scale.

This map was created from several existing shapefiles. Township and Range Lines of Indiana (line shapefile, 20020621), Land Survey Lines of Indiana (polygon shapefile, 20020621), and County Boundaries of Indiana (polygon shapefile, 20020621), were all from the Indiana Geological Survey and based on a 1:24,000 scale. Draft road shapefiles, System1 and System2 (line shapefiles, 2003), were from the Indiana Department of Transportation and based on a 1:24,000 scale. Populated Areas in Indiana 2000 (polygon shapefile, 20021000) was from the U.S. Census Bureau and based on a 1:100,000 scale. Hydrography, Streams (NHDI) (line shapefile, 20081218), Rivers (NHDI) (polygon shapefile, 20081218), Lakes (NHDI) (polygon shapefile, 20081218) was from the U.S. Geological Survey and the U.S. Environmental Protection Agency and based on a 1:24,000 scale. Managed Lands INDR IN (polygon shapefile, 20100920) was from INDR and based on a 1:24,000 scale. No Aquifer Material or Limited Data Hamilton County, Indiana (polygon shapefile, Grove, 2012). County Hillsbade image was from the U.S. Geological Survey National Elevation Dataset (raster image, 20120720). Potentiometric Surface Map of the Bedrock Aquifers of Hamilton County, Indiana (line shapefiles, Grove, 2012) was based on a 1:24,000 scale.

**Potentiometric Surface Map of the
 Bedrock Aquifers of Hamilton County, Indiana**
 by
 Glenn E. Grove
 Division of Water, Resource Assessment Section
 December 2012