## OUTREACH SERVICES OF INDIANA FSSA – State of Indiana Diagram Protects 1

## Diastat Protocol

Baseline Vital	Signs: Blood P	ressure Pu	lse Temper	rature Resp	oirations	
The following is intended as a guideline. This protocol does not supersede facility policy, nursing judgment or physician orders.						
Į.						
When to Give?  For Seizures lasting more than minutes. For more than seizures inminutes/hours  Special Instructions						
* How Much to Give? * Mg Always DBL check medication with the order/MAR prior to administering!						
Do Not Repeat in 24 hours unless otherwise ordered by physician.						
				<u> </u>		
How to Give?  See attached instructionsSee package insert located						
Call 911 if seizure continues for minutes after Diastat or if the person appears gravely ill or difficulty breathing.						
<ul> <li>Monitoring After Diastat Given</li> <li>Monitor for changes in breathing (shallow, labored), blue lips or fingernails. If noted, Call 911, attempt to sit person upright and attempt to arouse.</li> <li>Stay with the person for at least 4 hours after giving Diastat or until back to baseline.</li> <li>Monitor Vitals signs everyminutes for 4 hours.</li> <li>If BP is greater thanless than, Temp is greater thanless than, Pulse is greater thanless than, Notify nurse Supervisor</li> <li>Document Vital Signs on Vital Sign Flow Sheet Daily Notes</li> <li>Other</li> <li>Do not give anything to eat or drink until returns to normal activity. Start with small sips of water to be sure can tolerate.</li> <li>Ambulate with assist until fully awake.</li> <li>Instruct person not to operate heavy machinery, drive, swim or ride a bicycle for at least 4 hours and fully awake.</li> <li>Document overall condition during and after receiving diastat every minutes on Daily Note</li> <li>Other:</li> </ul>						
Completed by: Date:						
Physician Signature:					e:	(annually)
Review Date:						
Reviewer:						

Reference: www.diastat.com