

# STD MORBIDITY

## Chlamydia, Gonorrhea and Syphilis Morbidity – January – December 2017

Gender	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Female	23,479	69.0%	5,887	50.1%	28	9.1%
Male	10,637	31.0%	5,860	49.9%	281	90.9%
Unknown	11	<1%	6	<1%	0	0%
Total	34,127	100%	11,753	100%	309	100%
Race	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
White	15,253	44.7%	4,379	37.3%	135	43.7%
Black	11,736	34.4%	5,739	48.8%	139	45.0%
All Others	3,749	11.0%	881	7.5%	33	10.7%
Unknown	3,389	9.9%	754	6.4%	2	<1%
Total	34,127	100%	11,753	100%	309	100%
Ethnicity	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Hispanic	2,377	7.0%	472	4.0%	28	8.0%
Non-Hispanic	27,493	80.6%	10,251	87.2%	278	92.0%
Unknown	4,257	12.5%	1,030	8.8%	3	<1%
Total	34,127	100%	11,753	100%	309	100%
Age	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
0-19	9,954	29.2%	2,484	21.1%	21	6.8%
20-24	13,147	38.5%	3,601	30.6%	68	22.0%
25-29	6,078	17.8%	2,578	22.0%	84	27.2%
30-34	2,545	7.5%	1,286	10.9%	41	13.3%
35-39	1,236	3.6%	792	6.7%	30	9.7%
40-44	548	1.6%	397	3.4%	19	6.2%
45-54	485	1.4%	443	3.8%	35	11.3%
55+	134	<1%	172	1.5%	11	3.6%
Unknown	0	0%	0	0%	0	0%
Total	34,127	100%	11,753	100%	309	100%

Source: ISDH STD Database as of February 15, 2018

NOTE: Not all percents may add to 100 due to rounding.