

**Table 3F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Racial and Ethnic Distribution by Local Agency**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaska Native %</b>	<b>Asian/Pacific Islander %</b>	<b>Multiple Races %</b>	<b>All Other/Unknown %</b>
BCA	202	92.6	3.0	2.5	0.0	1.0	0.0	1.0
BCHD	526	81.9	1.7	14.3	0.0	0.6	0.8	0.8
BRSI	663	94.6	0.6	2.4	0.0	0.3	0.9	1.2
CAP	594	91.2	0.8	5.7	0.2	0.5	0.8	0.7
CCHD	243	95.5	1.2	0.4	0.0	0.4	1.6	0.8
CFS	1,039	92.5	0.5	5.0	0.1	0.1	0.8	1.1
DCH	577	95.1	0.7	1.0	0.3	0.0	0.9	1.9
DCHD	252	75.4	0.0	21.0	0.0	2.0	1.2	0.4
DMH	285	96.1	0.0	1.1	0.0	1.1	0.4	1.4
DMHC	1,048	88.5	4.1	4.1	0.2	0.7	1.4	1.0
ECHD	568	6.9	37.9	53.9	0.2	0.4	0.4	0.5
ELKCHD	1,943	55.8	9.8	31.9	0.0	0.9	1.0	0.6
FHHC	75	*	*	*	*	*	*	*
FLCHD	476	77.3	13.0	5.0	0.2	1.9	1.7	0.8
FULCHD	143	90.2	0.0	9.1	0.0	0.0	0.7	0.0
GCRC	298	94.3	2.7	1.0	0.7	0.7	0.3	0.3
HCHF	421	85.3	6.2	3.3	0.0	3.8	0.0	1.4
HCMH&HS	296	94.9	1.4	2.4	0.0	0.0	1.0	0.3
HIS	582	91.4	1.0	6.2	0.2	0.3	0.3	0.5
HU	57	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Racial and Ethnic Distribution by Local Agency**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaska Native %</b>	<b>Asian/Pacific Islander %</b>	<b>Multiple Races %</b>	<b>All Other/Unknown %</b>
ICAP	300	93.3	1.0	3.0	0.3	0.3	1.3	0.7
IHC	2,665	76.6	7.1	12.9	0.1	0.9	1.5	0.9
JEFCHD	224	85.7	1.3	7.6	0.0	0.9	2.2	2.2
JNHC	338	94.7	1.2	2.1	0.0	0.9	0.6	0.6
JOCHD	730	87.8	2.1	4.9	0.0	2.6	1.2	1.4
KCHD	441	77.6	1.1	20.0	0.0	0.5	0.0	0.9
LCHD	145	83.4	1.4	12.4	0.0	0.7	0.0	2.1
MADCHD	957	78.3	12.0	7.6	0.2	0.2	1.0	0.6
MARCHD	9,411	34.1	40.1	19.9	0.1	3.6	1.5	0.7
MARSCHD	373	76.4	0.3	22.0	0.0	0.3	1.1	0.0
MCMH	423	96.9	0.7	1.7	0.0	0.0	0.2	0.5
MHS	1,958	48.7	28.8	16.4	0.3	1.8	2.6	1.3
MW	419	89.3	1.9	2.9	0.0	0.5	5.3	0.2
NCCAA	878	68.1	19.1	9.0	0.0	0.8	2.2	0.8
NHC	3,361	57.0	21.8	13.9	0.1	4.4	2.1	0.7
NHS	1,301	79.6	5.8	11.5	0.2	0.6	1.6	0.7
NWFS	4,723	45.7	34.3	17.2	0.1	0.7	1.1	0.9
ODCS	848	83.4	11.3	2.0	0.1	0.6	2.1	0.5
PHNA	821	85.3	3.8	3.4	0.0	5.1	2.1	0.4
St Mary""s	549	95.1	1.5	1.6	0.0	0.2	0.5	1.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Racial and Ethnic Distribution by Local Agency**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
STARCHD	171	96.5	0.0	2.3	0.0	0.6	0.0	0.6
STEUBCHD	275	89.5	0.7	6.2	0.0	1.1	0.4	2.2
SVJH	284	95.4	0.7	2.1	0.7	0.4	0.7	0.0
TCHD	1,273	61.5	11.5	18.1	0.2	6.4	1.1	1.2
VCHD	1,021	73.5	16.7	5.8	0.3	0.8	2.3	0.7
WCHD	184	97.3	0.5	1.1	0.0	0.5	0.5	0.0
WCMH	295	82.4	0.0	15.9	0.0	0.7	1.0	0.0
WICAA	834	86.7	8.8	2.2	0.0	0.7	1.0	0.7
WVHS	435	87.1	0.7	8.5	0.0	0.2	1.6	1.8
Unknown Local Agencies	307	68.7	17.3	9.1	0.0	2.0	1.0	2.0
<b>Indiana</b>	<b>46,232</b>	<b>64.1</b>	<b>18.3</b>	<b>13.4</b>	<b>0.1</b>	<b>1.9</b>	<b>1.4</b>	<b>0.8</b>
<b>Nation (Prior Year)</b>	<b>1,162,961</b>	<b>41.6</b>	<b>22.7</b>	<b>29.2</b>	<b>1.1</b>	<b>3.4</b>	<b>1.3</b>	<b>0.7</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Age Distribution by Local Agency**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
BCA	202	0.0	5.0	15.3	67.3	11.4	1.0	0.0
BCHD	526	0.0	9.5	16.3	57.6	14.8	1.7	0.0
BRSI	663	0.2	3.8	15.5	64.1	15.5	0.6	0.3
CAP	594	0.0	6.2	15.5	62.5	14.8	1.0	0.0
CCHD	243	0.4	7.4	16.0	64.2	10.7	1.2	0.0
CFS	1,039	0.3	5.1	16.2	65.0	12.6	0.6	0.3
DCH	577	0.3	5.0	13.3	64.3	15.8	1.0	0.2
DCHD	252	0.0	2.8	9.9	64.7	20.2	2.4	0.0
DMH	285	0.0	9.8	15.4	58.9	14.4	1.4	0.0
DMHC	1,048	0.1	5.3	15.4	64.0	14.6	0.6	0.0
ECHD	568	1.1	5.5	12.9	57.4	21.1	1.9	0.2
ELKCHD	1,943	0.4	4.4	13.6	59.5	20.7	1.4	0.0
FHHC	75	*	*	*	*	*	*	*
FLCHD	476	0.4	6.3	16.2	62.4	13.9	0.6	0.2
FULCHD	143	0.0	3.5	12.6	71.3	11.2	1.4	0.0
GCRC	298	0.0	4.7	18.1	64.1	12.4	0.3	0.3
HCHF	421	0.0	4.3	14.5	60.1	19.5	1.7	0.0
HCMH&HS	296	0.0	5.1	13.5	61.5	18.9	1.0	0.0
HIS	582	0.2	7.6	13.9	63.7	13.4	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Age Distribution by Local Agency**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
HU	57	*	*	*	*	*	*	*
ICAP	300	0.3	6.0	14.0	65.3	13.3	0.7	0.3
IHC	2,665	0.5	4.8	14.3	62.5	16.5	1.1	0.3
JEFCHD	224	0.4	7.1	14.7	62.1	14.7	0.9	0.0
JNHC	338	0.3	4.7	15.7	63.3	15.4	0.6	0.0
JOCHD	730	0.1	5.1	15.1	63.0	14.8	1.9	0.0
KCHD	441	0.5	3.9	15.2	59.0	20.0	1.6	0.0
LCHD	145	0.7	6.2	16.6	56.6	17.9	0.7	1.4
MADCHD	957	0.3	6.3	15.3	61.4	15.3	1.4	0.1
MARCHD	9,411	0.4	4.7	10.7	62.7	20.1	1.3	0.1
MARSCHD	373	0.3	3.8	12.1	59.8	22.5	1.3	0.3
MCMH	423	0.0	5.0	11.3	67.4	15.1	0.9	0.2
MHS	1,958	0.1	6.7	13.1	58.1	20.9	1.1	0.0
MW	419	0.2	5.3	15.3	64.0	13.8	0.7	0.7
NCCAA	878	0.3	3.9	13.7	61.4	19.1	1.6	0.0
NHC	3,361	0.2	4.1	12.9	62.7	18.7	1.2	0.3
NHS	1,301	0.2	5.8	14.1	63.6	15.4	0.8	0.1
NWFS	4,723	0.3	5.1	12.0	58.7	22.3	1.5	0.2
ODCS	848	0.1	5.1	13.3	65.3	15.3	0.8	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Age Distribution by Local Agency**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
PHNA	821	0.1	3.4	13.8	63.8	17.4	1.0	0.5
St Mary's	549	0.2	7.7	15.1	63.2	12.6	0.9	0.4
STARCHD	171	1.2	2.9	10.5	64.9	18.1	1.8	0.6
STEUBCHD	275	0.0	5.5	19.3	63.6	11.3	0.4	0.0
SVJH	284	0.4	7.0	17.6	62.7	12.0	0.4	0.0
TCHD	1,273	0.1	4.5	12.8	60.5	20.5	1.6	0.1
VCHD	1,021	0.2	6.8	12.5	64.9	14.7	0.8	0.1
WCHD	184	0.0	3.3	11.4	68.5	15.8	1.1	0.0
WCMH	295	0.3	5.4	14.9	60.0	18.0	1.4	0.0
WICAA	834	0.2	7.8	17.4	62.5	11.5	0.6	0.0
WVHS	435	0.2	3.4	14.9	66.4	13.8	0.0	1.1
Unknown Local Agencies	307	0.3	6.5	11.7	65.8	14.3	1.3	0.0
<b>Indiana</b>	<b>46,232</b>	<b>0.3</b>	<b>5.2</b>	<b>13.2</b>	<b>62.1</b>	<b>17.9</b>	<b>1.2</b>	<b>0.1</b>
<b>Nation (Prior Year)</b>	<b>1,162,961</b>	<b>0.3</b>	<b>5.3</b>	<b>11.5</b>	<b>59.4</b>	<b>21.7</b>	<b>1.7</b>	<b>0.2</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Health Indicators by Local Agency**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Local Agency	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
BCA	158	5.1 (18)	50.0 (6)	155	14.8 (3)	54.2 (28)	30	* *	36	* *	198	2.0 (9)	198	0.5(2)
BCHD	424	7.3 (47)	53.8 (24)	412	18.9 (28)	54.4 (30)	69	* *	70	* *	516	4.7 (42)	516	7.4(42)
BRSI	531	4.5 (11)	56.5 (35)	506	21.1 (42)	48.2 (2)	69	* *	97	* *	643	5.0 (44)	643	7.8(43)
CAP	476	5.3 (23)	49.8 (5)	456	20.2 (36)	53.7 (23)	89	* *	112	22.3 (7)	573	4.0 (32)	573	4.0(32)
CCHD	207	6.3 (36)	57.5 (41)	199	22.1 (44)	48.2 (2)	28	* *	53	* *	237	3.4 (28)	237	2.5(19)
CFS	833	6.7 (40)	53.2 (19)	797	22.6 (46)	50.2 (7)	173	30.6 (3)	176	21.6 (6)	1,016	2.0 (9)	1,016	3.8(31)
DCH	428	5.4 (26)	53.0 (17)	408	15.4 (4)	58.3 (45)	74	* *	126	24.6(10)	558	2.9 (21)	558	3.4(27)
DCHD	191	5.8 (31)	60.7 (46)	185	16.8 (12)	51.9 (15)	41	* *	36	* *	249	4.4 (40)	249	2.8(20)
DMH	237	6.8 (42)	54.9 (28)	233	16.3 (8)	53.2 (20)	33	* *	51	* *	283	5.3 (45)	283	8.1(45)
DMHC	840	5.6 (28)	55.4 (30)	804	21.4 (43)	51.0 (10)	119	40.3 (13)	160	32.5(18)	1,020	4.3 (38)	1,020	3.1(23)
ECHD	444	2.7 (3)	63.5 (48)	428	22.4 (45)	51.4 (13)	86	* *	57	* *	558	3.4 (28)	558	5.2(38)
ELKCHD	1,656	4.0 (5)	57.5 (41)	1,599	19.9 (32)	51.2 (12)	191	27.2 (1)	247	17.4 (2)	1,893	2.7 (16)	1,893	2.1(16)
FHHC	59	* *	* *	58	* *	* *	8	* *	19	* *	75	* *	75	* *
FLCHD	377	5.0 (17)	53.6 (23)	361	20.2 (36)	52.9 (18)	74	* *	77	* *	461	6.3 (46)	461	9.8(47)
FULCHD	117	5.1 (18)	60.7 (46)	104	16.3 (8)	61.5 (47)	16	* *	36	* *	131	0.0 (1)	131	0.8(6)
GCRC	225	4.4 (9)	51.1 (10)	207	18.4 (21)	56.5 (37)	50	* *	49	* *	280	1.1 (4)	280	0.7(3)
HCHF	310	3.2 (4)	52.9 (16)	261	11.5 (1)	58.2 (44)	88	* *	217	47.9(24)	373	2.7 (16)	373	1.6(11)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Health Indicators by Local Agency**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
	(Rank)	(Rank)		(Rank)	(Rank)		(Rank)		(Rank)		(Rank)		(Rank)	
HCMH&HS	226	4.9 (15)	53.1 (18)	216	15.7 (6)	53.2 (20)	54	* *	68	* *	288	3.1 (25)	288	1.0(8)
HIS	457	6.3 (36)	50.8 (8)	442	17.0 (17)	49.3 (5)	81	* *	140	25.0(11)	569	4.0 (32)	569	4.4(34)
HU	47	* *	* *	46	* *	* *	4	* *	1	* *	56	* *	56	* *
ICAP	253	7.1 (45)	48.2 (1)	249	16.9 (14)	56.6 (38)	39	* *	26	* *	297	3.0 (22)	297	0.3(1)
IHC	2,139	5.1 (18)	52.6 (15)	2,051	18.6 (26)	51.1 (11)	420	34.0 (5)	601	25.6(12)	2,590	3.1 (25)	2,590	3.3(25)
JEFCHD	171	7.0 (44)	55.0 (29)	162	18.5 (24)	51.9 (15)	27	* *	36	* *	219	4.1 (35)	219	5.5(39)
JNHC	287	4.2 (7)	48.4 (2)	280	17.9 (19)	56.1 (35)	29	* *	72	* *	332	10.8 (47)	332	8.7(46)
JOCHD	526	6.8 (42)	50.8 (8)	506	15.6 (5)	56.9 (40)	133	42.1 (16)	144	19.4 (4)	712	3.8 (30)	712	8.0(44)
KCHD	342	5.6 (28)	53.5 (22)	325	20.6 (40)	48.3 (4)	75	* *	106	16.0 (1)	425	4.2 (37)	425	2.8(20)
LCHD	111	7.2 (46)	51.4 (11)	106	25.5 (47)	50.0 (6)	26	* *	35	* *	139	4.3 (38)	139	4.3(33)
MADCHD	735	4.5 (11)	56.5 (35)	702	18.4 (21)	52.4 (17)	152	48.7 (17)	165	36.4(22)	931	1.5 (5)	931	1.7(12)
MARCHD	6,641	4.4 (9)	58.2 (43)	6,156	20.2 (36)	51.6 (14)	1,384	36.9 (9)	1,391	26.2(14)	8,973	4.5 (41)	8,973	5.1(37)
MARSCHD	296	2.4 (1)	60.1 (45)	279	15.8 (7)	54.5 (31)	60	* *	63	* *	362	4.7 (42)	362	5.5(39)
MCMH	299	6.0 (34)	58.5 (44)	279	13.6 (2)	58.1 (43)	60	* *	65	* *	407	2.0 (9)	407	1.7(12)
MHS	1,561	4.9 (15)	55.5 (31)	1,489	20.3 (39)	50.2 (7)	290	39.0 (12)	417	32.9(20)	1,895	1.7 (7)	1,895	2.2(17)
MW	348	6.6 (39)	51.7 (13)	332	16.6 (11)	53.6 (22)	61	* *	243	23.5 (8)	405	3.2 (27)	405	6.7(41)
NCCAA	696	5.9 (33)	56.2 (32)	650	16.9 (14)	53.7 (23)	147	35.4 (7)	167	25.7(13)	839	2.4 (13)	839	1.7(12)

- (1) Reporting period is January 1 through December 31.
  - (2) Excludes records with unknown data or errors.
  - (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
  - (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
  - (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
  - (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 5F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Health Indicators by Local Agency**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	<b>Prepregnancy BMI</b> (4)			<b>Weight Gain</b> (5)			<b>Anemia</b> (Low Hb/Hct) (6)				<b>Gestational Diabetes</b> During Pregnancy		<b>Hypertension</b> During Pregnancy	
	<b>Under- weight</b>	<b>Over- weight</b>		<b>&lt; Ideal</b>	<b>&gt; Ideal</b>		<b>3rd Trimester</b>		<b>Postpartum</b>					
							<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
NHC	2,744	4.5 (11)	54.4 (26)	2,613	21.0 (41)	50.8 (9)	472	40.9 (15)	477	32.5 (18)	3,261	2.2 (12)	3,261	3.5 (28)
NHS	1,056	6.5 (38)	57.1 (39)	1,022	19.5 (30)	54.6 (32)	173	35.3 (6)	212	29.2 (16)	1,273	4.1 (35)	1,273	3.0 (22)
NWFS	3,409	4.3 (8)	56.9 (38)	3,096	17.5 (18)	55.7 (33)	737	37.6 (11)	714	28.3 (15)	4,428	2.5 (14)	4,428	3.5 (28)
ODCS	670	5.5 (27)	56.4 (33)	640	19.1 (29)	56.6 (38)	116	40.5 (14)	113	38.1 (23)	827	2.8 (18)	827	3.3 (25)
PHNA	655	5.2 (22)	53.9 (25)	607	18.8 (27)	54.2 (28)	129	30.2 (2)	189	33.9 (21)	776	3.0 (22)	776	0.8 (6)
St Mary's	445	4.0 (5)	51.5 (12)	422	18.5 (24)	53.1 (19)	83	* *	72	* *	531	1.9 (8)	531	4.9 (36)
STARCHD	150	6.7 (40)	49.3 (4)	146	19.9 (32)	56.2 (36)	20	* *	15	* *	167	3.0 (22)	167	1.2 (9)
STEUBCHD	234	5.1 (18)	56.4 (33)	227	19.8 (31)	53.7 (23)	26	* *	31	* *	269	0.7 (2)	269	0.7 (3)
SVJH	243	5.8 (31)	54.7 (27)	229	18.3 (20)	53.7 (23)	21	* *	48	* *	274	2.6 (15)	274	2.2 (17)
TCHD	1,045	4.6 (14)	52.2 (14)	984	20.1 (35)	48.0 (1)	172	37.2 (10)	156	18.6 (3)	1,227	2.8 (18)	1,227	1.3 (10)
VCHD	868	5.3 (23)	57.3 (40)	757	18.4 (21)	58.9 (46)	175	33.7 (4)	421	21.1 (5)	917	2.8 (18)	917	3.2 (24)
WCHD	150	6.0 (34)	53.3 (20)	142	16.9 (14)	57.0 (41)	28	* *	47	* *	178	3.9 (31)	178	1.7 (12)
WCMH	243	2.5 (2)	50.2 (7)	236	19.9 (32)	53.8 (27)	27	* *	42	* *	289	1.0 (3)	289	0.7 (3)
WICAA	696	5.3 (23)	53.3 (20)	608	16.8 (12)	57.4 (42)	122	36.1 (8)	393	23.7 (9)	753	1.6 (6)	753	3.5 (28)
WVHS	358	5.6 (28)	56.7 (37)	321	16.5 (10)	55.8 (34)	63	* *	236	29.7 (17)	402	4.0 (32)	402	4.7 (35)
Unknown Local Agencies	295	8.8 (48)	48.5 (3)	1	* *	* *	78	* *	2	* *	3	* *	3	* *
<b>Indiana</b>	<b>35,909</b>	<b>5.0</b>	<b>55.4</b>	<b>33,494</b>	<b>19.1</b>	<b>52.8</b>	<b>6,722</b>	<b>36.2</b>	<b>8,527</b>	<b>26.8</b>	<b>44,078</b>	<b>3.3</b>	<b>44,078</b>	<b>3.7</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Health Indicators by Local Agency**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>																									
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy												
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum																
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%											
												(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)	(Rank)				
<b>Nation (Prior Year)</b>												<b>1,106,370</b>	<b>4.5</b>	<b>53.4</b>	<b>835,927</b>	<b>21.5</b>	<b>48.0</b>	<b>164,201</b>	<b>33.9</b>	<b>346,981</b>	<b>28.1</b>	<b>419,956</b>	<b>5.2</b>	<b>467,675</b>	<b>6.3</b>

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Behavioral Indicators by Local Agency**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>									
Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prepregnancy <sup>(4)</sup>		During Pregnancy <sup>(5)</sup>	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
BCA	122	58.2 (31)	37.7 (32)	159	49.1 (15)	159	6.3 (48)	159	90.6 (8)
BCHD	354	44.6 (43)	45.2 (41)	425	44.5 (25)	425	18.6 (17)	426	90.1 (16)
BRSI	393	46.1 (42)	44.3 (39)	527	51.6 (12)	531	20.7 (9)	532	88.7 (22)
CAP	382	53.4 (35)	41.6 (35)	473	44.6 (23)	476	15.1 (35)	476	90.3 (11)
CCHD	141	53.2 (36)	41.1 (34)	207	54.6 (6)	206	18.9 (15)	207	89.9 (18)
CFS	623	60.8 (26)	35.0 (26)	837	49.3 (14)	833	17.6 (20)	836	90.6 (8)
DCH	279	70.3 (11)	24.7 (16)	433	38.1 (39)	431	18.3 (18)	431	87.7 (30)
DCHD	124	75.0 (5)	21.0 (6)	193	34.2 (45)	191	22.5 (3)	191	86.9 (33)
DMH	177	59.3 (30)	36.2 (29)	234	54.7 (5)	237	10.1 (46)	238	81.9 (41)
DMHC	648	49.1 (39)	43.7 (37)	834	52.2 (10)	775	13.7 (39)	822	79.0 (47)
ECHD	319	68.7 (16)	24.5 (15)	447	34.9 (44)	310	16.5 (27)	428	87.6 (31)
ELKCHD	1,424	40.2 (44)	56.1 (44)	1,654	57.3 (3)	1,658	21.2 (6)	1,658	81.4 (43)
FHHC	40	* *	* *	59	* *	59	* *	58	* *
FLCHD	267	65.9 (19)	25.8 (18)	385	40.3 (37)	378	20.4 (11)	378	89.2 (19)
FULCHD	74	* *	* *	120	46.7 (18)	117	15.4 (34)	117	88.0 (29)
GCRC	150	77.3 (3)	22.0 (10)	230	42.6 (31)	226	19.5 (13)	226	92.0 (5)
HCHF	194	82.0 (1)	11.3 (1)	314	29.0 (47)	306	22.5 (3)	311	92.6 (2)
HCMH&HS	181	64.1 (22)	31.5 (24)	226	42.5 (32)	228	20.6 (10)	228	90.8 (7)
HIS	331	63.1 (24)	31.4 (23)	458	43.0 (30)	458	13.1 (41)	458	85.4 (38)
HU	32	* *	* *	45	* *	47	* *	47	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Behavioral Indicators by Local Agency**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>											
<b>Local Agency</b>	<b>Medical Care</b>				<b>WIC Enrollment</b>			<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>		<b>1st Trimester</b>			<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
ICAP	215	61.9 (25)	36.3 (30)		254	52.4 (9)		254	15.0 (36)	254	81.5 (42)
IHC	1,539	65.6 (20)	28.5 (20)		2,150	42.0 (33)		2,141	15.6 (33)	2,144	89.0 (21)
JEFCHD	141	51.8 (37)	44.0 (38)		176	54.0 (7)		172	16.9 (26)	172	83.7 (40)
JNHC	217	47.9 (40)	49.3 (42)		285	53.3 (8)		287	22.0 (5)	286	80.4 (44)
JOCHD	418	63.4 (23)	29.4 (21)		532	33.3 (46)		508	16.5 (27)	526	90.3 (11)
KCHD	248	76.2 (4)	16.9 (2)		340	45.0 (22)		343	15.7 (32)	342	86.5 (34)
LCHD	76	* *	* *		114	43.9 (27)		111	18.9 (15)	111	86.5 (34)
MADCHD	563	64.8 (21)	23.4 (12)		739	41.1 (34)		737	14.9 (38)	737	90.1 (16)
MARCHD	4,976	67.1 (17)	24.4 (14)		6,648	35.7 (42)		6,605	11.7 (44)	6,659	87.5 (32)
MARSCHD	241	60.6 (27)	35.3 (28)		296	40.2 (38)		296	19.3 (14)	297	88.2 (28)
MCMH	224	69.2 (15)	24.1 (13)		303	35.0 (43)		300	21.0 (7)	300	90.3 (11)
MHS	1,214	69.4 (13)	25.1 (17)		1,577	40.5 (36)		1,563	17.1 (23)	1,563	92.5 (3)
MW	238	69.3 (14)	27.7 (19)		351	49.0 (16)		348	17.2 (22)	348	90.5 (10)
NCCAA	552	56.2 (33)	36.8 (31)		697	40.9 (35)		665	20.2 (12)	688	93.9 (1)
NHC	2,083	60.0 (29)	35.0 (26)		2,739	46.2 (19)		2,747	17.3 (21)	2,747	89.2 (19)
NHS	812	66.1 (18)	29.6 (22)		1,064	45.4 (21)		1,058	17.1 (23)	1,059	90.2 (15)
NWFS	2,244	69.8 (12)	21.8 (9)		3,428	37.1 (40)		3,408	16.4 (29)	3,413	88.3 (26)
ODCS	462	74.5 (6)	21.4 (8)		673	47.5 (17)		667	18.0 (19)	671	90.3 (11)
PHNA	420	71.7 (10)	23.3 (11)		653	44.6 (23)		480	20.8 (8)	635	92.4 (4)
St Mary's	326	60.4 (28)	34.4 (25)		448	44.0 (26)		446	9.2 (47)	447	88.6 (23)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Behavioral Indicators by Local Agency**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	<b>Medical Care</b>			<b>WIC Enrollment</b>			<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>	<b>1st Trimester</b>		<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)		
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	
STARCHD	126	54.0 (34)	44.4 (40)	149	55.0 (4)	150	16.0 (30)	150	79.3 (46)	
STEUBCHD	177	46.3 (41)	52.0 (43)	234	62.4 (2)	234	15.0 (36)	234	79.9 (45)	
SVJH	195	29.2 (45)	67.7 (45)	239	67.4 (1)	243	16.0 (30)	243	74.9 (48)	
TCHD	833	50.8 (38)	42.7 (36)	1,037	51.7 (11)	982	23.1 (2)	1,034	86.0 (37)	
VCHD	577	71.9 (9)	20.8 (5)	883	36.4 (41)	869	13.7 (39)	868	90.9 (6)	
WCHD	112	58.0 (32)	40.2 (33)	149	50.3 (13)	150	28.0 (1)	149	88.6 (23)	
WCMH	176	74.4 (7)	21.0 (6)	244	45.9 (20)	244	13.1 (41)	244	86.1 (36)	
WICAA	386	73.1 (8)	19.9 (4)	697	43.3 (29)	697	12.6 (43)	699	88.3 (26)	
WVHS	234	77.8 (2)	18.4 (3)	364	43.4 (28)	361	10.5 (45)	361	85.3 (39)	
Unknown Local Agencies	1	*	*	305	16.7 (48)	294	17.0 (25)	296	88.5 (25)	
<b>Indiana</b>	<b>26,281</b>	<b>62.8</b>	<b>30.8</b>	<b>36,028</b>	<b>42.6</b>	<b>35,411</b>	<b>16.1</b>	<b>35,904</b>	<b>88.0</b>	
<b>Nation (Prior Year)</b>	<b>429,770</b>	<b>83.1</b>	<b>4.1</b>	<b>1,074,863</b>	<b>36.0</b>	<b>380,154</b>	<b>19.2</b>	<b>418,448</b>	<b>85.0</b>	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7F**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Indiana**  
**Comparison of Smoking Indicators by Local Agency**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
BCA	202	41.1 (41)	198	33.3 (44)	81	* *	159	51.6 (35)	195	51.8 (44)
BCHD	526	37.5 (26)	516	26.6 (32)	192	30.2 (26)	426	54.7 (41)	510	50.4 (42)
BRSI	663	40.1 (39)	643	29.2 (38)	253	30.4 (24)	532	45.1 (27)	632	30.5 (15)
CAP	594	39.7 (37)	573	28.1 (36)	228	35.1 (8)	472	58.5 (46)	570	48.9 (40)
CCHD	243	37.9 (28)	237	26.2 (28)	89	* *	207	59.4 (47)	234	57.7 (46)
CFS	1,039	37.2 (25)	1,016	26.2 (28)	376	30.9 (20)	836	48.3 (31)	996	43.0 (28)
DCH	577	40.9 (40)	558	31.0 (41)	223	36.3 (6)	430	51.6 (35)	541	47.3 (35)
DCHD	252	27.0 (8)	249	21.3 (16)	66	* *	186	43.0 (20)	246	40.7 (24)
DMH	285	41.4 (42)	283	29.0 (37)	117	35.0 (9)	238	45.0 (26)	280	42.9 (27)
DMHC	1,048	38.3 (30)	1,020	26.9 (33)	385	30.4 (24)	841	54.8 (42)	1,005	44.8 (31)
ECHD	568	14.8 (1)	558	5.9 (1)	83	* *	445	21.1 (3)	549	13.1 (4)
ELKCHD	1,943	24.2 (5)	1,893	14.7 (3)	452	32.1 (16)	1,654	19.3 (2)	1,875	8.0 (3)
FHHC	75	* *	75	* *	33	* *	58	* *	75	* *
FLCHD	475	38.3 (30)	461	26.0 (25)	174	37.9 (4)	377	43.5 (22)	456	41.9 (25)
FULCHD	143	36.4 (24)	131	26.0 (25)	49	* *	117	47.9 (30)	131	42.0 (26)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7F**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Indiana**  
**Comparison of Smoking Indicators by Local Agency**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
GCRC	298	44.0 (45)	280	36.1 (46)	120	24.2 (32)	226	59.7 (48)	279	54.8 (45)
HCHF	421	27.6 (9)	373	15.3 (6)	94	* *	311	39.9 (17)	368	38.9 (21)
HCMH&HS	296	29.4 (15)	288	19.1 (8)	85	* *	226	24.8 (6)	286	14.0 (5)
HIS	582	37.8 (27)	569	26.4 (30)	209	32.5 (14)	459	53.6 (38)	562	51.1 (43)
HU	57	* *	56	* *	25	* *	47	* *	55	* *
ICAP	300	47.7 (48)	297	33.7 (45)	140	26.4 (30)	253	53.4 (37)	294	48.0 (36)
IHC	2,665	29.2 (14)	2,590	21.1 (14)	746	30.8 (21)	2,143	34.9 (11)	2,552	21.6 (9)
JEFCHD	224	44.6 (46)	219	36.5 (47)	96	* *	172	58.1 (45)	213	43.7 (29)
JNHC	338	46.4 (47)	332	32.2 (43)	155	32.3 (15)	287	53.7 (39)	330	49.7 (41)
JOCHD	730	27.8 (11)	712	21.2 (15)	195	32.8 (12)	527	42.7 (19)	699	29.9 (14)
KCHD	441	26.1 (6)	425	20.9 (12)	110	25.5 (31)	343	38.5 (13)	420	23.3 (10)
LCHD	144	26.4 (7)	139	20.9 (12)	35	* *	111	43.2 (21)	134	39.6 (23)
MADCHD	957	36.3 (23)	931	27.1 (35)	331	30.5 (22)	737	51.3 (34)	922	46.3 (34)
MARCHD	9,411	18.7 (2)	8,973	14.8 (4)	1,621	37.1 (5)	6,242	29.5 (7)	8,685	28.6 (13)
MARSCHD	373	28.7 (13)	362	19.9 (11)	100	34.0 (10)	296	17.6 (1)	361	5.8 (1)

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7F**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Indiana**  
**Comparison of Smoking Indicators by Local Agency**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
MCMH	423	34.5 (19)	407	30.2 (40)	136	29.4 (28)	298	51.0 (33)	403	48.4 (37)
MHS	1,958	27.7 (10)	1,895	14.9 (5)	516	43.8 (3)	1,560	24.7 (5)	1,868	16.0 (8)
MW	419	41.8 (43)	405	26.4 (30)	168	36.3 (6)	348	43.7 (24)	403	32.5 (17)
NCCAA	878	32.8 (17)	839	24.7 (21)	267	33.0 (11)	694	43.5 (22)	824	38.8 (20)
NHC	3,361	28.5 (12)	3,261	19.3 (9)	915	29.7 (27)	2,740	39.4 (16)	3,215	34.8 (18)
NHS	1,301	34.6 (20)	1,273	24.7 (21)	432	32.6 (13)	1,059	41.8 (18)	1,264	38.0 (19)
NWFS	4,722	22.3 (3)	4,428	14.2 (2)	970	45.1 (2)	3,407	32.6 (10)	4,374	27.0 (11)
ODCS	848	33.8 (18)	827	25.6 (23)	280	23.6 (34)	670	30.4 (9)	812	14.2 (6)
PHNA	820	34.8 (21)	776	23.6 (20)	262	30.5 (22)	655	39.1 (15)	771	28.0 (12)
St Mary's	549	43.0 (44)	531	29.4 (39)	228	29.4 (28)	446	56.1 (43)	520	59.2 (47)
STARCHD	171	35.1 (22)	167	26.9 (33)	59	* *	150	46.0 (28)	166	48.8 (39)
STEUBCHD	275	39.6 (36)	269	22.7 (19)	107	31.8 (18)	233	54.1 (40)	263	45.6 (32)
SVJH	284	39.8 (38)	274	31.4 (42)	108	18.5 (35)	243	56.8 (44)	271	48.7 (38)
TCHD	1,273	23.3 (4)	1,227	18.2 (7)	283	23.7 (33)	1,046	35.1 (12)	1,210	30.7 (16)
VCHD	1,021	39.0 (33)	917	19.3 (9)	347	45.8 (1)	868	22.4 (4)	906	7.9 (2)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 7F**  
**2011 Pregnancy Nutrition Surveillance** (1)  
**Indiana**  
**Comparison of Smoking Indicators by Local Agency**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
WCHD	184	38.0 (29)	178	21.3 (16)	68	* *	150	44.0 (25)	177	39.0 (22)
WCMH	295	30.5 (16)	289	22.1 (18)	89	* *	244	29.9 (8)	287	15.3 (7)
WICAA	834	39.0 (33)	753	25.9 (24)	285	31.2 (19)	699	50.2 (32)	745	44.2 (30)
WVHS	435	38.9 (32)	402	26.1 (27)	156	32.1 (16)	360	47.5 (29)	394	45.7 (33)
Unknown Local Agencies	306	39.2 (35)	3	* *	1	* *	291	38.5 (13)	3	* *
<b>Indiana</b>	<b>46,227</b>	<b>29.0</b>	<b>44,078</b>	<b>20.3</b>	<b>12,540</b>	<b>33.8</b>	<b>35,519</b>	<b>37.4</b>	<b>43,331</b>	<b>31.1</b>
<b>Nation (Prior Year)</b>	<b>814,631</b>	<b>23.0</b>	<b>715,034</b>	<b>12.1</b>	<b>137,883</b>	<b>38.1</b>	<b>660,570</b>	<b>16.3</b>	<b>562,126</b>	<b>15.6</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
BCA	189	5.3 (4)	6.9 (27)		151	13.9 (32)	130	2.3 (16)	189	56.1 (46)
BCHD	498	6.4 (9)	6.6 (22)		408	12.3 (20)	358	2.5 (21)	498	70.9 (18)
BRSI	609	6.2 (7)	5.9 (14)		491	12.4 (22)	430	2.1 (14)	609	74.2 (12)
CAP	549	7.5 (18)	8.7 (44)		441	11.8 (14)	388	2.8 (28)	550	68.0 (26)
CCHD	225	5.3 (4)	4.9 (4)		192	12.0 (17)	169	1.8 (7)	225	65.3 (33)
CFS	971	7.8 (24)	4.8 (3)		799	12.5 (24)	698	2.7 (26)	973	70.4 (21)
DCH	526	8.9 (37)	8.9 (45)		408	12.5 (24)	354	2.3 (16)	532	62.2 (38)
DCHD	242	4.5 (3)	5.8 (10)		187	12.8 (26)	163	0.6 (3)	242	76.9 (6)
DMH	274	9.9 (43)	6.6 (22)		229	9.6 (4)	207	4.3 (43)	274	70.4 (21)
DMHC	961	6.2 (7)	8.3 (41)		779	11.4 (13)	687	2.0 (12)	964	59.0 (44)
ECHD	535	10.7 (45)	5.8 (10)		428	16.8 (44)	355	2.5 (21)	537	64.4 (36)
ELKCHD	1,784	7.3 (17)	8.3 (41)		1,556	11.1 (9)	1,377	2.7 (26)	1,795	74.8 (11)
FHHC	75	* * *	* *		59	* *	51	* *	75	* *
FLCHD	447	9.4 (41)	5.8 (10)		367	19.3 (46)	293	2.4 (19)	450	65.6 (32)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Birthweight				Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)							
	Number	% (Rank)	%	(Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
FULCHD	129	3.1 (1)	7.8 (39)		107	9.3 (3)	97	* *	129	72.9 (15)
GCRC	275	8.4 (30)	5.8 (10)		212	14.6 (33)	180	2.2 (15)	278	55.0 (47)
HCHF	364	6.6 (11)	6.6 (22)		264	9.8 (5)	238	1.3 (5)	364	76.4 (9)
HCMH&HS	283	6.4 (9)	7.4 (31)		218	11.9 (16)	192	0.5 (1)	283	76.7 (8)
HIS	549	8.4 (30)	5.6 (8)		437	15.1 (39)	369	3.8 (40)	551	70.8 (19)
HU	55	* *	* *		44	* *	34	* *	55	* *
ICAP	288	6.9 (14)	7.6 (35)		248	6.5 (1)	230	4.3 (43)	290	70.0 (23)
IHC	2,496	7.6 (20)	6.2 (17)		2,041	11.2 (10)	1,803	3.0 (30)	2,507	62.2 (38)
JEFCHD	201	11.4 (46)	7.5 (32)		165	18.8 (45)	130	3.1 (31)	205	72.2 (16)
JNHC	321	6.9 (14)	6.9 (27)		274	12.0 (17)	240	3.3 (34)	322	68.6 (25)
JOCHD	693	7.6 (20)	7.6 (35)		509	13.2 (30)	440	2.3 (16)	695	71.1 (17)
KCHD	410	6.6 (11)	6.6 (22)		317	12.3 (20)	278	0.7 (4)	411	73.5 (14)
LCHD	130	9.2 (38)	6.9 (27)		104	15.4 (42)	88	* *	130	78.5 (3)
MADCHD	906	8.4 (30)	5.4 (7)		706	15.2 (40)	592	1.9 (8)	914	58.8 (45)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
MARCHD	8,754	8.8 (35)	6.2 (17)	6,142	14.6 (33)	5,224	2.6 (24)	8,805	64.4 (36)
MARSCHD	354	4.2 (2)	7.6 (35)	281	8.5 (2)	257	1.9 (8)	354	77.4 (4)
MCMH	399	9.3 (40)	8.5 (43)	284	12.0 (17)	250	2.4 (19)	401	73.6 (13)
MHS	1,830	8.7 (33)	7.8 (39)	1,490	10.7 (6)	1,326	3.4 (36)	1,837	76.9 (6)
MW	386	6.7 (13)	6.2 (17)	331	11.8 (14)	291	1.4 (6)	387	65.1 (34)
NCCAA	806	8.7 (33)	5.7 (9)	644	14.6 (33)	550	3.5 (38)	807	64.9 (35)
NHC	3,133	8.8 (35)	4.9 (4)	2,581	14.6 (33)	2,198	3.6 (39)	3,139	67.5 (27)
NHS	1,216	7.6 (20)	6.5 (20)	1,006	14.7 (37)	851	3.2 (33)	1,223	62.1 (40)
NWFS	4,262	8.2 (28)	6.6 (22)	3,055	13.5 (31)	2,631	2.8 (28)	4,278	70.0 (23)
ODCS	802	9.7 (42)	4.4 (2)	641	12.9 (28)	558	4.5 (45)	802	60.3 (42)
PHNA	748	8.2 (28)	9.1 (46)	595	11.3 (12)	527	3.4 (36)	749	77.3 (5)
St Mary""s	512	10.4 (44)	6.1 (15)	422	16.1 (43)	351	2.0 (12)	515	59.6 (43)
STARCHD	161	8.1 (26)	7.5 (32)	141	12.8 (26)	123	4.1 (42)	161	70.8 (19)
STEUBCHD	253	7.5 (18)	11.5 (47)	224	11.2 (10)	196	3.1 (31)	256	83.6 (1)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Infant Health Indicators by Local Agency**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
SVJH	260	8.1 (26)	6.5 (20)	223	14.8 (38)	190	0.5 (1)	260	67.3 (28)
TCHD	1,170	7.1 (16)	7.5 (32)	966	10.7 (6)	860	3.3 (34)	1,174	75.0 (10)
VCHD	896	9.2 (38)	6.1 (15)	773	15.3 (41)	651	2.6 (24)	902	66.1 (31)
WCHD	169	11.8 (47)	5.3 (6)	139	23.0 (47)	107	1.9 (8)	169	81.1 (2)
WCMH	272	5.9 (6)	7.7 (38)	228	11.0 (8)	201	2.5 (21)	275	66.5 (29)
WICAA	724	8.0 (25)	6.9 (27)	607	12.4 (22)	525	1.9 (8)	732	66.3 (30)
WVHS	380	7.6 (20)	3.4 (1)	321	13.1 (29)	279	3.9 (41)	381	60.4 (41)
Unknown Local Agencies	2	*	*	0	*	0	*	2	*
<b>Indiana</b>	<b>42,474</b>	<b>8.2</b>	<b>6.5</b>	<b>33,235</b>	<b>13.2</b>	<b>28,717</b>	<b>2.8</b>	<b>42,656</b>	<b>67.6</b>
<b>Nation (Prior Year)</b>	<b>880,933</b>	<b>8.0</b>	<b>6.7</b>	<b>948,286</b>	<b>10.7</b>	<b>749,107</b>	<b>3.4</b>	<b>871,811</b>	<b>68.2</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Racial and Ethnic Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaska Native %</b>	<b>Asian/Pacific Islander %</b>	<b>Multiple Races %</b>	<b>All Other/Unknown %</b>
BCA	612	94.0	1.1	2.9	0.0	0.7	0.2	1.1
BCHD	1,632	80.3	1.8	15.3	0.0	0.6	1.1	0.9
BRSI	1,977	94.8	0.6	2.5	0.1	0.5	0.8	0.8
CAP	1,838	91.7	0.4	6.0	0.1	0.4	0.6	0.9
CCHD	703	96.4	0.6	0.7	0.0	0.4	1.1	0.7
CFS	3,187	92.5	0.4	4.9	0.1	0.2	0.7	1.2
DCH	1,744	95.0	0.6	1.5	0.3	0.1	0.7	1.6
DCHD	782	73.0	0.0	23.7	0.1	1.7	0.6	0.9
DMH	931	96.1	0.4	1.4	0.0	1.2	0.3	0.5
DMHC	3,268	88.9	3.6	4.1	0.2	0.5	1.9	0.7
ECHD	1,852	6.3	36.2	56.4	0.1	0.2	0.3	0.5
ELKCHD	6,108	54.7	10.0	32.5	0.0	0.9	1.2	0.8
FHHC	240	96.7	0.0	1.7	0.4	0.0	0.4	0.8
FLCHD	1,410	76.0	12.5	6.1	0.1	2.2	1.3	1.8
FULCHD	517	89.0	0.8	9.1	0.0	0.0	0.2	1.0
GCRC	871	92.2	3.0	2.2	0.5	0.6	1.1	0.5
HCHF	1,273	86.6	4.6	4.1	0.1	3.0	0.1	1.5
HCMH&HS	866	94.3	1.4	1.8	0.1	0.2	0.8	1.3
HIS	1,694	90.9	0.7	7.1	0.1	0.2	0.2	0.8
HU	173	97.7	0.0	1.2	0.0	0.0	0.0	1.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Racial and Ethnic Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>White, Not Hispanic %</b>	<b>Black, Not Hispanic %</b>	<b>Hispanic %</b>	<b>American Indian/Alaska Native %</b>	<b>Asian/Pacific Islander %</b>	<b>Multiple Races %</b>	<b>All Other/Unknown %</b>
ICAP	1,003	94.5	0.8	2.1	0.1	0.3	1.2	1.0
IHC	8,198	76.6	6.7	13.7	0.1	1.0	1.4	0.6
JEFCHD	733	87.2	1.8	7.8	0.0	0.5	1.5	1.2
JNHC	1,120	95.5	0.6	1.6	0.0	0.7	0.7	0.8
JOCHD	2,227	87.4	1.4	5.4	0.0	3.1	1.1	1.5
KCHD	1,278	75.4	0.8	21.8	0.1	0.4	0.5	1.2
LCHD	435	82.1	1.1	12.0	0.0	1.1	1.4	2.3
MADCHD	2,920	77.0	12.1	8.3	0.2	0.3	1.4	0.8
MARCHD	29,030	34.3	39.6	20.6	0.2	3.0	1.5	0.8
MARSCHD	1,064	74.5	0.5	22.7	0.1	0.5	1.0	0.8
MCMH	1,275	96.9	0.4	1.2	0.0	0.2	0.6	0.8
MHS	5,934	49.8	28.3	16.5	0.3	1.7	2.4	1.1
MW	1,304	90.8	1.2	2.9	0.3	0.7	3.5	0.6
NCCAA	2,735	68.2	18.6	9.9	0.0	0.7	1.9	0.7
NHC	10,518	57.0	22.0	13.7	0.1	4.7	1.9	0.6
NHS	4,061	78.4	6.0	12.5	0.1	0.9	1.3	0.9
NWFS	14,908	45.9	34.6	16.7	0.1	0.6	1.1	0.9
ODCS	2,621	82.0	12.0	2.5	0.2	0.7	2.2	0.5
PHNA	2,546	84.4	4.6	3.5	0.0	4.8	1.7	1.1
St Mary""s	1,674	94.9	1.0	2.0	0.0	0.2	0.8	1.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 3F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Racial and Ethnic Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Racial and Ethnic Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
STARCHD	566	94.5	0.2	4.6	0.0	0.2	0.0	0.5
STEUBCHD	840	91.0	0.6	4.9	0.0	1.3	0.4	1.9
SVJH	864	95.0	0.7	2.5	0.6	0.1	0.3	0.7
TCHD	3,948	61.5	10.2	19.1	0.4	6.4	1.2	1.1
VCHD	3,081	71.9	18.3	5.0	0.1	1.0	2.1	1.6
WCHD	605	97.4	0.3	1.2	0.0	0.3	0.5	0.3
WCMH	916	81.6	0.0	17.0	0.1	0.4	0.4	0.4
WICAA	2,526	88.3	7.3	1.9	0.0	0.8	1.1	0.6
WVHS	1,295	86.6	0.6	8.6	0.1	0.2	1.7	2.2
Unknown Local Agencies	308	68.5	17.2	9.1	0.0	1.9	1.0	2.3
<b>Indiana</b>	<b>142,211</b>	<b>63.8</b>	<b>18.2</b>	<b>13.9</b>	<b>0.1</b>	<b>1.8</b>	<b>1.3</b>	<b>0.9</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 4F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Age Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
BCA	612	0.2	4.2	14.4	65.7	14.2	1.3	0.0
BCHD	1,632	0.1	8.1	16.4	59.3	15.0	1.1	0.0
BRSI	1,977	0.2	5.6	16.1	63.8	13.8	0.5	0.2
CAP	1,838	0.2	6.1	16.8	60.9	15.1	1.0	0.0
CCHD	703	0.1	8.0	15.4	63.7	11.9	0.9	0.0
CFS	3,187	0.3	5.1	15.7	64.6	13.3	0.6	0.4
DCH	1,744	0.3	5.4	15.3	63.0	14.7	1.1	0.2
DCHD	782	0.3	4.2	11.6	62.4	19.8	1.5	0.1
DMH	931	0.2	9.0	17.0	61.0	11.6	1.2	0.0
DMHC	3,268	0.2	6.6	15.3	63.3	13.9	0.7	0.0
ECHD	1,852	0.8	6.8	12.6	55.9	22.2	1.6	0.1
ELKCHD	6,108	0.4	5.8	13.9	58.6	19.8	1.3	0.0
FHHC	240	0.0	8.8	18.8	58.3	13.3	0.8	0.0
FLCHD	1,410	0.4	6.3	16.2	61.8	14.6	0.4	0.3
FULCHD	517	0.2	4.8	19.3	60.3	14.3	1.0	0.0
GCRC	871	0.0	6.9	15.3	64.1	13.0	0.7	0.1
HCHF	1,273	0.1	5.2	14.5	59.8	19.1	1.3	0.1
HCMH&HS	866	0.1	5.4	15.5	62.1	16.1	0.8	0.0
HIS	1,694	0.1	7.0	15.6	61.7	14.3	1.2	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Age Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
HU	173	0.0	9.8	13.9	61.8	13.9	0.6	0.0
ICAP	1,003	0.6	6.0	15.5	64.2	13.2	0.5	0.1
IHC	8,198	0.4	5.5	14.2	61.9	16.7	1.0	0.3
JEFCHD	733	0.5	6.7	16.1	61.8	14.1	0.8	0.0
JNHC	1,120	0.4	6.2	16.3	64.5	11.7	1.0	0.0
JOCHD	2,227	0.1	5.9	14.8	61.9	16.1	1.2	0.0
KCHD	1,278	0.3	6.7	15.2	58.2	18.4	1.3	0.0
LCHD	435	0.2	6.0	13.6	58.6	19.1	1.8	0.7
MADCHD	2,920	0.4	5.9	15.0	63.6	14.1	0.9	0.1
MARCHD	29,030	0.3	5.2	12.2	61.6	19.5	1.2	0.1
MARSCHD	1,064	0.1	5.3	12.1	57.7	22.7	1.9	0.3
MCMH	1,275	0.2	5.8	14.6	64.5	13.9	0.9	0.2
MHS	5,934	0.2	6.3	13.6	59.1	19.5	1.3	0.0
MW	1,304	0.3	5.3	16.4	63.9	13.0	0.8	0.2
NCCAA	2,735	0.3	5.5	13.5	60.6	18.9	1.2	0.0
NHC	10,518	0.2	4.6	13.3	63.2	17.0	1.0	0.6
NHS	4,061	0.2	6.4	15.7	62.7	14.1	0.7	0.1
NWFS	14,908	0.3	5.5	12.9	59.5	20.3	1.3	0.1
ODCS	2,621	0.2	5.5	14.4	64.6	14.2	0.9	0.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 4F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Age Distribution by Local Agency**  
**Combined 3 Years**

<b>Comparison of Age Distribution</b> <sup>(2)</sup>								
<b>Local Agency</b>	<b>Number Included in Analysis</b>	<b>&lt; 15 Years %</b>	<b>15 - 17 Years %</b>	<b>18 - 19 Years %</b>	<b>20 - 29 Years %</b>	<b>30 - 39 Years %</b>	<b>&gt;= 40 Years %</b>	<b>Unknown %</b>
PHNA	2,546	0.1	3.7	14.6	63.4	16.6	1.2	0.5
St Mary's	1,674	0.2	5.7	15.7	63.1	14.8	0.4	0.1
STARCHD	566	0.9	5.3	14.3	61.5	16.6	1.2	0.2
STEUBCHD	840	0.1	5.1	16.2	64.4	13.5	0.7	0.0
SVJH	864	0.2	8.9	21.1	56.4	13.1	0.3	0.0
TCHD	3,948	0.2	5.6	13.1	59.7	20.0	1.2	0.2
VCHD	3,081	0.4	6.3	14.0	64.6	14.3	0.5	0.0
WCHD	605	0.3	4.3	15.4	65.8	12.9	1.3	0.0
WCMH	916	0.7	6.1	12.9	60.8	18.0	1.5	0.0
WICAA	2,526	0.5	8.1	16.7	61.5	12.4	0.8	0.0
WVHS	1,295	0.2	5.9	15.0	63.7	14.0	0.5	0.8
Unknown Local Agencies	308	0.3	6.5	11.7	65.6	14.6	1.3	0.0
<b>Indiana</b>	<b>142,211</b>	<b>0.3</b>	<b>5.7</b>	<b>14.0</b>	<b>61.6</b>	<b>17.3</b>	<b>1.1</b>	<b>0.1</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
BCA	464	6.7 (44)	51.7 (11)	438	17.4 (17)	51.6 (6)	85	* *	155	21.3(13)	586	1.0 (3)	586	1.2(5)
BCHD	1,291	7.5 (49)	52.9 (21)	1,240	16.9 (12)	55.5 (33)	259	32.4 (14)	250	26.4(25)	1,590	5.3 (45)	1,590	6.6(44)
BRSI	1,614	4.5 (9)	56.3 (43)	1,533	21.5 (49)	49.5 (1)	220	26.8 (4)	247	19.0 (7)	1,907	3.4 (30)	1,907	6.0(40)
CAP	1,493	5.4 (27)	51.8 (12)	1,425	18.5 (28)	56.6 (40)	288	32.3 (13)	383	23.2(21)	1,781	4.5 (40)	1,781	4.9(32)
CCHD	589	4.9 (18)	58.6 (47)	561	20.1 (43)	53.1 (18)	82	* *	123	18.7 (5)	684	2.8 (25)	684	4.1(27)
CFS	2,575	7.0 (48)	52.0 (15)	2,460	18.7 (33)	53.8 (26)	472	32.8 (15)	522	27.8(29)	3,094	2.2 (16)	3,094	5.4(35)
DCH	1,362	5.4 (27)	51.8 (12)	1,289	15.5 (4)	57.2 (44)	240	44.6 (36)	364	26.6(27)	1,674	2.6 (22)	1,674	4.6(31)
DCHD	589	5.1 (21)	61.1 (48)	570	17.7 (20)	53.5 (23)	135	26.7 (3)	133	17.3 (4)	767	5.0 (43)	767	3.1(19)
DMH	746	6.4 (41)	53.6 (26)	704	17.6 (19)	51.4 (4)	103	49.5 (38)	176	32.4(42)	901	4.9 (42)	901	7.7(46)
DMHC	2,655	5.8 (36)	54.6 (34)	2,528	20.8 (47)	52.8 (16)	409	38.6 (28)	568	33.1(43)	3,171	3.9 (36)	3,171	3.5(23)
ECHD	1,439	2.8 (1)	62.8 (50)	1,378	20.7 (46)	51.9 (10)	290	50.0 (39)	158	39.9(46)	1,805	4.2 (39)	1,805	6.9(45)
ELKCHD	5,123	3.9 (3)	57.4 (46)	4,883	19.7 (40)	51.8 (9)	623	27.1 (5)	969	20.6(10)	5,897	3.1 (27)	5,897	3.1(19)
FHHC	199	6.5 (42)	49.2 (2)	197	16.2 (7)	56.9 (43)	20	* *	37	* *	240	1.7 (9)	240	2.5(16)
FLCHD	1,093	5.3 (23)	53.8 (29)	1,019	19.4 (37)	52.3 (12)	201	34.8 (21)	409	25.4(24)	1,342	8.3 (49)	1,342	12.4(49)
FULCHD	423	4.3 (5)	52.2 (19)	390	18.5 (28)	58.2 (47)	71	* *	113	28.3(30)	491	0.0 (1)	491	0.2(1)
GCRC	663	5.3 (23)	54.0 (30)	614	17.8 (21)	55.5 (33)	133	31.6 (12)	158	20.9(12)	824	1.2 (6)	824	1.5(7)
HCHF	942	4.7 (12)	52.1 (16)	801	12.2 (2)	57.7 (46)	248	41.5 (35)	674	39.6(45)	1,136	2.4 (19)	1,136	2.5(16)

- (1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Health Indicators by Local Agency**  
**Combined 3 Years**

Comparison of Maternal Health Indicators <sup>(2, 3)</sup>														
Local Agency	Prepregnancy BMI <sup>(4)</sup>			Weight Gain <sup>(5)</sup>			Anemia (Low Hb/Hct) <sup>(6)</sup>				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under-weight	Over-weight		< Ideal	> Ideal		3rd Trimester		Postpartum					
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
HCMH&HS	651	5.7 (33)	52.1 (16)	614	14.7 (3)	59.1 (48)	149	28.9 (6)	193	16.1 (2)	834	3.5 (32)	834	2.4(14)
HIS	1,304	6.2 (40)	53.5 (25)	1,243	17.5 (18)	53.8 (26)	219	33.3 (17)	603	23.1(20)	1,636	3.4 (30)	1,636	3.7(26)
HU	143	4.2 (4)	51.0 (9)	137	11.7 (1)	62.0 (49)	21	* *	14	* *	166	1.2 (6)	166	5.4(35)
ICAP	828	6.8 (47)	50.7 (6)	797	18.2 (25)	57.5 (45)	127	31.5 (11)	127	18.9 (6)	979	1.8 (11)	979	0.9(4)
IHC	6,516	5.0 (19)	53.4 (24)	6,189	19.6 (39)	51.6 (6)	1,349	39.5 (30)	1,740	28.3(30)	7,913	2.7 (24)	7,913	4.1(27)
JEFCHD	581	5.7 (33)	54.2 (31)	547	16.3 (8)	52.3 (12)	100	31.0 (10)	126	11.1 (1)	704	5.4 (46)	704	6.3(42)
JNHC	939	5.3 (23)	53.2 (23)	895	16.9 (12)	56.1 (36)	131	30.5 (7)	236	25.0(23)	1,083	6.0 (48)	1,083	7.9(47)
JOCHD	1,628	6.1 (39)	51.1 (10)	1,546	16.5 (10)	56.0 (35)	377	35.5 (22)	424	20.8(11)	2,156	3.5 (32)	2,156	7.9(47)
KCHD	1,001	4.7 (12)	55.5 (39)	926	17.8 (21)	53.1 (18)	245	33.9 (18)	336	21.4(14)	1,205	4.5 (40)	1,205	5.0(33)
LCHD	337	5.3 (23)	50.1 (4)	316	18.4 (26)	55.1 (30)	74	* *	122	23.0(18)	412	4.1 (38)	412	6.1(41)
MADCHD	2,169	4.7 (12)	55.4 (36)	2,076	17.3 (15)	53.1 (18)	529	45.6 (37)	587	34.1(44)	2,843	1.9 (12)	2,843	2.5(16)
MARCHD	20,240	4.4 (7)	56.7 (45)	18,550	19.1 (35)	53.3 (22)	4,409	38.1 (27)	4,342	27.6(28)	27,472	3.9 (36)	27,472	5.3(34)
MARSCHD	856	3.2 (2)	61.1 (48)	814	16.3 (8)	56.3 (39)	187	23.0 (2)	165	17.0 (3)	1,029	5.7 (47)	1,029	6.5(43)
MCMH	926	6.6 (43)	55.3 (35)	872	17.3 (15)	55.2 (32)	173	40.5 (32)	224	30.8(38)	1,228	1.7 (9)	1,228	2.2(12)
MHS	4,704	4.6 (11)	55.5 (39)	4,448	18.5 (28)	53.5 (23)	906	39.3 (29)	1,284	31.8(41)	5,710	1.5 (8)	5,710	2.2(12)
MW	1,066	6.7 (44)	52.3 (20)	989	18.1 (24)	53.2 (21)	195	20.0 (1)	808	21.9(16)	1,232	3.7 (35)	1,232	5.8(38)
NCCAA	2,146	5.2 (22)	54.5 (33)	1,962	16.1 (6)	54.8 (29)	444	34.0 (19)	793	26.4(25)	2,567	2.5 (20)	2,567	2.1(11)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).
- (5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).
- (6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 5F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Health Indicators</b> <sup>(2, 3)</sup>														
<b>Local Agency</b>	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight	Over- weight	%	< Ideal	> Ideal	%	3rd Trimester		Postpartum					
							Number	(Rank)	Number	(Rank)	Number	(Rank)	Number	(Rank)
NHC	8,533	4.7 (12)	53.6 (26)	8,121	20.3 (45)	51.3 (3)	1,572	40.3 (31)	1,609	30.9(39)	10,179	2.2 (16)	10,179	3.5(23)
NHS	3,277	5.5 (32)	55.6 (41)	3,128	20.0 (42)	52.1 (11)	609	36.5 (24)	766	28.5(33)	3,932	5.1 (44)	3,932	3.6(25)
NWFS	10,630	4.4 (7)	55.4 (36)	9,498	17.9 (23)	55.1 (30)	2,223	40.8 (34)	2,006	28.6(34)	13,844	2.2 (16)	13,844	3.4(22)
ODCS	1,976	5.4 (27)	54.4 (32)	1,860	18.9 (34)	56.2 (38)	388	40.7 (33)	311	31.2(40)	2,515	2.1 (15)	2,515	2.0(10)
PHNA	2,039	5.7 (33)	49.9 (3)	1,910	18.5 (28)	53.0 (17)	379	30.6 (8)	585	24.4(22)	2,428	2.6 (22)	2,428	1.7(8)
St Mary's	1,292	6.0 (38)	51.9 (14)	1,225	17.1 (14)	54.4 (28)	263	30.8 (9)	264	22.7(17)	1,615	3.2 (28)	1,615	5.4(35)
STARCHD	466	5.4 (27)	50.9 (8)	446	15.7 (5)	56.1 (36)	60	* *	64	* *	550	2.5 (20)	550	4.4(30)
STEUBCHD	704	4.8 (16)	55.8 (42)	677	21.4 (48)	52.6 (14)	76	* *	108	20.4 (8)	816	1.1 (5)	816	0.6(2)
SVJH	756	6.7 (44)	53.0 (22)	727	19.5 (38)	51.6 (6)	77	* *	118	28.8(35)	840	1.0 (3)	840	1.7(8)
TCHD	3,240	4.8 (16)	50.7 (6)	3,060	19.3 (36)	51.5 (5)	512	32.8 (15)	466	20.4 (8)	3,805	2.0 (14)	3,805	1.4(6)
VCHD	2,576	4.5 (9)	55.4 (36)	2,174	18.4 (26)	56.8 (42)	513	37.2 (26)	1,801	23.0(18)	2,690	2.8 (25)	2,690	4.2(29)
WCHD	476	5.9 (37)	50.6 (5)	455	20.2 (44)	52.7 (15)	89	* *	131	21.4(14)	585	3.6 (34)	585	3.2(21)
WCMH	745	4.3 (5)	52.1 (16)	722	19.8 (41)	51.1 (2)	108	36.1 (23)	132	29.5(37)	896	0.7 (2)	896	0.8(3)
WICAA	2,125	5.0 (19)	53.6 (26)	1,823	16.7 (11)	56.6 (40)	373	34.3 (20)	1,214	28.4(32)	2,238	1.9 (12)	2,238	2.4(14)
WVHS	1,050	5.4 (27)	56.6 (44)	917	18.6 (32)	53.7 (25)	191	36.6 (25)	695	28.9(36)	1,168	3.2 (28)	1,168	5.8(38)
Unknown Local Agencies	295	8.8 (50)	48.5 (1)	1	* *	* *	78	* *	3	* *	4	* *	4	* *
<b>Indiana</b>	<b>109,475</b>	<b>5.0</b>	<b>54.7</b>	<b>101,695</b>	<b>18.7</b>	<b>53.5</b>	<b>21,025</b>	<b>37.0</b>	<b>27,836</b>	<b>26.9</b>	<b>135,164</b>	<b>3.0</b>	<b>135,164</b>	<b>4.1</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).  
(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).  
(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Behavioral Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	Medical Care			WIC Enrollment		Multivitamin Consumption				
	1st Trimester		No Care	1st Trimester		Pregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
BCA	364	63.2 (28)	31.3 (28)	468	44.4 (21)	465	11.8 (46)	227	91.2 (5)	
BCHD	1,066	56.0 (39)	35.4 (35)	1,299	36.0 (42)	1,291	18.4 (13)	661	90.3 (10)	
BRSI	1,173	51.7 (42)	42.6 (41)	1,612	54.2 (5)	1,614	17.7 (17)	806	89.2 (19)	
CAP	1,223	51.2 (43)	43.2 (43)	1,499	42.6 (28)	1,495	16.9 (21)	736	89.4 (18)	
CCHD	404	57.7 (36)	38.1 (37)	592	54.9 (3)	589	17.3 (18)	298	86.6 (34)	
CFS	1,918	57.7 (36)	38.3 (38)	2,598	49.6 (13)	2,578	18.2 (15)	1,294	89.7 (14)	
DCH	924	68.3 (15)	26.7 (17)	1,372	40.0 (31)	1,364	15.0 (31)	690	88.1 (25)	
DCHD	399	78.4 (3)	17.0 (4)	596	34.2 (45)	589	21.6 (7)	293	86.3 (35)	
DMH	555	56.4 (38)	39.3 (39)	749	53.8 (6)	744	13.4 (41)	360	82.2 (42)	
DMHC	1,930	59.1 (34)	35.3 (34)	2,664	50.4 (11)	2,576	12.9 (43)	1,274	78.6 (47)	
ECHD	1,054	68.9 (13)	22.5 (8)	1,443	34.7 (44)	948	16.1 (28)	670	88.1 (25)	
ELKCHD	4,372	42.2 (47)	53.4 (47)	5,125	53.5 (7)	5,135	19.8 (9)	2,567	81.5 (44)	
FHHC	152	52.6 (41)	42.8 (42)	198	50.5 (10)	178	23.0 (4)	89	* *	
FLCHD	787	60.5 (32)	33.2 (31)	1,116	39.3 (33)	1,096	19.5 (11)	568	89.1 (21)	
FULCHD	293	67.6 (16)	27.6 (18)	426	43.0 (26)	423	14.9 (33)	210	89.0 (22)	
GCRC	458	72.1 (8)	24.7 (14)	678	44.7 (19)	665	15.0 (31)	352	91.5 (4)	
HCHF	596	78.7 (2)	15.1 (3)	953	30.3 (48)	935	22.4 (5)	498	92.6 (2)	
HCMH&HS	460	65.4 (21)	29.8 (22)	654	36.7 (37)	653	19.8 (9)	349	91.7 (3)	
HIS	964	63.8 (26)	30.5 (26)	1,316	40.5 (30)	1,309	14.9 (33)	683	84.9 (40)	
HU	94	* *	* *	142	54.9 (3)	143	21.7 (6)	72	* *	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 6F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Behavioral Indicators by Local Agency**  
**Combined 3 Years**

Comparison of Maternal Behavioral Indicators <sup>(2, 3)</sup>									
Local Agency	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Prenatal (4)		During Pregnancy (5)	
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
ICAP	642	67.0 (18)	29.8 (22)	835	49.3 (14)	832	13.9 (40)	402	81.6 (43)
IHC	4,707	65.4 (21)	28.0 (19)	6,542	39.7 (32)	6,526	14.8 (35)	3,303	87.8 (28)
JEFCHD	480	60.6 (31)	36.3 (36)	589	46.9 (15)	583	14.4 (38)	288	86.1 (36)
JNHC	705	48.1 (44)	48.8 (45)	937	53.1 (8)	937	15.9 (29)	447	82.6 (41)
JOCHD	1,303	66.2 (19)	26.3 (16)	1,636	31.7 (47)	1,610	17.2 (19)	795	89.6 (15)
KCHD	732	76.8 (5)	14.9 (2)	1,005	29.3 (49)	1,002	16.4 (24)	542	88.0 (27)
LCHD	227	79.7 (1)	12.8 (1)	342	36.0 (42)	337	16.3 (26)	182	87.4 (32)
MADCHD	1,698	61.1 (30)	24.2 (13)	2,173	36.4 (40)	2,172	14.6 (36)	1,127	89.5 (16)
MARCHD	14,872	69.3 (12)	21.9 (7)	20,267	32.4 (46)	20,012	11.6 (47)	10,402	86.9 (33)
MARSCHD	688	65.7 (20)	28.6 (20)	860	36.5 (39)	857	18.4 (13)	444	87.6 (30)
MCMH	691	64.7 (24)	29.5 (21)	933	36.4 (40)	927	18.8 (12)	461	90.0 (12)
MHS	3,568	71.3 (9)	22.5 (8)	4,742	37.5 (35)	4,708	16.2 (27)	2,409	91.2 (5)
MW	730	67.5 (17)	29.9 (24)	1,078	46.3 (17)	1,068	16.4 (24)	539	89.2 (19)
NCCAA	1,618	58.7 (35)	33.1 (30)	2,159	39.0 (34)	1,889	20.5 (8)	1,049	93.7 (1)
NHC	6,360	59.8 (33)	34.0 (32)	8,555	44.0 (22)	8,533	14.6 (36)	4,274	89.5 (16)
NHS	2,523	63.7 (27)	30.6 (27)	3,294	43.7 (24)	3,281	16.5 (23)	1,643	90.1 (11)
NWFS	7,091	68.4 (14)	23.6 (12)	10,696	36.7 (37)	10,609	16.9 (21)	5,311	87.6 (30)
ODCS	1,389	72.4 (7)	22.6 (10)	1,993	43.8 (23)	1,969	17.9 (16)	1,019	90.5 (9)
PHNA	1,380	64.9 (23)	30.2 (25)	2,053	45.8 (18)	1,408	26.3 (1)	969	91.1 (8)
St Mary's	955	64.0 (25)	32.1 (29)	1,300	42.3 (29)	1,296	10.3 (49)	702	88.5 (23)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 6F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Maternal Behavioral Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Maternal Behavioral Indicators</b> <sup>(2, 3)</sup>										
<b>Local Agency</b>	<b>Medical Care</b>			<b>WIC Enrollment</b>			<b>Multivitamin Consumption</b>			
	<b>1st Trimester</b>		<b>No Care</b>	<b>1st Trimester</b>			<b>Prepregnancy</b> (4)		<b>During Pregnancy</b> (5)	
	<b>Number</b>	<b>% (Rank)</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>		<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
STARCHD	370	53.5 (40)	41.9 (40)	463	52.7 (9)		467	15.4 (30)	230	80.9 (45)
STEUBCHD	517	47.2 (46)	49.9 (46)	709	62.2 (2)		701	12.6 (44)	366	80.6 (46)
SVJH	624	28.0 (48)	68.3 (48)	756	65.7 (1)		756	12.3 (45)	362	77.9 (48)
TCHD	2,690	47.6 (45)	45.7 (44)	3,238	50.1 (12)		3,009	23.5 (3)	1,574	86.1 (36)
VCHD	1,699	71.1 (10)	22.7 (11)	2,621	37.5 (35)		2,586	11.6 (47)	1,314	91.2 (5)
WCHD	337	62.0 (29)	34.7 (33)	479	46.6 (16)		475	26.1 (2)	221	90.0 (12)
WCMH	539	69.8 (11)	25.4 (15)	751	44.6 (20)		749	14.3 (39)	379	85.2 (39)
WICAA	1,149	76.5 (6)	18.7 (6)	2,139	43.2 (25)		2,127	13.0 (42)	1,111	87.7 (29)
WVHS	644	77.5 (4)	18.0 (5)	1,064	43.0 (26)		1,053	9.5 (50)	577	85.8 (38)
Unknown Local Agencies	1	*	*	305	16.7 (50)		294	17.0 (20)	296	88.5 (23)
<b>Indiana</b>	<b>80,115</b>	<b>63.3</b>	<b>30.0</b>	<b>110,014</b>	<b>40.7</b>		<b>107,563</b>	<b>15.6</b>	<b>55,435</b>	<b>87.7</b>

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Indiana**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
BCA	612	39.1 (32)	586	29.2 (40)	223	33.2 (20)	464	51.3 (35)	579	48.4 (36)
BCHD	1,632	35.8 (23)	1,590	25.3 (24)	564	35.6 (11)	1,289	51.6 (36)	1,570	46.5 (34)
BRSI	1,977	42.1 (43)	1,907	29.3 (41)	799	32.2 (29)	1,614	49.6 (31)	1,879	36.7 (21)
CAP	1,838	43.1 (47)	1,781	31.0 (44)	768	32.6 (25)	1,491	59.1 (49)	1,763	53.3 (46)
CCHD	703	39.4 (36)	684	27.2 (33)	267	33.3 (19)	590	56.6 (48)	676	52.7 (45)
CFS	3,185	39.2 (35)	3,094	27.2 (33)	1,199	30.4 (37)	2,568	46.5 (26)	3,038	41.0 (27)
DCH	1,744	42.6 (44)	1,674	31.3 (45)	699	33.0 (21)	1,362	53.5 (39)	1,637	49.9 (40)
DCHD	782	26.6 (6)	767	18.6 (8)	203	34.5 (14)	578	41.0 (18)	757	37.8 (22)
DMH	931	40.5 (41)	901	27.7 (35)	363	36.6 (8)	745	50.3 (33)	895	46.4 (33)
DMHC	3,267	37.5 (28)	3,171	26.8 (29)	1,178	31.9 (32)	2,657	52.5 (37)	3,130	46.1 (32)
ECHD	1,852	13.8 (1)	1,805	5.8 (1)	246	56.9 (1)	1,438	19.1 (1)	1,784	13.5 (5)
ELKCHD	6,105	23.7 (4)	5,895	14.8 (3)	1,376	32.8 (22)	5,122	21.8 (3)	5,832	7.8 (2)
FHHC	240	44.2 (49)	240	28.8 (37)	105	35.2 (12)	198	54.0 (41)	237	51.9 (43)
FLCHD	1,409	38.2 (29)	1,342	25.1 (23)	508	40.0 (6)	1,093	48.9 (30)	1,313	46.0 (31)
FULCHD	517	35.6 (21)	491	24.8 (22)	175	36.6 (8)	422	38.9 (16)	486	33.5 (16)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Indiana**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
GCRC	870	43.2 (48)	824	33.9 (48)	347	29.1 (41)	667	61.0 (50)	818	55.0 (48)
HCHF	1,273	31.3 (13)	1,136	15.7 (6)	324	46.3 (3)	943	43.3 (22)	1,114	39.4 (24)
HCMH&HS	866	34.5 (17)	834	22.7 (15)	289	34.3 (16)	650	34.5 (9)	823	16.8 (6)
HIS	1,693	37.1 (26)	1,636	27.1 (30)	591	31.0 (35)	1,306	55.0 (45)	1,621	52.4 (44)
HU	172	43.0 (45)	166	30.7 (43)	68	* *	143	41.3 (19)	165	25.5 (11)
ICAP	1,003	46.1 (50)	979	31.8 (46)	447	30.6 (36)	827	53.0 (38)	966	49.5 (38)
IHC	8,196	29.6 (9)	7,913	22.1 (13)	2,314	29.6 (40)	6,525	35.9 (13)	7,813	24.9 (10)
JEFCHD	733	42.0 (42)	704	34.7 (49)	293	27.0 (44)	581	40.6 (17)	694	28.4 (13)
JNHC	1,120	43.0 (45)	1,083	28.8 (37)	462	32.7 (23)	934	54.2 (42)	1,072	49.0 (37)
JOCHD	2,227	30.8 (10)	2,156	23.1 (18)	649	32.5 (27)	1,624	41.9 (20)	2,116	34.2 (18)
KCHD	1,275	30.8 (10)	1,205	22.0 (12)	356	32.3 (28)	1,000	44.4 (25)	1,190	34.1 (17)
LCHD	434	31.6 (14)	412	23.8 (19)	125	30.4 (37)	337	48.4 (29)	400	42.5 (28)
MADCHD	2,920	34.9 (19)	2,843	27.1 (30)	981	32.0 (31)	2,171	50.3 (33)	2,814	46.8 (35)
MARCHD	29,025	18.0 (2)	27,471	14.6 (2)	4,769	36.7 (7)	19,086	28.1 (6)	26,505	27.0 (12)
MARSCHD	1,063	32.9 (16)	1,029	21.9 (11)	331	34.4 (15)	857	20.1 (2)	1,020	5.2 (1)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Indiana**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	<b>Smoking</b>				<b>Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)</b>		<b>Smoking in Household</b>			
	<b>3 Months Prior to Pregnancy</b>		<b>Last 3 Months of Pregnancy</b>				<b>Prenatal</b>		<b>Postpartum</b>	
	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>	<b>Number</b>	<b>% (Rank)</b>
MCMH	1,275	38.4 (30)	1,228	33.6 (47)	462	26.6 (46)	917	56.3 (47)	1,214	55.0 (48)
MHS	5,934	27.9 (7)	5,710	15.1 (5)	1,568	47.8 (2)	4,701	24.7 (4)	5,637	17.2 (7)
MW	1,304	39.1 (32)	1,232	25.8 (26)	468	36.1 (10)	1,065	47.0 (27)	1,220	35.3 (20)
NCCAA	2,734	31.7 (15)	2,567	24.4 (21)	790	31.8 (33)	2,146	43.1 (21)	2,526	39.4 (24)
NHC	10,515	28.3 (8)	10,179	20.1 (9)	2,830	31.1 (34)	8,495	37.6 (14)	10,045	34.2 (18)
NHS	4,061	36.7 (25)	3,931	25.5 (25)	1,426	34.3 (16)	3,273	43.4 (23)	3,887	38.8 (23)
NWFS	14,904	22.1 (3)	13,844	15.0 (4)	3,003	42.2 (4)	10,606	33.0 (8)	13,665	29.0 (14)
ODCS	2,620	34.5 (17)	2,515	28.3 (36)	862	28.4 (42)	1,892	35.3 (11)	2,486	23.3 (8)
PHNA	2,544	35.3 (20)	2,428	22.7 (15)	840	32.6 (25)	1,893	35.8 (12)	2,395	23.5 (9)
St Mary's	1,674	39.7 (37)	1,615	28.9 (39)	637	28.1 (43)	1,292	54.7 (43)	1,593	54.2 (47)
STARCHD	566	40.1 (40)	550	27.1 (30)	217	32.7 (23)	464	53.9 (40)	547	51.4 (42)
STEUBCHD	838	40.0 (38)	816	26.5 (28)	324	30.2 (39)	701	54.8 (44)	802	49.8 (39)
SVJH	864	37.4 (27)	840	30.6 (42)	310	15.5 (48)	756	56.0 (46)	829	50.9 (41)
TCHD	3,948	24.4 (5)	3,805	18.0 (7)	917	26.7 (45)	3,180	34.8 (10)	3,754	30.8 (15)
VCHD	3,077	38.6 (31)	2,690	20.7 (10)	995	41.7 (5)	2,569	26.7 (5)	2,640	9.3 (3)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 7F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** (1)  
**Indiana**  
**Comparison of Smoking Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Smoking Indicators</b> (2, 3)										
<b>Local Agency</b>	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
WCHD	604	35.6 (21)	585	22.2 (14)	206	35.0 (13)	474	50.2 (32)	583	44.4 (30)
WCMH	916	30.9 (12)	896	23.8 (19)	273	21.2 (47)	746	28.7 (7)	884	13.1 (4)
WICAA	2,525	40.0 (38)	2,238	26.2 (27)	853	33.5 (18)	2,128	47.6 (28)	2,219	43.1 (29)
WVHS	1,295	36.4 (24)	1,168	22.9 (17)	414	32.1 (30)	1,054	43.8 (24)	1,136	40.1 (26)
Unknown Local Agencies	307	39.1 (32)	4	* *	1	* *	291	38.5 (15)	3	* *
<b>Indiana</b>	<b>142,169</b>	<b>29.0</b>	<b>135,160</b>	<b>20.5</b>	<b>38,415</b>	<b>34.2</b>	<b>107,925</b>	<b>37.5</b>	<b>132,772</b>	<b>31.6</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) Among women who smoked 3 months prior to pregnancy.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
BCA	568	8.1 (30)	6.2 (20)	435	12.2 (13)	382	2.9 (31)	568	56.7 (46)
BCHD	1,542	6.8 (7)	6.5 (22)	1,242	13.8 (29)	1,069	2.1 (12)	1,542	68.4 (17)
BRSI	1,804	8.8 (39)	5.9 (17)	1,480	15.1 (37)	1,254	3.3 (40)	1,804	70.1 (14)
CAP	1,706	7.5 (22)	7.6 (35)	1,405	13.6 (26)	1,211	2.4 (17)	1,709	62.5 (33)
CCHD	655	4.9 (1)	7.0 (29)	553	10.1 (1)	497	1.4 (3)	654	62.5 (33)
CFS	2,945	8.3 (34)	4.8 (5)	2,429	13.4 (24)	2,102	2.7 (25)	2,949	68.2 (18)
DCH	1,593	7.2 (15)	7.7 (39)	1,282	11.6 (8)	1,121	2.0 (11)	1,608	58.8 (42)
DCHD	737	6.5 (6)	6.8 (26)	570	13.2 (22)	493	1.4 (3)	737	75.4 (4)
DMH	872	9.7 (46)	5.3 (8)	703	10.4 (3)	627	4.6 (49)	874	65.4 (27)
DMHC	3,040	7.0 (10)	6.9 (27)	2,501	12.8 (18)	2,171	2.3 (13)	3,045	58.3 (43)
ECHD	1,744	10.0 (48)	5.7 (13)	1,367	17.5 (46)	1,125	2.8 (28)	1,748	62.4 (35)
ELKCHD	5,567	7.3 (17)	7.9 (42)	4,751	11.6 (8)	4,182	2.3 (13)	5,586	74.6 (6)
FHHC	235	8.1 (30)	3.8 (1)	197	15.2 (39)	165	3.6 (45)	237	59.1 (41)
FLCHD	1,297	9.6 (44)	4.9 (6)	1,037	19.1 (48)	833	3.0 (34)	1,302	65.7 (26)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
FULCHD	474	7.4 (20)	5.7 (13)	396	15.9 (40)	328	1.5 (6)	475	66.5 (25)
GCRC	798	7.8 (25)	4.0 (2)	623	13.6 (26)	534	3.2 (38)	806	53.6 (49)
HCHF	1,107	7.6 (23)	7.8 (41)	806	11.2 (6)	716	2.7 (25)	1,107	74.3 (7)
HCMH&HS	808	6.1 (2)	7.4 (33)	614	12.9 (19)	535	1.1 (2)	808	71.5 (11)
HIS	1,590	7.1 (13)	6.9 (27)	1,241	12.1 (11)	1,084	3.4 (43)	1,598	67.5 (21)
HU	161	6.2 (3)	7.5 (34)	132	19.7 (49)	106	0.0 (1)	161	54.7 (48)
ICAP	958	7.8 (25)	7.6 (35)	804	10.1 (1)	721	4.3 (48)	960	68.1 (19)
IHC	7,610	7.9 (28)	6.6 (24)	6,137	12.5 (16)	5,344	3.0 (34)	7,632	62.4 (35)
JEFCHD	672	9.8 (47)	7.9 (42)	549	16.4 (44)	455	2.6 (22)	673	67.8 (20)
JNHC	1,039	7.3 (17)	8.8 (47)	881	13.5 (25)	755	2.9 (31)	1,045	66.9 (23)
JOCHD	2,073	7.3 (17)	7.6 (35)	1,535	13.3 (23)	1,324	2.6 (22)	2,080	70.2 (13)
KCHD	1,161	7.4 (20)	6.5 (22)	912	14.0 (31)	784	1.4 (3)	1,162	70.3 (12)
LCHD	392	8.7 (37)	5.9 (17)	313	16.3 (43)	260	2.3 (13)	394	73.4 (9)
MADCHD	2,782	8.1 (30)	5.5 (11)	2,079	13.8 (29)	1,783	3.3 (40)	2,784	57.3 (44)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.



**Table 8F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
MARCHD	26,744	9.0 (40)	6.1 (19)	18,478	14.8 (36)	15,683	2.8 (28)	26,826	63.1 (32)
MARSCHD	995	6.4 (4)	8.1 (44)	808	11.0 (4)	719	1.9 (9)	995	72.7 (10)
MCMH	1,198	8.0 (29)	6.7 (25)	872	12.5 (16)	762	2.4 (17)	1,200	67.4 (22)
MHS	5,517	8.5 (36)	7.3 (32)	4,445	12.4 (15)	3,885	2.5 (19)	5,528	73.6 (8)
MW	1,178	7.0 (10)	7.2 (30)	983	12.0 (10)	863	1.5 (6)	1,180	63.8 (31)
NCCAA	2,464	9.1 (41)	5.5 (11)	1,941	14.5 (33)	1,655	3.3 (40)	2,463	60.9 (38)
NHC	9,765	8.7 (37)	5.4 (9)	8,013	16.0 (41)	6,709	3.4 (43)	9,762	63.9 (30)
NHS	3,764	7.8 (25)	5.7 (13)	3,079	14.1 (32)	2,631	3.1 (37)	3,765	65.0 (28)
NWFS	13,357	8.4 (35)	6.2 (20)	9,406	14.7 (35)	7,993	2.9 (31)	13,392	69.0 (16)
ODCS	2,435	9.6 (44)	4.6 (4)	1,848	13.7 (28)	1,586	4.1 (47)	2,447	60.2 (39)
PHNA	2,321	7.2 (15)	8.5 (45)	1,879	11.5 (7)	1,660	2.5 (19)	2,323	74.8 (5)
St Mary's	1,556	9.1 (41)	5.8 (16)	1,214	17.1 (45)	999	1.9 (9)	1,562	55.4 (47)
STARCHD	528	6.4 (4)	9.1 (48)	435	12.9 (19)	379	3.2 (38)	525	66.7 (24)
STEUBCHD	751	7.6 (23)	9.1 (48)	662	12.1 (11)	573	2.8 (28)	762	80.1 (1)

- (1) Reporting period is January 1 through December 31.
- (2) Excludes records with unknown data or errors.
- (3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.
- (4) <2500 g.
- (5) >4000 g.
- (6) <37 weeks gestation.
- (7) 37 or more weeks gestation and <2500 g.
- \* Percentages are not calculated if <100 records are available for analysis after exclusions.





**Table 8F**  
**2009 - 2011 Pregnancy Nutrition Surveillance** <sup>(1)</sup>  
**Indiana**  
**Comparison of Infant Health Indicators by Local Agency**  
**Combined 3 Years**

<b>Comparison of Infant Health Indicators</b> <sup>(2, 3)</sup>									
<b>Local Agency</b>	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)		High (5)						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
SVJH	791	7.0 (10)	7.6 (35)	705	14.6 (34)	601	1.8 (8)	792	61.7 (37)
TCHD	3,621	6.9 (8)	8.5 (45)	2,997	11.0 (4)	2,656	2.6 (22)	3,630	75.5 (3)
VCHD	2,593	9.3 (43)	5.4 (9)	2,207	16.1 (42)	1,834	2.7 (25)	2,600	59.5 (40)
WCHD	568	10.9 (49)	5.1 (7)	450	18.7 (47)	366	2.5 (19)	567	76.5 (2)
WCMH	844	7.1 (13)	7.7 (39)	699	12.2 (13)	609	3.0 (34)	849	69.1 (15)
WICAA	2,160	6.9 (8)	7.2 (30)	1,821	13.0 (21)	1,566	2.3 (13)	2,176	64.8 (29)
WVHS	1,094	8.2 (33)	4.0 (2)	910	15.1 (37)	769	3.9 (46)	1,103	56.8 (45)
Unknown Local Agencies	3	*	*	0	*	0	*	3	*
<b>Indiana</b>	<b>130,177</b>	<b>8.3</b>	<b>6.4</b>	<b>100,826</b>	<b>13.9</b>	<b>86,459</b>	<b>2.8</b>	<b>130,498</b>	<b>65.8</b>

(1) Reporting period is January 1 through December 31.  
(2) Excludes records with unknown data or errors.  
(3) Rank compares one local agency's rate to other local agencies. Rank 1 = best rate.  
(4) <2500 g.  
(5) >4000 g.  
(6) <37 weeks gestation.  
(7) 37 or more weeks gestation and <2500 g.  
\* Percentages are not calculated if <100 records are available for analysis after exclusions.

