



Class III State Regulated Wetlands

(317) 232-8603 • (800) 451-6027

www.idem.IN.gov

100 N. Senate Ave., Indianapolis, IN 46204

Class III State Regulated Wetland Types

Indiana State Regulated Class III wetlands are defined at IC 13-11-2-25.8 (3). Class III wetlands include any non-jurisdictional wetland that is any of the following rare and ecologically important types:

- Acid bog
- Acid seep
- Circumneutral bog
- Circumneutral seep
- Cypress swamp
- Dune and swale
- Fen
- Forested Fen
- Forested swamp
- Marl beach
- Muck flat
- Panne
- Sand flat
- Sedge meadow
- Shrub swamp
- Sinkhole pond
- Sinkhole swamp
- Wet floodplain forest
- Wet prairie
- Wet sand prairie

A state regulated wetland may also be classified as Class III when it is located in a setting undisturbed or minimally disturbed by human activity or development that supports more than minimal wildlife, aquatic habitat, or hydrologic function.

A wetland may also be considered Class III even if it does not meet the above criteria when the wetland contains the presence of, or habitat for rare, threatened, or endangered species. That is, a wetland that would otherwise meet the definition of a Class II wetland but contains rare, threatened, or endangered species is considered a Class III wetland for permitting purposes.

After the State Regulated Wetland Law first became effective in 2003, IDEM facilitated a Wetland Science Advisory Group, made up of a cross-section of wetland science experts, to develop definitions for the list of rare and ecologically important types.

Definitions of the Rare and ecologically important wetland types:

Acid Bog

Acid bog is an acidic wetland of kettle holes in glacial terrain. Bogs can be graminoid (*Carex* spp. and *Sphagnum* spp.) or low shrub (*Chamaedaphne calyculata* and *Betula pumila*). The graminoid bog can be a floating, quaking mat. The soils in acid bogs are saturated and acidic peat. Bogs have non-flowing or very slow flowing water. The water level fluctuates seasonally. When a sphagnum mat floats, it rises and falls with the water table. Acid bogs can be found in northern Indiana.

Acid Seep

Acid seep is a bog-like wetland typically found in unglaciated hill regions. This community is a small groundwater-fed wetland located primarily in upland terrain. A thin layer of muck may lie over a mineral substrate. The soil reaction is acid. This seep community is characterized by flowing water during at least part of the year. Acid seeps are located primarily in southern Indiana.

Circumneutral bog

Circumneutral bog is a bog-like wetland that receives groundwater. Circumneutral bogs can be a mosaic of tall shrub bog, graminoid bog, and other communities. The graminoid bog often occurs on a quaking or floating mat. Although a few bogs occur in unglaciated regions, most are found in glacial ice-block depressions. The soils in circumneutral bogs are usually peat, or other low nutrient organic substrates, which are saturated and circumneutral to slightly acid. Circumneutral bogs have non-flowing or very slow flowing water. The water level fluctuates seasonally. Circumneutral bogs are usually found in northern Indiana.

Circumneutral Seep

The circumneutral seep (or seep-spring) is a groundwater-fed wetland on organic soil. It is primarily herbaceous. Species typically include marsh marigold (*Caltha palustris*) and skunk cabbage (*Symplocarpus foetidus*) with a scattered tree canopy. Circumneutral seep is typically situated on or near the base of a slope. The soil is typically circumneutral muck. This seep community is characterized by slowly flowing water during at least part of the year. Circumneutral seeps can be found scattered throughout Indiana.

Cypress swamp

Bald cypress swamps are seasonally to permanently inundated wetlands found in depressions and sloughs of large bottomlands associated with the Wabash/Ohio River system. Poorly to very poorly drained soils characterize this environment. Bald cypress (*Taxodium distichum*) is present, and green ash (*Fraxinus pennsylvanica*), silver maple (*Acer saccharinum*), and overcup oak (*Quercus lyrata*) are also usually present. This community is restricted to extreme southwest Indiana.

Dune and Swale

Dune and swale is an ecological system consisting of a mixture of upland (black oak sand savanna, dry to mesic sand prairie) and wetland (pond, panne, sedge meadow, marsh, wet prairie) natural communities. These communities occur in long, narrow, linear complexes, with

the dry communities occupying sand ridges, and the wet communities occurring in the intervening swales. Black oak (*Quercus velutina*), paper birch (*Betula papyrifera*), jack pine (*Pinus banksiana*), and prairie vegetation typically occur on the ridges, and sedges, reeds, and marsh/aquatic vegetation line are found in the swales. Water levels are directly influenced by ground water, with the interdunal swales controlled largely by lateral flow through porous beach ridges. Dune and swale is restricted to extreme northwest Indiana, near Lake Michigan.

Fen

Fen is a calcareous, groundwater-fed wetland. Fens are often a mosaic of grassy areas, sedgy areas, graminoid-shrubby cinquefoil, and tall shrub areas. The extent of the tall shrub component of fens may be determined by fire frequency and/or soil moisture. Drying of the soil increases the growth of shrubs. Fens typically occur in the vicinity of glacial moraines. Fens typically have a muck or peat substrate. The water level fluctuates seasonally and is fed by groundwater. Fens can be found in central and northern Indiana.

Forested fen

Forested fen is a tree-dominated wetland on organic soil which receives groundwater. Forested fens are often a mosaic of treed areas, tall shrub areas, and herbaceous areas. A tall shrub layer is often well developed in forested fens. Indicative species typically include tamarack (*Larix laricina*), black ash (*Fraxinus nigra*), yellow birch (*Betula alleghaniensis*), poison sumac (*Toxicodendron vernix*), and red maple (*Acer rubrum*). Forested fens occur in wet lowlands, where moraines meet outwash features or depressions. Forested fens have saturated, poorly to very poorly drained soils that are often muck, but some seasonal flooding can occur in forested fens that are especially level. This community is a late successional stage of fen or circumneutral bog. Forested fens occur in northern Indiana.

Forested Swamp

Forested swamp is a seasonally inundated to intermittently exposed wetland of large river bottoms. Forested swamps do not receive direct flow from river flooding except under exceptional circumstances. Forested swamps occur in depressions, sloughs and large bottomlands, typically dominated by tree species such as swamp cottonwood (*Populus heterophylla*), green ash (*Fraxinus pennsylvanica*), and swamp white oak (*Quercus bicolor*). In northern Indiana important tree species include black ash (*Fraxinus nigra*), yellow birch (*Betula alleghaniensis*), and red maple (*Acer rubrum*). Poorly to very poorly drained and aerated soils characterize the swamp environment. Soils usually are mineral not muck or peat. This community type is found throughout Indiana.

Marl beach

Marl beach is a fen-like community located on the marly muck shorelines of lakes. Marl precipitate is evident. A thin layer of water is present in spring, but dries down in summer. Draw-down of a lake creates additional area for this community to develop on. Marl beaches can be found in extreme northern Indiana, primarily in the northeast.

Muck flat

Muck flat is a shoreline and lake community possessing a unique flora of sedges and annual plants, many of which are also found on the Atlantic and Gulf Coastal Plains. This community is

found at the margins of lakes or covering shallow basins. This community has a peat substrate. The muck flats can float on the water surface, but during high water periods are usually inundated. The water level of a basin fluctuates during a season or from year to year in response to the amount of precipitation. This exposes bare substrate needed for germination by species of the community. Muck flats are found in northern Indiana.

Panne

Panne is a groundwater fed herbaceous wetland occupying interdunal swales near Lake Michigan. Pannes are located on the lee side of the first or second line of dunes from the lakeshore. The soil is wet, calcareous sand. Pannes are located in counties bordering Lake Michigan.

Sand flat

Sand flat is a shoreline and lake community possessing a unique flora of sedges and annual plants, many of which are also found on the Atlantic and Gulf Coastal Plains. This community is found at the margins of lakes or covering shallow basins. This community has a sand substrate. During high water periods sand flats at the margins of lakes or ponds are inundated. The water level of a basin fluctuates during a season or from year to year in response to the amount of precipitation. This exposes bare substrate needed for germination by species of the community. Sand flats occur in northern Indiana, and in the Plainville Sand Section of southwest Indiana.

Sedge meadow

Sedge meadow is an herbaceous wetland typically dominated by graminoid species such as flat sedge (*Cyperus* spp.), spike rush (*Eleocharis* spp.), rushes (*Juncus* spp.) and sedges (*Carex* spp.). Sedge meadow is an herbaceous wetland of stream margins and river floodplains, and lake margins or upland depressions. Streamside sedge meadows are frequently flooded in the spring and early summer. Sedge meadows of lake margins and depressions often contain standing water during wet months and after heavy rains; during dry periods, the water level is at or just below the substrate. Sedge meadow usually occupies the ground between a marsh and the uplands, or a shrub swamp or wet forest. Periodic high water can kill trees and shrubs invading sedge meadows. Sedge meadows can be found in the northern half of the state.

Shrub swamp

Shrub swamp is a shrub-dominated wetland that is seasonally inundated to intermittently exposed. This community occurs in depressions and the substrate is either mineral soils or muck, as opposed to peat which is characteristic of bogs. Shrub swamp is characterized by non-flowing or very slowly flowing water with levels that fluctuate seasonally. Shrub swamps are persistent, though considered successional. Two opportunistic native shrubs, sandbar willow (*Salix exigua*) and gray dogwood (*Cornus racemosa*), by themselves, are not indicative of shrub swamps. This community type is found throughout Indiana.

Sinkhole pond

Sinkhole ponds are water-containing depressions in karst topography. Sinkhole ponds are found in the Mitchell Karst Plain in south-central Indiana.

Sinkhole Swamp

Sinkhole swamps are depressions in karst topography dominated by tree or shrub species. Sinkhole swamps are found in the Mitchell Karst Plain in south-central Indiana.

Wet floodplain forest

Wet floodplain forest is a broadleaf deciduous forest of river floodplains. Wet floodplain forests occur in depressions and flats on narrow to wide floodplains and also on recently exposed substrates that are frequently flooded. Wet floodplain forests are frequently flooded and may have standing water seasonally to permanently present. Wet floodplain forests occur statewide.

Wet prairie

Wet prairie is an herbaceous wetland typically dominated by graminoid species such as prairie cordgrass (*Spartina pectinata*), bluejoint (*Calamagrostis canadensis*), and sedges (*Carex* spp.). Vegetation height is often 2-3 m. The species diversity of wet prairies is lower than that of mesic prairies. Wet prairies occur in deep swales and the substrate ranges from very deep black mineral soils (which are high in organic matter) to muck. Ponding in spring lasts for several weeks prior to drainage. Wet prairies commonly occur in the Grand Prairie Natural Region, the Tipton Till Plain and the Bluffton Till Plain, with a few examples found in the Northern Lakes Natural Region.

Wet sand prairie

Wet sand prairie is an herbaceous wetland typically dominated by graminoid species such as prairie cordgrass (*Spartina pectinata*), bluejoint (*Calamagrostis canadensis*), and sedges (*Carex* spp.). Vegetation height is often 2-3 m. The species diversity of wet prairies is lower than that of mesic prairies. Wet lowland prairies occur in deep swales and the substrate is sand, sometimes mixed with muck. Flooding is a regular springtime occurrence in wet sand prairie and may last several weeks. This community occurs in a mosaic with marsh and other wetlands, and with upland prairies and sand savannas. Fire was frequent occurrence, but more common in the fall when waters had receded. This community occurs in northwest Indiana and in the Plainsville Sands area.

IDEM's Role

- IDEM is responsible for preserving and protecting the chemical, physical and biological integrity of Indiana's waters, including wetlands.
- IDEM is responsible for the implementation of the State Regulated Wetland Law.

Citizen's Role

- Property owners should always seek out confirmation of regulation applicability through the USACE and IDEM.

More Information

- For information about IDEM's wetland regulations visit:
<https://www.in.gov/idem/wetlands/index.htm>.