





# 3rd Congressional District Nutrient and Sediment Load Reductions

## Accomplished By Private Landowners and the Indiana Conservation Partnership



**Sediment Reduced: 291,676,395 lbs.**  
 Enough to fill 1,458 freight cars!



**Phosphorus Reduced: 192,985 lbs.**  
 Enough to fill 192 truck beds (8' bed)!



**Nitrogen Reduced: 417,390 lbs.**  
 Enough to fill 417 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	2,992	2,992	331,967,615	194,690	403,180
2014	2,018	2,365	254,369,525	154,220	322,525
2015	2,448	3,031	309,059,770	190,600	395,075
2016	2,055	2,976	288,755,885	181,145	376,160
2017	2,526	3,596	324,149,870	201,450	416,745
2018	2,712	4,113	291,676,395	192,985	417,390
<b>13-18</b>	<b>14,751</b>		<b>1,799,979,060</b>	<b>1,115,090</b>	<b>2,331,075</b>

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Load reduction calculations have been rounded to the multiple of 5.