


First Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Comparison of Land Use Across District



Sediment Reduced: 31,338,980 lbs.

Enough to fill 157 freight cars!



Phosphorus Reduced: 22,949 lbs.

Enough to fill 23 truck beds (8' bed)!



Nitrogen Reduced: 38,756 lbs.

Enough to fill 39 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	110	183	6,203	7,209	14,380
2017	131	230	9,350	11,623	23,200
2018	155	274	9,314	11,584	23,109
2019	340	494	16,316	20,741	41,421
2020	256	437	15,817	20,168	40,284
2021	231	431	15,669	22,949	38,756
2013-21	1,672	-	96,172	123,700	239,984

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.



For more information visit:
<http://www.in.gov/isda/2991.htm>
 or contact ISDANutrientReduction@isda.in.gov
 Last updated: 4/20/2022

Second Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership




Comparison of Land Use Across District



Sediment Reduced: 187,110,972 lbs.

Enough to fill 936 freight cars!



x 936

Phosphorus Reduced: 105,928 lbs.

Enough to fill 106 truck beds (8' bed)!



x 106

Nitrogen Reduced: 212,086 lbs.

Enough to fill 212 truck beds (8' bed)!



x 212

Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	1,435	1,913	94,178	106,108	212,518
2017	1,581	2,286	99,346	111,931	224,225
2018	1,366	2,211	84,232	95,494	191,263
2019	1,480	2,558	105,502	113,574	227,219
2020	1,994	3,340	102,236	111,154	222,422
2021	1,110	3,062	93,555	105,928	212,086
2013-21	14,434	-	914,361	1,026,159	2,055,348

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

For more information visit:

<http://www.in.gov/isda/2991.htm>

or contact ISDANutrientReduction@isda.in.gov


Last updated: 4/20/2022

Third Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Comparison of Land Use Across District



Sediment Reduced: 363,224,065 lbs.

Enough to fill 1816 freight cars!



Phosphorus Reduced: 230,857 lbs.

Enough to fill 231 truck beds (8' bed)!



Nitrogen Reduced: 492,810 lbs.

Enough to fill 493 truck beds (8' bed)!



Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	2,055	2,886	143,648	180,196	374,260
2017	2,526	3,506	161,345	200,498	414,845
2018	2,712	4,023	145,109	192,036	415,490
2019	3,615	5,190	221,533	281,570	594,395
2020	2,929	4,861	182,439	234,946	501,196
2021	2,933	5,247	181,612	230,857	492,810
2013-21	24,228	-	1,483,382	1,859,611	3,913,779

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.



For more information visit:
<http://www.in.gov/isda/2991.htm>
 or contact ISDANutrientReduction@isda.in.gov
 Last updated: 4/20/2022

Fourth Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership




Comparison of Land Use Across District



Sediment Reduced: 544,377,465 lbs.

Enough to fill 2722 freight cars!



x 2722

Phosphorus Reduced: 312,407 lbs.

Enough to fill 312 truck beds (8' bed)!



x 312

Nitrogen Reduced: 631,797 lbs.

Enough to fill 632 truck beds (8' bed)!



x 632

Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	1,533	3,266	136,069	156,932	322,263
2017	1,727	3,952	165,706	191,195	390,141
2018	2,002	4,495	186,140	213,171	433,679
2019	1,870	4,784	197,248	223,927	454,975
2020	2,634	5,910	244,433	275,912	558,681
2021	2,635	6,547	272,188	312,407	631,797
2013-21	17,501	-	1,558,122	1,791,285	3,632,657

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

For more information visit:

<http://www.in.gov/isda/2991.htm>

or contact ISDANutrientReduction@isda.in.gov



Last updated: 4/20/2022

Fifth Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Comparison of Land Use Across District



Sediment Reduced: 46,629,399 lbs.

Enough to fill 233 freight cars!



x 233

Phosphorus Reduced: 30,638 lbs.

Enough to fill 31 truck beds (8' bed)!



x 31

Nitrogen Reduced: 62,256 lbs.

Enough to fill 62 truck beds (8' bed)!



x 62

Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	73	225	7,665	9,789	20,617
2017	142	315	11,725	14,769	30,564
2018	162	378	12,073	15,319	31,670
2019	206	444	17,436	22,490	45,995
2020	202	464	19,993	26,422	53,831
2021	249	545	23,314	30,638	62,256
2013-21	1,640	-	128,305	166,799	340,656

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

For more information visit:

<http://www.in.gov/isda/2991.htm>

or contact ISDA.NutrientReduction@isda.in.gov


Last updated: 4/20/2022

Sixth Congressional District Nutrient and Sediment Load Reductions


Accomplished By Private Landowners and the Indiana Conservation Partnership



Comparison of Land Use Across District




- Agriculture - 62%
- Developed - 9%
- Forest and Wetland - 28%
- Other - 1%



Sediment Reduced: 501,690,052 lbs.

Enough to fill 2508 freight cars!



x 2508

Phosphorus Reduced: 257,540 lbs.

Enough to fill 258 truck beds (8' bed)!



x 258

Nitrogen Reduced: 513,232 lbs.

Enough to fill 513 truck beds (8' bed)!



x 513

Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	1,554	2,742	148,540	149,175	298,660
2017	1,476	3,097	150,861	153,530	307,362
2018	2,026	3,943	185,690	191,380	383,701
2019	2,656	5,152	241,723	250,025	500,822
2020	2,801	5,677	276,675	283,094	567,084
2021	2,395	5,798	250,845	257,540	513,232
2013-21	17,378	-	1,639,676	1,672,223	3,346,757

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.


For more information visit:
<http://www.in.gov/isda/2991.htm>
 or contact ISDANutrientReduction@isda.in.gov
 Last updated: 4/20/2022

Seventh Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Comparison of Land Use Across District



Sediment Reduced: 2,511,799 lbs.

Enough to fill 13 freight cars!



x 13

Phosphorus Reduced: 1,215 lbs.

Enough to fill 1 truck beds (8' bed)!



x 1

Nitrogen Reduced: 2,374 lbs.

Enough to fill 2 truck beds (8' bed)!



x 2

Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	33	45	1,009	917	1,770
2017	36	58	1,088	1,039	2,013
2018	29	57	858	723	1,381
2019	23	60	915	789	1,524
2020	25	77	834	696	1,336
2021	26	82	1,255	1,215	2,374
2013-21	225	-	8,320	7,652	14,812

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

For more information visit:

<http://www.in.gov/isda/2991.htm>

or contact ISDA.NutrientReduction@isda.in.gov




Last updated: 4/20/2022

Eighth Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Comparison of Land Use Across District



Sediment Reduced: 868,710,075 lbs.

Enough to fill 4344 freight cars!



x 4344

Phosphorus Reduced: 423,958 lbs.

Enough to fill 424 truck beds (8' bed)!



x 424

Nitrogen Reduced: 853,608 lbs.

Enough to fill 854 truck beds (8' bed)!



x 854

Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	2,254	3,987	373,447	350,236	701,177
2017	2,426	4,697	415,208	388,757	778,483
2018	3,638	6,226	450,756	422,708	846,365
2019	3,308	6,887	478,830	455,184	911,277
2020	3,773	7,924	511,281	482,568	965,875
2021	2,875	7,740	434,355	423,958	853,608
2013-21	26,736	-	3,828,055	3,612,900	7,237,131

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

For more information visit:

<http://www.in.gov/isda/2991.htm>

or contact ISDANutrientReduction@isda.in.gov


Last updated: 4/20/2022

Ninth Congressional District Nutrient and Sediment Load Reductions

Accomplished By Private Landowners and the Indiana Conservation Partnership



Comparison of Land Use Across District




- Agriculture - 36%
- Developed - 10%
- Forest and Wetland - 52%
- Other - 2%



Sediment Reduced: 653,875,450 lbs.

Enough to fill 3269 freight cars!



x 3269

Phosphorus Reduced: 314,359 lbs.

Enough to fill 314 truck beds (8' bed)!



x 314

Nitrogen Reduced: 643,093 lbs.

Enough to fill 643 truck beds (8' bed)!



x 643

Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model.

Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2016	1,650	2,840	284,420	264,469	531,468
2017	1,865	3,462	351,967	326,905	656,426
2018	2,004	3,955	316,046	295,120	593,183
2019	2,392	4,765	333,441	311,967	626,907
2020	2,504	5,301	328,545	307,856	618,591
2021	2,431	5,712	326,937	314,359	643,093
2013-21	17,178	-	2,575,325	2,409,132	4,849,266

The "practices installed" column indicates the number of newly installed practices within a given year, while the "active practices" column indicates the number of practices that are actively reducing sediment and nutrient loading regardless of the year of installation. Please Note: Calendar year 2013 through 2015 metrics are excluded from the table due to space limitations, but are present in the "2013-21" summations.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

For more information visit:
<http://www.in.gov/isda/2991.htm>
 or contact ISDANutrientReduction@isda.in.gov
 Last updated: 4/20/2022