



No. 1086	AUTHOR: Chad M. Stewart, Deer Research Biologist	DATE 7/2/14
	TITLE: 2013 Bovine Tuberculosis Surveillance & Monitoring Summary	

Abstract: *Bovine Tuberculosis (TB) is a chronic bacterial infection that can affect most mammals. In 2009, three captive cervid farms had animals that tested positive for TB, with all animals originating from one location in Northwest Franklin County. To ensure that the disease did not escape into the wild deer population, the Division of Fish and Wildlife and their collaborators, implemented a surveillance program to test hunter harvested deer in the locations near the three captive cervid farms. An additional cattle farm tested positive in 2011, expanding surveillance efforts into Dearborn County. Results from 229 deer sampled in Franklin, Fayette, and Dearborn counties in 2013 failed to detect the presence of TB. Since 2009, 1,209 deer have been sampled for TB in Indiana.*

Bovine Tuberculosis (TB) is a chronic bacterial disease caused by the bacterium *Mycobacterium bovis*, and can affect nearly any mammal. *M. bovis* is most commonly transmitted by inhalation of aerosols or by ingestion. TB is most commonly maintained in cattle, but several species can propagate the disease, and are classified as reservoir hosts. These species make eradication of the disease difficult. In Michigan, white-tailed deer appear to be reservoir hosts, and significantly complicate eradication efforts.

In 2009, deer in a captive cervid farm in Franklin County tested positive for TB. Investigations conducted by the Indiana Board of Animal Health (BOAH) determined that two additional farms obtained cervids from the Franklin County farm, one each in Harrison and Wayne Counties. Further testing within these herds and subsequent depopulation found that the disease had not spread to other captive animals. Results from sampling free range deer in 2009 failed to detect the presence of TB. Still, the Division of Fish and Wildlife (DFW), the BOAH, and the United States Department of Agriculture (USDA) decided to continue its efforts to collect tissue samples from free ranging deer in Franklin and Fayette counties in 2011 to monitor if the disease had spread to wild deer.

Additionally, a cattle farm in northern Dearborn County had 4 cattle test positive for the same strain of TB in 2011. Additional cattle were determined to be positive after the herd was depopulated. Due to this significant finding, surveillance efforts were expanded to include Dearborn County beginning in 2012.



These management notes are issued periodically to provide a quick source of information on wildlife surveys and investigations, and various wildlife programs prior to more terminal reports. Any information provided is subject to further analysis and therefore is not for publication without permission.

In 2013 DFW staff manned 1 check station in Franklin, Fayette, and Ripley counties, as well as 3 check stations in Dearborn County during the opening weekend of firearms season, to voluntarily collect deer heads from hunters. Tissue samples were prepared for submission to Purdue's Animal Disease and Diagnostic Lab (ADDL) and the National Veterinary Services Laboratory (NVSL) in Ames, Iowa by BOAH and USDA personnel. A total of 102 deer heads were collected, with 16 coming from Dearborn County, 20 coming from Fayette County, 64 coming from Franklin County, and 2 coming from just outside of the targeted surveillance area.

Analysis of the samples by either the ADDL or NVSL indicated that TB was not been detected in any of the collected deer. Hunters in these regions are still encouraged to report any harvested deer that exhibit symptoms of TB (white lesions on the internal organs or ribcage of harvested deer) to their local district wildlife biologist or the Indiana Board of Animal Health. Since 2009, 1,309 deer have been specifically sampled for TB in Indiana, with no positives detected.