


POTENTIOMETRIC SURFACE MAP OF THE BEDROCK AQUIFERS OF HENRY COUNTY, INDIANA



Henry County, Indiana is located in the east-central portion of the state and is within the boundaries of three river basins. The northern area of the county is within the White and West Fork White River Basin; to the east is the Whitewater River Basin; and the remainder and largest part of the county, in the East Fork White River Basin.



The potentiometric surface mapped (PSM) contour elevations represent lines of equal elevation relative to the measured groundwater levels in wells. In general, wells completed in a confined aquifer system are bound by impermeable layers and will have static water levels under hydrostatic pressure causing the water level to rise above the elevation of the aquifer resource. In contrast, an unconfined aquifer system is not bound by impermeable layers; therefore, the water level will not be under hydrostatic pressure and will not rise above the aquifer resource. The potentiometric contour lines crossing through Westwood Park Reservoir represent the potentiometric surface of the groundwater in the immediate area, not the water level of the reservoir, which is a man made feature.

Static water level measurements in individual wells used to construct the potentiometric surface map are indicative of the water level at the time of well completion. Therefore, current site specific conditions may differ due to local or seasonal variations in measured static water levels.

Coordinate locations of water well records were physically obtained in the field, determined through address geocoding, or reported on water well records. Elevation data were obtained from a digital elevation model (DEM). Elevation and location quality assurance procedures were utilized to refine or remove data where errors were readily apparent.

In Henry County wells producing from bedrock are extremely limited with much of the county lacking in data and/or covered by more prolific unconsolidated deposits that limit the necessity to complete wells in bedrock. Therefore, potentiometric surface elevation contours have not been extended through the majority of the county. Depth to bedrock generally ranges from 10 to 345 feet with wells completed in carbonate deposits of the Silurian and Devonian Carbonates Aquifer System or, to a much lesser extent, shale with interbedded limestone of the Ordovician Maquoctus Group. There are 169 located wells that are completed in bedrock and utilized towards the mapping of the bedrock potentiometric surface.

Potentiometric surface elevations range from an isolated high of 1020 feet mean sea level (msl) along the north-central part county, to a low of 880 feet msl in the southwest part of the county along a portion of the Big Blue River. Generalized groundwater flow direction for the county is towards major drainage relevant to the basin. However, due to limited data, potentiometric surface contours are limited to isolated sections along the Big Blue River in the East Fork White River Basin; and, along Buck Creek and Fall Creek in the White and West Fork White River Basin.



- EXPLANATION**
- 800 — Line of equal elevation, in feet above mean sea level
 - Potentiometric Contour interval 10 feet
 - Stream
 - County Road
 - State Road
 - US Highway
 - Interstate
 - Basin Boundary
 - Municipal Boundary
 - Lake & River
 - State Managed Property
 - No Aquifer Material or Limited Data

Map Use and Disclaimer Statement
We request that the following agency be acknowledged in products derived from this map: Indiana Department of Natural Resources, Division of Water.
This map was compiled by staff of the Indiana Department of Natural Resources, Division of Water using data believed to be reasonably accurate. However, a degree of error is inherent in all maps. This product is distributed "as is" without warranties of any kind, either expressed or implied. This map is intended for use only at the published scale.

This map is created from several existing shapefiles. Township and Range Lines of Indiana (line shapefile, 20020621), Land Survey Lines of Indiana (polygon shapefile, 20020621), and County Boundaries of Indiana (polygon shapefile, 20020621), are all from the Indiana Geological Survey and based on a 1:24,000 scale. Roads (TIGER and INDOT) (line shapefile, 2005) is from the Indiana Department of Transportation and based on a 1:100,000 scale. System1 (line shapefile, 2003) is from the Indiana Department of Transportation and based on a 1:24,000 scale. Basin boundaries are modified from Watershed Boundary Dataset (polygon shapefile, 2008) from the Natural Resource Conservation Service and based on a 1:24,000 scale. Populated Areas in Indiana 2000 (polygon shapefile, 20021000) is from the U.S. Census Bureau and based on a 1:100,000 scale. Managed Lands IDNR IN (polygon shapefile, 20100920) is from the Indiana Department of Natural Resources and based on a 1:24,000 scale. Hydrography, Streams (NH) (line shapefile, 20081218), Rivers/NHD (polygon shapefile, 20081218), and Lakes (NHD) (polygon shapefile, 20081218) are from the U.S. Geological Survey and based on a 1:24,000 scale. Digital Elevation Hillshade image is derived from the Indiana OrthoLiDAR Statewide Collection Program (2011). Henry County Bedrock No Aquifer Material or Limited Data (polygon shapefile, Maier, 2014) and Potentiometric Surface Contours of the Bedrock Aquifers of Henry County, Indiana (line shapefile, Maier, 2014) are based on a 1:24,000 scale.

Potentiometric Surface Map of the
Bedrock Aquifers of Henry County, Indiana
by
Randal D. Maier
Division of Water, Resource Assessment Section
May 2014

Map generated by Joel Sanderson and Randy Maier
IDNR, Division of Water, Resource Assessment Section