



Indiana 2011 and United States 2011 Results

High School Youth Risk Behavior Survey						
Question	Indiana 2011	United States 2011	p-value	Indiana 2011 More Likely Than United States 2011	United States 2011 More Likely Than Indiana 2011	No Difference
Unintentional Injuries and Violence						
Rarely or never wore a bicycle helmet (among students who had ridden a bicycle during the 12 months before the survey)	93.3 (90.5–95.3) 1,931 [†]	87.5 (85.0–89.7) 9,526	0.00	●		
Rarely or never wore a seat belt (when riding in a car driven by someone else)	8.6 (7.0–10.6) 2,842	7.7 (6.5–9.1) 14,940	0.39			●
Rode with a driver who had been drinking alcohol one or more times (in a car or other vehicle during the 30 days before the survey)	21.7 (18.7–25.0) 2,853	24.1 (22.9–25.3) 14,908	0.15			●
Drove when drinking alcohol one or more times (a car or other vehicle during the 30 days before the survey)	5.3 (4.1–6.8) 2,795	8.2 (7.6–8.8) 15,090	0.00		●	
Carried a weapon on school property on at least 1 day (for example, a gun, knife, or club during the 30 days before the survey)	3.7 (2.9–4.8) 2,818	5.4 (4.7–6.1) 14,926	0.01		●	
Did not go to school because they felt unsafe at school or on their way to or from school on at least 1 day (during the 30 days before the survey)	4.9 (3.2–7.6) 2,854	5.9 (5.1–6.9) 15,320	0.37			●
Threatened or injured with a weapon on school property one or more times (for example, a gun, knife, or club during the 12 months before the survey)	6.8 (4.8–9.5) 2,851	7.4 (6.8–8.1) 15,344	0.58			●
In a physical fight on school property one or more times (during the 12 months before the survey)	8.9 (7.3–10.6) 2,829	12.0 (11.3–12.8) 15,182	0.00		●	
Bullied on school property (during the 12 months before the survey)	25.0 (22.3–27.9) 2,795	20.1 (18.7–21.5) 14,695	0.00	●		
Felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	29.1 (26.3–31.9) 2,797	28.5 (27.2–29.7) 15,269	0.69			●

Seriously considered attempting suicide (during the 12 months before the survey)	18.9 (15.8–22.5) 2,847	15.8 (15.1–16.5) 15,293	0.06			●
Made a plan about how they would attempt suicide (during the 12 months before the survey)	13.6 (11.9–15.5) 2,840	12.8 (12.0–13.6) 15,278	0.40			●
Attempted suicide one or more times (during the 12 months before the survey)	11.0 (8.9–13.4) 2,401	7.8 (7.1–8.5) 13,514	0.01	●		
Suicide attempt resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	3.9 (3.2–4.9) 2,412	2.4 (2.0–2.9) 12,959	0.00	●		
Carried a weapon on at least 1 day (for example, a gun, knife, or club during the 30 days before the survey)	17.0 (14.2–20.2) 2,803	16.6 (15.4–18.0) 15,024	0.84			●
Carried a gun on at least 1 day (during the 30 days before the survey)	4.6 (3.6–5.9) 2,817	5.1 (4.6–5.7) 14,507	0.40			●
In a physical fight one or more times (during the 12 months before the survey)	29.0 (26.3–31.8) 2,784	32.8 (31.5–34.1) 15,106	0.01		●	
Injured in a physical fight one or more times (injuries had to be treated by a doctor or nurse, during the 12 months before the survey)	3.7 (2.7–5.1) 2,839	3.9 (3.5–4.4) 14,570	0.68			●
Hit, slapped, or physically hurt on purpose by their boyfriend or girlfriend (during the 12 months before the survey)	11.3 (9.0–14.0) 2,824	9.4 (8.6–10.3) 15,270	0.15			●
Ever physically forced to have sexual intercourse (when they did not want to)	9.8 (8.1–11.9) 2,727	8.0 (7.3–8.8) 15,199	0.07			●
Ever been electronically bullied (including through e-mail, chat rooms, instant messaging, Web sites, or texting during the 12 months before the survey)	18.7 (16.4–21.2) 2,845	16.2 (15.3–17.2) 13,877	0.05			●
Tobacco Use						
Ever tried cigarette smoking (even one or two puffs)	49.5 (45.9–53.0) 2,721	44.7 (42.3–47.2) 13,359	0.03	●		
Smoked a whole cigarette for the first time before age 13 years	11.1 (9.6–12.9) 2,678	10.3 (9.3–11.5) 14,845	0.40			●
Smoked cigarettes on at least 1 day (during the 30 days before the survey)	18.1 (15.9–20.4) 2,721	18.1 (16.7–19.5) 14,743	0.99			●
Smoked cigarettes on 20 or more days (during the 30 days before the survey)	8.1 (6.7–9.8) 2,721	6.4 (5.8–7.1) 14,743	0.05			●
Smoked more than 10 cigarettes per day (among students who currently smoked cigarettes, on the days they smoked during the 30 days before the survey)	7.7 (5.1–11.4) 474	7.8 (6.3–9.7) 2,390	0.94			●
Smoked cigarettes on school property on at least 1 day (during the 30 days before the survey)	4.5 (3.5–5.8) 2,809	4.9 (4.4–5.4) 14,488	0.52			●
Ever smoked at least one cigarette every day for 30 days	13.8 (11.7–16.3) 2,727	10.2 (9.2–11.2) 13,940	0.00	●		

Did not try to quit smoking cigarettes (among students who currently smoked cigarettes, during the 12 months before the survey)	43.2 (36.4–50.3) 473	50.1 (47.0–53.1) 2,428	0.07			●
Usually obtained their own cigarettes by buying them in a store or gas station (among the students who were aged <18 years and who currently smoked cigarettes, during the 30 days before the survey)	13.9 (10.1–19.0) 378	14.0 (11.5–16.9) 1,835	0.98			●
Used chewing tobacco, snuff, or dip on at least 1 day (during the 30 days before the survey)	8.2 (7.2–9.3) 2,820	7.7 (6.6–9.0) 14,967	0.52			●
Used chewing tobacco, snuff, or dip on school property on at least 1 day (during the 30 days before the survey)	4.4 (3.5–5.6) 2,820	4.8 (4.0–5.9) 13,912	0.57			●
Smoked cigars, cigarillos, or little cigars on at least 1 day (during the 30 days before the survey)	14.6 (12.6–16.9) 2,840	13.1 (12.2–14.1) 14,566	0.18			●
Smoked cigarettes; smoked cigars, cigarillos, or little cigars; or used chewing tobacco, snuff, or dip on at least 1 day (during the 30 days before the survey)	24.5 (21.8–27.3) 2,695	23.4 (21.8–25.1) 13,890	0.51			●
Alcohol and Other Drug Use						
Ever had at least one drink of alcohol on at least 1 day (during their life)	70.4 (66.3–74.1) 2,778	70.8 (69.0–72.5) 14,168	0.84			●
Drank alcohol for the first time before age 13 years (other than a few sips)	17.6 (15.6–19.8) 2,825	20.5 (19.2–21.8) 15,094	0.02		●	
Had at least one drink of alcohol on at least 1 day (during the 30 days before the survey)	33.4 (30.2–36.9) 2,614	38.7 (37.2–40.3) 13,943	0.00		●	
Had five or more drinks of alcohol in a row within a couple of hours on at least 1 day (during the 30 days before the survey)	19.8 (17.0–22.9) 2,787	21.9 (21.0–22.8) 14,956	0.16			●
Usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol during the 30 days before the survey)	39.0 (36.2–41.9) 827	40.0 (37.5–42.5) 4,992	0.61			●
Had at least one drink of alcohol on school property on at least 1 day (during the 30 days before the survey)	2.0 (1.4–2.9) 2,801	5.1 (4.5–5.8) 15,033	0.00		●	
Ever used marijuana one or more times (during their life)	37.2 (33.5–41.1) 2,786	39.9 (37.8–42.1) 14,925	0.21			●
Tried marijuana for the first time before age 13 years	6.9 (5.6–8.6) 2,795	8.1 (7.3–9.0) 15,031	0.17			●
Used marijuana one or more times (during the 30 days before the survey)	20.0 (17.8–22.4) 2,789	23.1 (21.5–24.7) 14,970	0.03		●	
Used marijuana on school property one or more times (during the 30 days before the survey)	3.3 (2.2–5.0) 2,816	5.9 (5.1–6.7) 14,638	0.00		●	
Ever used any form of cocaine one or more times (for example, powder, crack, or freebase, during their life)	5.6 (4.1–7.7) 2,820	6.8 (6.2–7.5) 15,138	0.20			●

Used any form of cocaine one or more times (for example, powder, crack, or freebase, during the 30 days before the survey)	2.3 (1.7–3.2) 2,818	3.0 (2.6–3.5) 14,435	0.10			●
Ever sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high one or more times (during their life)	10.6 (8.1–13.6) 2,838	11.4 (10.7–12.1) 14,556	0.55			●
Ever used heroin one or more times (also called "smack", "junk", or "China white", during their life)	2.8 (1.7–4.5) 2,833	2.9 (2.5–3.3) 14,069	0.91			●
Ever used methamphetamines one or more times (also called "speed", "crystal", "crank", or "ice", during their life)	3.9 (2.3–6.5) 2,789	3.8 (3.4–4.3) 14,933	0.93			●
Ever used ecstasy one or more times (also called "MDMA", during their life)	6.3 (4.6–8.5) 2,833	8.2 (7.2–9.4) 14,270	0.07			●
Ever took steroid pills or shots without a doctor's prescription one or more times (during their life)	3.4 (2.4–4.8) 2,840	3.6 (3.2–4.1) 14,824	0.73			●
Ever took prescription drugs one or more times without a doctor's prescription (such as Oxycontin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, during their life)	21.4 (18.3–24.8) 2,834	20.7 (19.2–22.2) 14,011	0.67			●
Ever used a needle to inject any illegal drug into their body one or more times (during their life)	2.1 (1.3–3.2) 2,837	2.3 (1.9–2.7) 14,062	0.68			●
Offered, sold, or given an illegal drug by someone on school property (during the 12 months before the survey)	28.3 (25.6–31.1) 2,824	25.6 (23.6–27.6) 15,146	0.11			●
Sexual Behaviors						
Ever had sexual intercourse	51.0 (46.4–55.5) 2,513	47.4 (45.0–49.9) 14,344	0.17			●
Had sexual intercourse for the first time before age 13 years	5.2 (4.0–6.8) 2,523	6.2 (5.6–6.9) 14,335	0.17			●
Had sexual intercourse with four or more persons (during their life)	16.8 (14.1–20.0) 2,511	15.3 (14.2–16.4) 14,284	0.30			●
Had sexual intercourse with at least one person (during the 3 months before the survey)	38.5 (34.4–42.8) 2,516	33.7 (31.8–35.7) 14,296	0.04	●		
Drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	19.9 (15.8–24.8) 843	22.1 (20.6–23.6) 5,110	0.35			●
Did not use a condom during last sexual intercourse (among students who were currently sexually active)	42.2 (37.5–47.2) 840	39.8 (37.1–42.5) 5,035	0.37			●
Did not use birth control pills before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)	73.2 (66.2–79.2) 828	82.0 (79.8–84.1) 4,922	0.01		●	
Were never taught in school about AIDS or HIV infection	10.4 (8.9–12.1) 2,792	16.0 (14.4–17.7) 14,497	0.00		●	

Did not use Depo-Provera (or any injectable birth control), Nuva Ring (or any birth control ring), implanon (or any implant), or any IUD use before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)	92.6 (89.7–94.8) 828	94.7 (93.6–95.6) 4,922	0.12			●
Did not use birth control pills or Depo-Provera (or any injectable birth control), Nuva Ring (or any birth control ring), implanon (or any implant), or any IUD before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)	65.8 (58.7–72.3) 828	76.7 (74.3–79.0) 4,922	0.00		●	
Did not use both a condom during last sexual intercourse and birth control pills or Depo-Provera (or any injectable birth control), Nuva Ring (or any birth control ring), implanon (or any implant), or any IUD before last sexual intercourse (to prevent pregnancy, among students who were currently sexually active)	82.5 (76.5–87.1) 821	90.5 (89.1–91.8) 4,898	0.00		●	
Did not use any method to prevent pregnancy during last sexual intercourse (among students who were currently sexually active)	13.0 (9.6–17.5) 828	12.9 (11.6–14.2) 4,922	0.93			●
Dietary Behaviors						
Did not drink 100% fruit juices (during the 7 days before the survey)	23.1 (20.8–25.7) 2,816	19.2 (18.0–20.5) 14,748	0.00	●		
Did not eat fruit (during the 7 days before the survey)	13.0 (11.2–15.0) 2,833	11.7 (10.8–12.6) 14,699	0.20			●
Did not eat green salad (during the 7 days before the survey)	41.1 (38.9–43.4) 2,830	38.0 (36.1–39.9) 14,761	0.03	●		
Did not eat potatoes (excluding French fries, fried potatoes, or potato chips, during the 7 days before the survey)	32.1 (29.5–34.7) 2,825	31.2 (30.0–32.6) 14,512	0.57			●
Did not eat carrots (during the 7 days before the survey)	56.7 (53.3–60.0) 2,833	52.0 (49.9–54.1) 14,511	0.02	●		
Did not eat other vegetables (excluding green salad, potatoes, or carrots, during the 7 days before the survey)	17.3 (15.4–19.5) 2,832	17.0 (15.8–18.2) 14,724	0.77			●
Drank a can, bottle, or glass of soda or pop (not counting diet soda or diet pop, during the 7 days before the survey)	78.4 (75.5–81.1) 2,820	79.1 (77.6–80.5) 14,304	0.69			●
Drank a can, bottle, or glass of soda or pop one or more times per day (not counting diet soda or diet pop, during the 7 days before the survey)	28.5 (24.8–32.4) 2,820	27.8 (25.6–30.1) 14,304	0.75			●
Drank a can, bottle, or glass of soda or pop two or more times per day (not counting diet soda or diet pop, during the 7 days before the survey)	19.2 (17.0–21.6) 2,820	19.0 (16.9–21.3) 14,304	0.91			●

Drank a can, bottle, or glass of soda or pop three or more times per day (not counting diet soda or diet pop, during the 7 days before the survey)	10.7 (9.0–12.5) 2,820	11.3 (9.9–12.9) 14,304	0.58			●
Did not eat fruit or drink 100% fruit juices (during the 7 days before the survey)	6.5 (5.0–8.4) 2,809	4.8 (4.3–5.4) 14,654	0.06			●
Ate fruit or drank 100% fruit juices less than one time per day (during the 7 days before the survey)	44.7 (41.7–47.8) 2,809	36.0 (34.3–37.8) 14,654	0.00	●		
Ate fruit or drank 100% fruit juices less than two times per day (during the 7 days before the survey)	75.4 (73.5–77.3) 2,809	66.0 (64.4–67.7) 14,654	0.00	●		
Ate fruit or drank 100% fruit juices less than three times per day (during the 7 days before the survey)	85.5 (83.9–87.0) 2,809	77.6 (76.2–79.0) 14,654	0.00	●		
Did not eat vegetables (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	6.1 (5.0–7.4) 2,810	5.7 (5.1–6.4) 14,441	0.58			●
Ate vegetables less than one time per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	42.0 (40.0–43.9) 2,810	37.7 (36.0–39.5) 14,441	0.00	●		
Ate vegetables less than two times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	78.2 (76.4–79.8) 2,810	71.7 (70.3–73.1) 14,441	0.00	●		
Ate vegetables less than three times per day (green salad, potatoes [excluding French fries, fried potatoes, or potato chips], carrots, or other vegetables, during the 7 days before the survey)	91.0 (89.7–92.1) 2,810	84.7 (83.6–85.7) 14,441	0.00	●		
Overweight (students who were \geq 85th percentile but $<$ 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts)	15.4 (13.5–17.5) 2,746	15.2 (14.4–16.1) 14,285	0.86			●
Obese (students who were \geq 95th percentile for body mass index, based on sex- and age-specific reference data from the 2000 CDC growth charts)	14.7 (13.1–16.6) 2,746	13.0 (11.7–14.4) 14,285	0.12			●
Described themselves as slightly or very overweight	31.8 (28.9–34.8) 2,831	29.2 (27.7–30.6) 15,140	0.11			●
Did not eat for 24 or more hours to lose weight or to keep from gaining weight (during the 30 days before the survey)	13.1 (11.1–15.4) 2,818	12.2 (11.3–13.1) 15,068	0.41			●
Took diet pills, powders or liquids to lose weight or to keep from gaining weight (without a doctor's advice, during the 30 days before the survey)	6.3 (5.3–7.6) 2,822	5.1 (4.5–5.6) 15,136	0.04	●		

Vomited or took laxatives to lose weight or to keep from gaining weight (during the 30 days before the survey)	6.7 (4.5–9.7) 2,811	4.3 (3.8–4.7) 15,101	0.06			●
Physical Activity						
Physically active at least 60 minutes per day on less than 5 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	56.5 (53.1–59.9) 2,805	50.5 (48.5–52.6) 15,048	0.00		●	
Did not attend physical education classes in an average week (when they were in school)	65.1 (58.3–71.3) 2,802	48.2 (42.4–54.0) 14,896	0.00		●	
Did not attend physical education classes 5 days in an average week (when they were in school)	80.3 (73.0–86.1) 2,802	68.5 (62.6–73.9) 14,896	0.01		●	
Did not play on sports teams (run by their school or community groups during the 12 months before the survey)	44.6 (40.3–48.9) 2,818	41.6 (39.3–44.0) 14,891	0.23			●
Watched television 3 or more hours per day (on an average school day)	27.0 (24.2–29.9) 2,821	32.4 (30.9–34.0) 15,010	0.00		●	
Used computers 3 or more hours per day (played video or computer games or used a computer for something that was not school work on an average school day)	29.0 (26.2–31.9) 2,818	31.1 (29.3–32.9) 14,997	0.21			●
Physically active at least 60 minutes per day on less than 7 days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	75.8 (73.1–78.4) 2,805	71.3 (69.7–72.9) 15,048	0.00		●	
Did not participate in at least 60 minutes of physical activity on any day (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	15.9 (13.2–18.9) 2,805	13.8 (12.8–14.8) 15,048	0.16			●
Other - Asthma						
Ever told by a doctor or nurse that they had asthma	23.7 (21.0–26.7) 2,812	23.0 (21.7–24.3) 14,340	0.62			●
Ever told by a doctor or nurse that they had asthma and still have asthma	11.3 (9.9–12.9) 2,793	11.9 (10.9–12.9) 14,186	0.53			●

Footnotes

†	Percentage, confidence interval, cell size
---	--

Application URL: <http://qanccd.cdc.gov/youthonline/App/Results.aspx?>

TT=G&OUT=0&SID=HS&QID=QQ&LID=IN&YID=2011&LID2=XX&YID2=2011&COL=&ROW1=&ROW2=&HT=QQ&LCT=&FS=1&FR=1&FG=1&FSL=&FRL=&FGL=&PV=&TST=True&C1=IN2011&C2=XX2011&QP=G&DP=1&VA=CI&CS=Y&SYID=&EYID=&SC=DEFAULT&SO=ASC&PF=1

Centers for Disease Control and Prevention 1600 Clifton Rd. Atlanta, GA 30333, USA
800-CDC-INFO (800-232-4636) TTY: (888) 232-6348, 24 Hours/Every Day - cdcinfo@cdc.gov

