


Floyd County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding Percentage by Source



Comparison of Land Use Across County



Sediment Reduced 6,361,500 lbs.

Enough to fill 32 train cars.



x 32

Nitrogen Reduced 5,528 lbs.

Enough to fill 6 8' truck beds.



x 6

Phosphorus Reduced 2,762 lbs.

Enough to fill 3 8' truck beds.



x 3

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	16	16	1,885,790	846	1,693
2014	26	29	8,208,330	3,673	7,350
2015	32	39	9,134,710	4,090	8,185
2016	38	48	10,665,920	4,792	9,589
2017	79	92	10,738,265	4,703	9,413
2018	86	109	11,650,485	5,122	10,250
2019	17	65	6,361,500	2,762	5,528

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA