



Blackford County Nutrient and Sediment Load




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 10,724,525 lbs.
Enough to fill 54 train cars.



x 54

Nitrogen Reduced 14,852 lbs.
Enough to fill 15 8' truck beds.



x 15

Phosphorus Reduced 7,437 lbs.
Enough to fill 7 8' truck beds.



x 7

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	7	7	102,375	73	146
2014	3	7	102,375	73	146
2015	18	22	1,050,410	777	1,552
2016	38	45	2,047,600	1,534	3,064
2017	44	55	4,876,885	3,191	6,369
2018	50	79	6,645,390	4,533	9,050
2019	68	98	10,724,525	7,437	14,852

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA