


Franklin County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding Percentage by Source




Comparison of Land Use Across County



Sediment Reduced 21,411,545 lbs.

Enough to fill 107 train cars.



x 107

Nitrogen Reduced 20,666 lbs.

Enough to fill 21 8' truck beds.



x 21

Phosphorus Reduced 10,302 lbs.

Enough to fill 10 8' truck beds.



x 10

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	78	78	9,296,670	4,353	8,707
2014	18	69	7,696,705	3,543	7,165
2015	63	118	10,896,525	5,102	10,284
2016	88	172	18,228,065	8,760	17,601
2017	107	215	22,751,640	11,002	22,088
2018	92	204	20,034,830	9,628	19,337
2019	107	246	21,411,545	10,302	20,666

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA