



Wabash County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 36,109,320 lbs.
Enough to fill 181 train cars.



x 181

Nitrogen Reduced 42,062 lbs.
Enough to fill 42 8' truck beds.



x 42

Phosphorus Reduced 21,007 lbs.
Enough to fill 21 8' truck beds.



x 21

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	552	552	61,338,600	37,188	74,332
2014	521	602	66,448,090	40,060	80,071
2015	435	538	56,987,995	34,475	69,044
2016	334	449	49,819,215	29,872	59,783
2017	195	356	35,814,995	21,164	42,375
2018	159	338	35,664,070	20,800	41,649
2019	201	408	36,109,320	21,007	42,062

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA