



Putnam County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding Percentage by Source



Comparison of Land Use Across County



Sediment Reduced 76,136,215 lbs.
Enough to fill 381 train cars.



x 381

Nitrogen Reduced 81,468 lbs.
Enough to fill 81 8' truck beds.



x 81

Phosphorus Reduced 39,863 lbs.
Enough to fill 40 8' truck beds.



x 40

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	154	154	12,589,305	6,509	13,722
2014	151	256	16,999,595	8,833	19,175
2015	147	360	20,940,800	11,097	23,950
2016	104	381	19,508,490	10,200	22,173
2017	90	443	25,078,415	13,205	28,181
2018	361	787	59,518,900	31,772	65,281
2019	351	914	76,136,215	39,863	81,468

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA