

STD MORBIDITY

Chlamydia, Gonorrhea and Syphilis Morbidity – January – December 2009*

Gender	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Female	16,146	75%	3,951	58%	16	11%
Male	5,515	25%	2,838	42%	136	89%
Unknown	98	<1%	23	<1%	0	0%
Total	21,759	100%	6,812	100%	152	100%
Race	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
White	8,229	42%	1,534	25%	75	49%
Black	8,438	40%	4,107	60%	62	40%
Amer. Indian	28	<1%	7	<1%	1	<1%
Asian	58	<1%	13	<1%	2	<1%
Multiple Races	74	0%	25	0%	1	<1%
All Others	4,234	17%	1,009	13%	6	<1%
Unknown	518	<1%	117	<1%	5	<1%
Total	21,759	100%	6,812	100%	152	100%
Ethnicity	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Hispanic	1,049	4%	181	2%	7	5%
Non-Hispanic	14,613	68%	5,021	74%	134	88%
Unknown	6,097	28%	1,610	24%	11	7%
Total	21,759	100%	6,812	100%	152	100%
Age	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
0-9	5	<1%	3	<1%	0	0%
10-14	124	1%	29	<1%	0	0%
15-19	6,597	30%	1,652	24%	5	3%
20-24	8,830	40%	2,496	37%	27	18%
25-29	3,395	15%	1,267	20%	29	19%
30-34	1,480	7%	626	9%	18	11%
35-39	655	3%	309	4%	25	12%
40-44	241	1%	175	2%	17	18%
45-54	176	1%	150	2%	25	16%
55-64	36	<1%	39	<1%	6	3%
65+	12	<1%	8	<1%	0	0%
Unknown	208	1%	58	1%	0	0%
Total	21,759	100%	6,812	100%	152	100%

Source: ISDH STD*MIS Database as of January 2, 2010

NOTE: *Not all percents may add to 100 due to rounding. Data is considered provisional and will be updated later in 2011