



Martin County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 34,298,930 lbs.
Enough to fill 171 train cars.



x 171

Nitrogen Reduced 32,658 lbs.
Enough to fill 33 8' truck beds.



x 33

Phosphorus Reduced 16,274 lbs.
Enough to fill 16 8' truck beds.



x 16

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	136	136	30,642,365	14,583	29,175
2014	46	83	16,626,200	8,068	16,138
2015	52	98	20,384,850	10,013	20,117
2016	11	69	13,300,880	6,686	13,460
2017	78	141	36,836,590	17,580	35,253
2018	70	128	18,699,470	9,000	18,104
2019	107	182	34,298,930	16,274	32,658

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA