



Henry County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 20,077,945 lbs.
Enough to fill 100 train cars.



x 100

Nitrogen Reduced 24,328 lbs.
Enough to fill 24 8' truck beds.



x 24

Phosphorus Reduced 12,176 lbs.
Enough to fill 12 8' truck beds.



x 12

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	73	73	7,439,225	4,353	8,699
2014	36	65	6,970,355	4,093	8,180
2015	61	90	9,954,835	5,974	11,944
2016	25	63	5,540,240	3,179	6,347
2017	60	107	13,989,550	8,421	16,827
2018	82	127	14,671,950	8,808	17,600
2019	113	160	20,077,945	12,176	24,328

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA