


Clark County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding Percentage by Source




Comparison of Land Use Across County



Sediment Reduced 21,270,690 lbs.

Enough to fill 106 train cars.



x 106

Nitrogen Reduced 21,346 lbs.

Enough to fill 21 8' truck beds.



x 21

Phosphorus Reduced 10,670 lbs.

Enough to fill 11 8' truck beds.



x 11

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	99	99	17,412,715	8,792	17,585
2014	82	106	17,031,720	8,425	16,852
2015	110	155	25,353,205	12,687	25,379
2016	39	98	20,010,490	9,689	19,383
2017	31	93	11,565,745	5,909	11,820
2018	80	133	17,269,300	8,459	16,925
2019	97	161	21,270,690	10,670	21,346

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA