



Allen County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 46,002,135 lbs.
Enough to fill 230 train cars.



x 230

Nitrogen Reduced 59,921 lbs.
Enough to fill 60 8' truck beds.



x 60

Phosphorus Reduced 29,971 lbs.
Enough to fill 30 8' truck beds.



x 30

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	110	110	11,022,195	6,899	13,853
2014	130	142	12,538,395	8,120	16,292
2015	329	345	24,600,390	16,443	32,922
2016	250	273	17,461,965	11,548	23,141
2017	568	595	31,614,120	20,631	41,303
2018	408	622	40,750,535	26,143	52,320
2019	541	759	46,002,135	29,971	59,921

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA