


Adams County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 38,192,375 lbs.
Enough to fill 191 train cars.



x 191

Nitrogen Reduced 53,214 lbs.
Enough to fill 53 8' truck beds.



x 53

Phosphorus Reduced 25,932 lbs.
Enough to fill 26 8' truck beds.



x 26

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	244	244	10,605,140	7,681	16,795
2014	117	206	10,357,940	6,940	15,322
2015	172	289	14,544,300	9,589	20,617
2016	168	312	15,722,170	10,431	22,300
2017	195	350	17,280,855	11,057	23,550
2018	359	564	27,878,790	18,582	38,531
2019	498	716	38,192,375	25,932	53,214

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA