



Fulton County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 6,413,380 lbs.

Enough to fill 32 train cars.



x 32

Nitrogen Reduced 7,320 lbs.

Enough to fill 7 8' truck beds.



x 7

Phosphorus Reduced 3,649 lbs.

Enough to fill 4 8' truck beds.



x 4

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	86	86	13,410,285	7,606	15,283
2014	182	190	34,618,550	18,749	37,635
2015	225	233	31,216,610	17,692	36,101
2016	55	65	5,033,455	2,774	5,567
2017	103	118	6,582,760	3,678	7,380
2018	83	123	7,787,750	4,408	8,852
2019	102	162	6,413,380	3,649	7,320

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA