


Tipton County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 7,239,820 lbs.
Enough to fill 36 train cars.



x 36

Nitrogen Reduced 9,334 lbs.
Enough to fill 9 8' truck beds.



x 9

Phosphorus Reduced 4,673 lbs.
Enough to fill 5 8' truck beds.



x 5

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	16	16	3,599,405	2,468	4,922
2014	40	42	6,602,095	4,299	8,580
2015	13	25	3,164,800	1,963	3,911
2016	27	43	5,274,320	3,388	6,758
2017	26	42	5,645,615	3,638	7,259
2018	29	44	5,345,430	3,408	6,808
2019	46	62	7,239,820	4,673	9,334

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA