


Pike County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 79,379,855 lbs.

Enough to fill 397 train cars.



x 397

Nitrogen Reduced 72,347 lbs.

Enough to fill 72 8' truck beds.



x 72

Phosphorus Reduced 36,164 lbs.

Enough to fill 36 8' truck beds.



x 36

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	114	114	39,169,990	17,526	35,069
2014	165	193	48,051,480	21,378	42,776
2015	168	204	73,016,540	31,822	63,668
2016	136	193	94,345,480	40,928	81,894
2017	147	212	73,085,605	32,454	64,929
2018	144	217	60,511,025	27,298	54,609
2019	196	304	79,379,855	36,164	72,347

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA