


Vigo County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source




Comparison of Land
Use Across County



Sediment Reduced 36,315,730 lbs.

Enough to fill 182 train cars.



x 182

Nitrogen Reduced 38,975 lbs.

Enough to fill 39 8' truck beds.



x 39

Phosphorus Reduced 19,528 lbs.

Enough to fill 20 8' truck beds.



x 20

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	124	124	17,518,525	9,161	18,318
2014	43	115	13,777,450	7,191	14,376
2015	85	181	22,524,705	12,024	24,039
2016	26	124	14,577,535	7,599	15,192
2017	52	158	19,474,525	10,399	20,765
2018	177	302	33,277,100	18,060	36,042
2019	183	361	36,315,730	19,528	38,975

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA