


Kosciusko County Nutrient and Sediment Load




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 37,122,125 lbs.
Enough to fill 186 train cars.



x 186

Nitrogen Reduced 40,253 lbs.
Enough to fill 40 8' truck beds.



x 40

Phosphorus Reduced 20,085 lbs.
Enough to fill 20 8' truck beds.



x 20

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	672	672	62,270,510	33,502	67,211
2014	592	631	64,728,945	34,765	69,739
2015	328	393	35,638,225	20,143	40,435
2016	385	462	52,098,530	28,663	57,523
2017	484	581	58,045,785	31,579	63,361
2018	389	515	45,817,455	24,757	49,654
2019	274	462	37,122,125	20,085	40,253

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA