


Harrison County Nutrient and Sediment Load




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source




Comparison of Land
Use Across County



Sediment Reduced 108,228,270

Enough to fill 541 train cars.



x 541

Nitrogen Reduced 99,198 lbs.

Enough to fill 99 8' truck beds.



x 99

Phosphorus Reduced 48,361 lbs.

Enough to fill 48 8' truck beds.



x 48

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	177	177	41,022,695	17,608	35,237
2014	183	227	49,950,855	21,258	42,544
2015	267	325	72,418,155	31,802	66,040
2016	425	564	153,298,850	69,006	140,488
2017	320	547	146,933,600	66,769	136,022
2018	411	752	164,808,500	74,622	151,741
2019	271	727	108,228,270	48,361	99,198

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA