


Steuben County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source




Comparison of Land
Use Across County



Sediment Reduced 62,537,925 lbs.

Enough to fill 313 train cars.



x 313

Nitrogen Reduced 66,944 lbs.

Enough to fill 67 8' truck beds.



x 67

Phosphorus Reduced 33,466 lbs.

Enough to fill 33 8' truck beds.



x 33

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	131	131	32,398,750	17,157	34,311
2014	86	96	20,371,925	10,700	21,399
2015	114	133	33,330,665	17,643	35,284
2016	114	140	29,999,060	16,097	32,200
2017	80	111	20,379,890	10,891	21,794
2018	55	92	12,127,940	6,432	12,877
2019	299	349	62,537,925	33,466	66,944

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA