

Appendix M. Drift Plains Region SGCN Habitat and Subhabitat Tables

Table L-1. SGCN Occurring in Agricultural Systems in the Drift Plains Region.

| Taxa | Group | Scientific Name | Common Name |
|-------------------|----------------|-------------------------------|-------------------------|
| Amphibians | Frogs | <i>Lithobates areolatus</i> | Crawfish Frog |
| Birds | Waterfowl | <i>Cygnus buccinator</i> | Trumpeter Swan |
| | Quail & Grouse | <i>Colinus virginianus</i> | Northern Bobwhite |
| | Cranes | <i>Grus canadensis</i> | Sandhill Crane |
| | | <i>Grus americana</i> | Whooping Crane |
| | Shorebirds | <i>Pluvialis dominica</i> | American Golden-plover |
| | | <i>Bartramia longicauda</i> | Upland Sandpiper |
| | | <i>Calidris subruficollis</i> | Buff-breasted Sandpiper |
| | | <i>Tringa solitaria</i> | Solitary Sandpiper |
| | Raptors | <i>Circus cyaneus</i> | Northern Harrier |
| | | <i>Tyto alba</i> | Barn Owl |
| | | <i>Asio flammeus</i> | Short-eared Owl |
| | Songbirds | <i>Lanius ludovicianus</i> | Loggerhead Shrike |
| | | <i>Cistothorus platensis</i> | Sedge Wren |
| | Bears | <i>Ursus Americana</i> | Black Bear |
| Mammals | Mustelids | <i>Mustela nivalis</i> | Least Weasel |
| | | <i>Taxidea taxus</i> | American Badger |
| Reptiles | Turtles | <i>Terrapene carolina</i> | Eastern Box Turtle |

Table M-2. SGCN Occurring in Aquatic Systems in the Drift Plains Region.

| Taxa | Group | Scientific Name | Common Name | Big Rivers | Medium Rivers | Low Gradient | Moderate Gradient | High Gradient | Pools | Riffles | Springs/ Spring Brooks | Creeks | Shallow Water Lakes | Deep Water Lakes | Lake Michigan | Comments | |
|-------------------|---------------------|---------------------------|---------------------------------------|------------------------|---------------|--------------|-------------------|---------------|-------|---------|------------------------------|--------|------------------------|---------------------|------------------|------------------------------|----------------------|
| Amphibians | Aquatic Salamanders | <i>Necturus maculosus</i> | Common Mudpuppy | X | X | | | | | | | X | X | X | | Requires silt-free areas for | |
| | Salamanders | <i>Ambystoma barbouri</i> | Streamside Salamander | | | | | | X | | | X | | | | | |
| | Frogs | <i>Acris blanchardi</i> | Blanchard's Cricket Frog | | | X | | | X | | | X | X | | | | |
| | | | <i>Lithobates areolatus</i> | Crawfish Frog | | | | | | | | | X | | | | |
| Birds | Waterfowl | <i>Cygnus buccinator</i> | Trumpeter Swan | | | | | | | | | | X | | | Can use impoundments. | |
| | Rails | <i>Gallinula galeata</i> | Common Gallinule | | | | | | | | | | X | | | | |
| | Cranes | <i>Grus canadensis</i> | Sandhill Crane | | | X | X | | | | | | X | | | | |
| | | | <i>Grus americana</i> | Whooping Crane | | | | | | | | | X | | | | |
| | Shorebirds | <i>Charadrius melodus</i> | Piping Plover | | | | | | | | | | | | X | | |
| | | | <i>Arenaria interpres</i> | Ruddy Turnstone | | | | | | | | | | X | | X | |
| | | | <i>Limnodromus griseus</i> | Short-billed Dowitcher | | | | | | | | | | X | | X | |
| | | | <i>Tringa solitaria</i> | Solitary Sandpiper | | X | X | | | | | | X | X | | | |
| | | | <i>Tringa melanoleuca</i> | Greater Yellowlegs | | | | | | | | | | X | | | |
| | | | <i>Phalaropus tricolor</i> | Wilson's Phalarope | | | | | | | | | | X | | | |
| | | Terns | <i>Sternula antillarum athalassos</i> | Interior Least Tern | X | X | | | | | | | | X | X | | Can use borrow pits. |

| Taxa | Group | Scientific Name | Common Name | Big Rivers | Medium Rivers | Low Gradient | Moderate Gradient | High Gradient | Pools | Riffles | Springs/Spring Brooks | Creeks | Shallow Water Lakes | Deep Water Lakes | Lake Michigan | Comments |
|-------------|-----------|---------------------------------|----------------------------|------------|---------------|--------------|-------------------|---------------|-------|---------|-----------------------|--------|---------------------|------------------|---------------|---------------------------------------|
| | | <i>Chlidonias niger</i> | Black Tern | X | X | X | | | | | | | X | X | | Can use impoundments. |
| | Herons | <i>Botaurus lentiginosus</i> | American Bittern | | | | | | | | | | X | | | Can use impoundments. |
| | | <i>Ixobrychus exilis</i> | Least Bittern | | | | | | | | | | X | | | |
| | | <i>Ardea alba</i> | Great Egret | X | X | X | X | | X | X | X | X | X | | X | |
| | | <i>Nycticorax nycticorax</i> | Black-crowned Night-heron | | | X | X | | X | | | | X | | X | |
| | | <i>Nyctanassa violacea</i> | Yellow-Crowned Night-heron | | | | | | | | | | X | | | |
| | Raptors | <i>Pandion haliaetus</i> | Osprey | X | X | | X | X | | | | | X | X | X | Can use impoundments and borrow pits. |
| | | <i>Ictinia mississippiensis</i> | Mississippi Kite | X | X | | X | X | | | | | X | X | | |
| | | <i>Haliaeetus leucocephalus</i> | Bald Eagle | X | X | | X | X | | | | | X | X | X | Can use impoundments and borrow pits. |
| | | <i>Falco peregrinus</i> | Peregrine Falcon | | | | | | | | | | | | X | Can use impoundments. |
| | Songbirds | <i>Cistothorus palustris</i> | Marsh Wren | | | | | | | | | | X | | | |
| Fish | Eels | <i>Anguilla rostrata</i> | American Eel | X | X | X | X | X | | | | | X | X | X | |
| | Catfish | <i>Noturus stigmosus</i> | Northern Madtom | | X | X | X | | | X | | | X | | | |
| | Perches | <i>Etheostoma variatum</i> | Variagate Darter | | X | | | X | X | X | | | X | | | |
| | Perches | <i>Percina copelandi</i> | Channel Darter | | X | X | X | | X | X | | | X | X | | |

| Taxa | Group | Scientific Name | Common Name | Big Rivers | Medium Rivers | Low Gradient | Moderate Gradient | High Gradient | Pools | Riffles | Springs/ Spring Brooks | Creeks | Shallow Water Lakes | Deep Water Lakes | Lake Michigan | Comments |
|-----------------|---------------|---------------------------------------|---------------------------|------------|---------------|--------------|-------------------|---------------|-------|---------|------------------------------|--------|------------------------|---------------------|------------------|----------|
| | Trout-perches | <i>Percopsis omiscomaycus</i> | Trout-perch | X | X | | X | | X | | | X | X | X | X | |
| Mollusks | Mussels | <i>Plethobasus cyphus</i> | Sheepnose | X | X | X | X | | | X | | | | | | |
| | | <i>Pleurombema cordatum</i> | Ohio Pigtoe | X | X | X | X | | | X | | | X | | | |
| | | <i>Ptychobranhus fasciolaris</i> | Kidneyshell | X | X | | X | X | | X | | X | X | X | | |
| | | <i>Simpsonaias ambigua</i> | Salamander Mussel | X | X | | X | | | | | X | X | | | |
| | | <i>Toxolasma lividum</i> | Purple Lilliput | | X | X | | X | | X | | X | | | | |
| | | <i>Villosa lienosa</i> | Little Spectaclecase | X | X | X | X | | X | X | | X | | | | |
| Reptiles | Snakes | <i>Nerodia erythrogaster neglecta</i> | Copper-bellied Watersnake | | X | X | | | | | | X | X | | | |

Table M-3. SGCN Occurring in Barren Lands in the Drift Plains Region.

| Taxa | Group | Scientific Name | Common Name | Sand/ Dunes | Cliffs/Rock Outcrops | Bare Rock/ Talus | Comments |
|-------------------|------------|---|--------------------------|----------------|-------------------------|---------------------|--------------------------------------|
| Amphibians | Frogs | <i>Acris blanchardi</i> | Blanchard's Cricket Frog | | | X | Can use quarries. |
| Birds | Shorebirds | <i>Pluvialis dominica</i> | American Golden-plover | X | | | |
| | | <i>Charadrius melodus</i> | Piping Plover | X | | | |
| | Terns | <i>Sternula antillarum athalassos</i> | Interior Least Tern | X | | | |
| | Raptors | <i>Haliaeetus leucocephalu</i> | Bald Eagle | | X | | |
| | | <i>Falco peregrinus</i> | Peregrine Falcon | | X | | |
| Reptiles | Snakes | <i>Crotalus horridus</i> | Timber Rattlesnake | | X | X | Requires rocky areas for hibernation |

Table M-4. SGCN Occurring in Developed Lands in the Drift Plains Region.

| Taxa | Group | Scientific Name | Common Name | Suburban Areas | Urban Areas |
|-------------------|-----------|-----------------------------|---------------------|----------------|-------------|
| Amphibians | Frogs | <i>Lithobates areolatus</i> | Crawfish Frog | X | |
| Birds | Nightjars | <i>Chordeiles minor</i> | Common Nighthawk | X | X |
| | Raptors | <i>Accipiter striatus</i> | Sharp-shinned Hawk | X | |
| | | <i>Falco peregrinus</i> | Peregrine Falcon | | X |
| Mammals | Bats | <i>Myotis lucifugus</i> | Little Brown Myotis | X | |
| | | <i>Myotis sodalis</i> | Indiana Myotis | X | |
| | | <i>Perimyotis subflavus</i> | Tri-colored Bat | X | |
| | | <i>Nycticeius humeralis</i> | Evening Bat | X | |
| | Bears | <i>Ursus Americana</i> | Black Bear | X | |
| | Mustelids | <i>Mustela nivalis</i> | Least Weasel | X | |
| Reptiles | Snakes | <i>Clonophis kirtlandii</i> | Kirtland's Snake | X | X |
| | | <i>Opheodrys aestivus</i> | Rough Greensnake | X | |

Table M-5. SGCN Occurring in Forests in the Drift Plains Region.

| Taxa | Group | Scientific Name | Common Name | Hardwood Forests | Conifer Forests | Mixed Forests | Hardwood Woodlands | Conifer Woodlands | Mixed Woodlands | |
|----------------------------------|----------------|---------------------------------|----------------------------|----------------------|-----------------|---------------|--------------------|-------------------|-----------------|---|
| Amphibians | Salamanders | <i>Ambystoma barbouri</i> | Streamside Salamander | X | | | | | | |
| | | <i>Lithobates areolatus</i> | Crawfish Frog | | | | | X | | |
| Birds | Quail & Grouse | <i>Bonasa umbellus</i> | Ruffed Grouse | X | X | X | X | X | X | |
| | Nightjars | <i>Antrostomus vociferus</i> | Eastern Whip-poor-will | X | | X | X | | X | |
| | Raptors | <i>Ictinia mississippiensis</i> | Mississippi Kite | X | X | X | X | X | X | |
| | | <i>Haliaeetus leucocephalus</i> | Bald Eagle | X | X | X | X | X | X | |
| | | <i>Accipiter striatus</i> | Sharp-shinned Hawk | X | X | X | X | X | X | |
| | | <i>Buteo platypterus</i> | Broad-winged Hawk | X | | X | X | | X | |
| | Songbirds | <i>Helmitheros vermivorum</i> | Worm-eating Warbler | X | | | X | | | |
| | | <i>Mniotilta varia</i> | Black-and-white Warbler | X | | X | X | | X | |
| | | <i>Setophaga citrina</i> | Hooded Warbler | X | | | X | | | |
| | | <i>Setophaga cerulea</i> | Cerulean Warbler | X | | X | X | | X | |
| | Mammals | Shrews | <i>Sorex hoyi</i> | American Pygmy Shrew | X | X | X | | | |
| | | Bats | <i>Myotis lucifugus</i> | Little Brown Myotis | X | | X | X | | X |
| <i>Myotis septentrionalis</i> | | | Northern Long-eared Myotis | X | X | X | X | X | X | |
| <i>Myotis sodalis</i> | | | Indiana Myotis | X | | X | X | | X | |
| <i>Lasionycteris noctivagans</i> | | | Silver-haired Bat | X | X | X | X | X | X | |
| <i>Perimyotis subflavus</i> | | | Tri-colored Bat | X | | X | X | | X | |
| <i>Nycticeius humeralis</i> | | | Evening Bat | X | | X | X | | X | |
| <i>Lasiurus borealis</i> | | | Eastern Red Bat | X | | X | X | | X | |
| <i>Lasiurus cinereus</i> | | | Hoary Bat | X | X | X | X | X | X | |
| <i>Corynorhinus rafinesquii</i> | | | Rafinesque's Big-eared Bat | X | | X | X | | X | |
| Bears | | | <i>Ursus Americana</i> | Black Bear | X | X | X | X | X | X |
| Mustelids | | <i>Mustela nivalis</i> | Least Weasel | X | | X | X | X | X | |

| Taxa | Group | Scientific Name | Common Name | Hardwood Forests | Conifer Forests | Mixed Forests | Hardwood Woodlands | Conifer Woodlands | Mixed Woodlands |
|-----------------|---------|---------------------------------------|----------------------------|------------------|-----------------|---------------|--------------------|-------------------|-----------------|
| Reptiles | Turtles | <i>Terrapene carolina</i> | Eastern Box Turtle | X | X | X | X | X | X |
| | Snakes | <i>Nerodia erythrogaster neglecta</i> | Copper-bellied Watersnake | | | | X | X | X |
| | | <i>Opheodrys aestivus</i> | Rough Greensnake | X | | | X | | |
| | | <i>Cemophora coccinea</i> | Scarletsnake | X | X | X | X | X | X |
| | | <i>Tantilla coronate</i> | Southeastern Crowned Snake | | | | X | X | X |
| | | <i>Crotalus horridus</i> | Timber Rattlesnake | X | X | X | X | X | X |

Table M-6. SGCN Occurring in Grasslands in the Drift Plains Region.

| Taxa | Group | Scientific Name | Common Name | Savannas | Shrublands | Herbaceous Grasslands | Old Fields (Early Successional) | Comments |
|-------------------|------------------------------|---------------------------------|--------------------------|----------|------------|-----------------------|----------------------------------|----------|
| Amphibians | Frogs | <i>Acris blanchardi</i> | Blanchard's Cricket Frog | X | | X | | |
| | | <i>Lithobates areolatus</i> | Crawfish Frog | X | | X | | |
| Birds | Gamebirds | <i>Colinus virginianus</i> | Northern Bobwhite | X | X | X | X | |
| | Nightjars | <i>Chordeiles minor</i> | Common Nighthawk | | | X | X | |
| | Cranes | <i>Grus canadensis</i> | Sandhill Crane | | | X | | |
| | | <i>Grus americana</i> | Whooping Crane | | | X | | |
| | Herons | <i>Botaurus lentiginosus</i> | American Bittern | | | X | | |
| | | <i>Ardea alba</i> | Great Egret | | | X | | |
| | Shorebirds | <i>Pluvialis dominica</i> | American Golden-plover | | | X | | |
| | | <i>Bartramia longicauda</i> | Upland Sandpiper | | | X | X | |
| | | <i>Calidris subruficollis</i> | Buff-breasted Sandpiper | | | X | | |
| | Raptors | <i>Ictinia mississippiensis</i> | Mississippi Kite | X | X | X | | |
| | | <i>Circus cyaneus</i> | Northern Harrier | | | X | X | |
| | | <i>Accipiter striatus</i> | Sharp-shinned Hawk | | | | X | |
| | | <i>Tyto alba</i> | Barn Owl | X | | X | X | |
| | | <i>Asio flammeus</i> | Short-eared Owl | X | X | X | X | |
| | Songbirds | <i>Lanius ludovicianus</i> | Loggerhead Shrike | X | X | X | X | |
| | <i>Cistothorus platensis</i> | Sedge Wren | | | X | X | | |
| | <i>Ammodramus henslowii</i> | Henslow's Sparrow | | | X | X | Can use Farm Bill Program lands. | |
| | <i>Vermivora chrysoptera</i> | Golden-winged Warbler | X | X | | X | | |

| Taxa | Group | Scientific Name | Common Name | Savannas | Shrublands | Herbaceous Grasslands | Old Fields (Early Successional) | Comments |
|-----------------|-----------|---------------------------------------|----------------------------|----------|------------|-----------------------|---------------------------------|--|
| Mammals | Shrews | <i>Sorex hoyi</i> | American Pygmy Shrew | | X | X | | |
| | Bats | <i>Myotis lucifugus</i> | Little Brown Myotis | X | | | | Can use recovering mine lands. |
| | | <i>Myotis septentrionalis</i> | Northern Long-eared Myotis | X | | | | |
| | | <i>Myotis sodalis</i> | Indiana Myotis | X | | | | Can use recovering mine lands. |
| | Mustelids | <i>Mustela nivalis</i> | Least Weasel | | X | X | X | Can use hay lands and recovering mine lands. |
| | | <i>Taxidea taxus</i> | American Badger | X | X | X | | Can use recovering mine lands and dunes. |
| Reptiles | Turtles | <i>Terrapene carolina</i> | Eastern Box Turtle | | | X | | |
| | Snakes | <i>Nerodia erythrogaster neglecta</i> | Copper-bellied Watersnake | | | X | | Can use recovering mine lands. |
| | | <i>Clonophis kirtlandii</i> | Kirtland's Snake | X | | X | X | |
| | | <i>Opheodrys aestivus</i> | Rough Greensnake | | X | X | X | Can use recovering mine lands. |
| | | <i>Cemophora coccinea</i> | Scarletsnake | | | X | | |
| | | <i>Crotalus horridus</i> | Timber Rattlesnake | | X | X | X | Can use recovering mine lands. |

Table M-7. SGCN Occurring in Subterranean Systems in the Drift Plains Region.

| Taxa | Group | Scientific Name | Common Name | Subaquatic | Subterrestrial | Comments |
|-----------------|--------|----------------------------------|----------------------------|------------|----------------|---------------------|
| Mammals | Bats | <i>Myotis grisescens</i> | Gray Myotis | | X | Requires wet caves. |
| | | <i>Myotis lucifugus</i> | Little Brown Myotis | | X | |
| | | <i>Myotis septentrionalis</i> | Northern Long-eared Myotis | | X | |
| | | <i>Myotis sodalis</i> | Indiana Myotis | | X | |
| | | <i>Lasiorycteris noctivagans</i> | Silver-haired Bat | | X | |
| | | <i>Perimyotis subflavus</i> | Tri-colored Bat | | | |
| | | <i>Corynorhinus rafinesquii</i> | Rafinesque's Big-eared Bat | | X | |
| Reptiles | Snakes | <i>Crotalus horridus</i> | Timber Rattlesnake | | X | |

Table M-8. SGCN Occurring in Wetlands in the Drift Plains Region.

| Taxa | Group | Scientific Name | Common Name | Bogs/Fens | Herbaceous Wetlands | Forested Wetlands | Shrub Wetlands | Ephemeral/Temporary Wetlands | Mudflats | Riparian Zones |
|-------------------|---------------------------------|---------------------------------------|---------------------------|-----------|---------------------|-------------------|----------------|------------------------------|----------|----------------|
| Amphibians | Salamanders | <i>Hemidactylium scutatum</i> | Four-toed Salamander | X | | X | X | X | | X |
| | Frogs | <i>Acris blanchardi</i> | Blanchard's Cricket Frog | X | X | | X | X | | X |
| | | <i>Lithobates areolatus</i> | Crawfish Frog | | X | | | X | | X |
| Birds | Waterfowl | <i>Cygnus buccinator</i> | Trumpeter Swan | | X | | | X | | X |
| | Rails | <i>Rallus elegans</i> | King Rail | | X | | X | | | |
| | | <i>Gallinula galeata</i> | Common Gallinule | | X | | | | | |
| | Cranes | <i>Grus canadensis</i> | Sandhill Crane | X | X | | | | | X |
| | | <i>Grus americana</i> | Whooping Crane | X | X | | | X | | |
| | Shorebirds | <i>Pluvialis dominica</i> | American Golden-plover | | X | | | X | X | X |
| | | <i>Charadrius melodus</i> | Piping Plover | | X | | | | X | X |
| | | <i>Arenaria interpres</i> | Ruddy Turnstone | | X | | | | X | X |
| | | <i>Calidris subruficollis</i> | Buff-breasted Sandpiper | | | | | X | X | X |
| | | <i>Limnodromus griseus</i> | Short-billed Dowitcher | | X | | | X | X | X |
| | | <i>Scolopax minor</i> | American Woodcock | X | X | X | X | X | | X |
| | | <i>Tringa solitaria</i> | Solitary Sandpiper | X | X | X | | X | X | X |
| | | <i>Tringa melanoleuca</i> | Greater Yellowlegs | | X | | | X | X | X |
| | | <i>Phalaropus tricolor</i> | Wilson's Phalarope | | X | | | X | X | X |
| | Terns | <i>Sternula antillarum athalassos</i> | Interior Least Tern | | X | | | | | X |
| | | <i>Chlidonias niger</i> | Black Tern | | X | | | | | X |
| | Herons | <i>Botaurus lentiginosus</i> | American Bittern | | X | | X | X | | X |
| | | <i>Ixobrychus exilis</i> | Least Bittern | X | X | | X | | | X |
| | | <i>Ardea alba</i> | Great Egret | | X | X | X | X | | X |
| | | <i>Nycticorax nycticorax</i> | Black-crowned Night-heron | | X | X | X | X | | X |
| | <i>Nyctanassa violacea</i> | Yellow-crowned Night-heron | | X | X | | | | X | |
| Raptors | <i>Pandion haliaetus</i> | Osprey | | X | X | | | | X | |
| | <i>Ictinia mississippiensis</i> | Mississippi Kite | | | X | | | | X | |

| Taxa | Group | Scientific Name | Common Name | Bogs/Fens | Herbaceous Wetlands | Forested Wetlands | Shrub Wetlands | Ephemeral/ Temporary Wetlands | Mudflats | Riparian Zones |
|----------------|-----------|--------------------------------------|----------------------------|-----------|---------------------|-------------------|----------------|-------------------------------|----------|----------------|
| | | <i>Haliaeetus leucocephalus</i> | Bald Eagle | | X | X | | | | X |
| | | <i>Circus cyaneus</i> | Northern Harrier | X | X | | | | | |
| | | <i>Accipiter striatus</i> | Sharp-shinned Hawk | | | | | | | X |
| | | <i>Buteo platypterus</i> | Broad-winged Hawk | | | X | | | | X |
| | | <i>Tyto alba</i> | Barn Owl | | X | | | | | X |
| | | <i>Asio flammeus</i> | Short-eared Owl | X | X | | | | | |
| | | <i>Falco peregrinus</i> | Peregrine Falcon | | X | | | | | X |
| | Songbirds | <i>Cistothorus platensis</i> | Sedge Wren | X | X | | | | | X |
| | | <i>Cistothorus palustris</i> | Marsh Wren | | X | | | | | |
| | | <i>Xanthocephalus xanthocephalus</i> | Yellow-headed Blackbird | | X | | | | | |
| | | <i>Vermivora chrysoptera</i> | Golden-winged Warbler | X | | X | | | | X |
| | | <i>Mniotilta varia</i> | Black-and-white Warbler | | | X | | | | X |
| | | <i>Setophaga cerulea</i> | Cerulean Warbler | | | X | | | | X |
| Mammals | Shrews | <i>Sorex hoyi</i> | American Pygmy Shrew | X | X | X | | | | X |
| | Bats | <i>Myotis grisescens</i> | Gray Myotis | | | | | | | X |
| | | <i>Myotis lucifugus</i> | Little Brown Myotis | | | X | | | | X |
| | | <i>Myotis septentrionalis</i> | Northern Long-eared Myotis | | | X | | | | X |
| | | <i>Myotis sodalis</i> | Indiana Myotis | | | X | | | | X |
| | | <i>Lasionycteris noctivagans</i> | Silver-haired Bat | | | | | | | X |
| | | <i>Perimyotis subflavus</i> | Tri-colored Bat | | | | | | | X |
| | | <i>Nycticeius humeralis</i> | Evening Bat | | | | | | | X |
| | | <i>Lasiurus borealis</i> | Eastern Red Bat | | | | | | | X |
| | | <i>Lasiurus cinereus</i> | Hoary Bat | | | | | | | X |
| | | <i>Corynorhinus rafinesquii</i> | Rafinesque's Big-eared Bat | | | X | | | | X |
| | Mustelids | <i>Mustela nivalis</i> | Least Weasel | | X | | | | | X |

| Taxa | Group | Scientific Name | Common Name | Bogs/Fens | Herbaceous Wetlands | Forested Wetlands | Shrub Wetlands | Ephemeral/ Temporary Wetlands | Mudflats | Riparian Zones |
|-----------------|--------|---------------------------------------|---------------------------|-----------|---------------------|-------------------|----------------|-------------------------------|----------|----------------|
| Reptiles | Snakes | <i>Nerodia erythrogaster neglecta</i> | Copper-bellied Watersnake | X | X | X | | X | | X |
| | | <i>Clonophis kirtlandii</i> | Kirtland's Snake | X | X | | X | X | | X |
| | | <i>Opheodrys aestivus</i> | Rough Greensnake | | | X | X | | | X |
| | | <i>Cemophora coccinea</i> | Scarletsnake | | | | | | | X |
| | | <i>Crotalus horridus</i> | Timber Rattlesnake | | | X | | | | X |