	Food A	llergy Action Plan		
Student's Name:		D.O.B:Teach	er:	Place
ALLERGY	TO:			Child's Picture
Asthmatic Y	Yes* No *	Higher risk for severe reaction		Here
	♦ STF	P 1: TREATMENT ◆		,
Symptoms:			Give Checked Medication**: (To be determined by physician authorizing treatment)	
 If a food a 	allergen has been ingested, but	no symptoms:	☐ Epinephrine	☐ Antihistamine
Mouth	Itching, tingling, or swellin	g of lips, tongue, mouth	☐ Epinephrine	☐ Antihistamine
Skin	Hives, itchy rash, swelling	of the face or extremities	☐ Epinephrine	☐ Antihistamine
Gut	Nausea, abdominal cramps	, vomiting, diarrhea	☐ Epinephrine	☐ Antihistamine
■ Throat†	Tightening of throat, hoars	eness, hacking cough	☐ Epinephrine	☐ Antihistamine
■ Lung†	Shortness of breath, repetit	ive coughing, wheezing	☐ Epinephrine	☐ Antihistamine
 Heart† 	Thready pulse, low blood p	ressure, fainting, pale, blueness	☐ Epinephrine	☐ Antihistamine
Other†			☐ Epinephrine	☐ Antihistamine
 If reaction 	is progressing (several of the	above areas affected), give	☐ Epinephrine	☐ Antihistamine
Antihistamiı	ne: give			
Othom sive		medication/dose/route		
Other: give_		medication/dose/route		
	◆ <u>S</u>	ΓΕΡ 2: EMERGENCY CAL	<u>LS</u> ♦	
1. Call 911 (c may be nee	or Rescue Squad:) . State that an a	allergic reaction has been	treated, and additional epinephri
2. Dr		at		
3. Emergency Name/Relation		Phone Number(s)		
a		1.)	2.)	
b		1.)	2.)	
c		1.)	2.)	
EVEN IF PARE	ENT/GUARDIAN CANNOT BE R	EACHED, DO NOT HESITATE TO M	EDICATE OR TAKE CHII	D TO MEDICAL FACILITY!
Parent/Guardia	an Signature		Date	
Doctor's Signa			Date	
· ·	(Required)			

TRAINED STAFF MEMBERS				
1	Room			
2	Room			
3	Room			

EpiPen® and EpiPen® Jr. Directions

Pull off gray activation cap.



 Hold black tip near outer thigh (always apply to thigh).



Swing and jab firmly into outer thigh until Auto-Injector mechanism functions. Hold in place and count to 10. Remove the EpiPen® unit and massage the injection area for 10 seconds. Twinject™ 0.3 mg and Twinject™ 0.15 mg Directions



- Pull off green end cap, then red end cap.
- Put gray cap against outer thigh, press down firmly until needle penetrates. Hold for 10 seconds, then remove.



SECOND DOSE ADMINISTRATION:

If symptoms don't improve after 10 minutes, administer second dose:

- Unscrew gray cap and pull syringe from barrel by holding blue collar at needle base.
- Slide yellow or orange collar off plunger.
- Put needle into thigh through skin, push plunger down all the way, and remove.



Once EpiPen® or Twinject™ is used, call the Rescue Squad. Take the used unit with you to the Emergency Room. Plan to stay for observation at the Emergency Room for at least 4 hours.

For children with multiple food allergies, consider providing separate Action Plans for different foods.



**Medication checklist adapted from the Authorization of Emergency Treatment form developed by the Mount Sinai School of Medicine. Used with permission.