

# STD MORBIDITY

## Chlamydia, Gonorrhea and Syphilis Morbidity – January – December 2015

Gender	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Female	20,385	70.6%	3,984	50.8%	28	9.8%
Male	8,492	29.4%	3,854	49.1%	257	90.2%
Unknown	9	<1%	5	<1%	0	0%
Total	28,886	100%	7,843	100%	285	100%
Race	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
White	13,357	46.2%	2,832	36.1%	148	51.9%
Black	10,163	35.2%	4,127	52.6%	104	36.5%
All Others	5,366	18.6%	884	11.3%	33	11.6%
Unknown	0	0%	0	0%	0	0%
Total	28,886	100%	7,843	100%	285	100%
Ethnicity	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
Hispanic	1,645	5.7%	258	3.3%	26	8.7%
Non-Hispanic	27,241	94.3%	7,585	96.7%	272	91.3%
Unknown	0	0%	0	0%	0	0%
Total	28,886	100%	7,843	100%	285	100%
Age	Chlamydia		Gonorrhea		Syphilis (Primary and Secondary)	
0-19	8,214	28.4%	1,676	21.4%	11	3.9%
20-24	11,618	40.2%	2,783	35.5%	56	19.6%
25-29	5,004	17.3%	1,596	20.3%	66	23.2%
30-34	2,157	7.5%	798	10.1%	55	19.3%
35-39	993	3.4%	449	5.7%	36	12.6%
40-44	483	1.7%	240	3.1%	20	7.0%
45-54	319	1.1%	221	2.8%	26	9.1%
55+	98	<1%	80	1.1%	15	5.3%
Unknown	0	0%	0	0%	0	0%
Total	28,886	100%	7,843	100%	285	100%

Source: ISDH STD Database as of January 23, 2017

NOTE: Not all percents may add to 100 due to rounding.