

COVID-19 Vaccination Schedule

Pfizer

Pfizer (Most People)					
Age Group	6 months – 4 years	5-11 years	12 years +	Bivalent Booster 5-11 yrs	Bivalent Booster 12 years and older
Vial Cap Color/Label	Maroon Cap	Orange Cap	Gray Cap	Orange Cap Orange Border	Gray Cap
Description	Dilute Before Use 2.2 ml 10 doses per vial	Dilute Before Use 1.3 ml 10 doses per vial	Do Not Dilute 6 doses per vial	Dilute Before Use 1.3ml 10 doses per vial	Do Not Dilute Dose volume 0.3 ml 6 doses per vial
Primary Dose #1	3 mcg Dose volume 0.2 ml	10 mcg Dose volume 0.2 ml	30 mcg Dose volume 0.3ml	N/A	N/A
Primary Dose #2 3-8 weeks from 1 st primary dose	3 mcg Dose volume 0.2 ml	10 mcg Dose volume 0.2 ml	30 mcg Dose volume 0.3ml	N/A	N/A
Primary Dose #3	3 mcg (8 weeks from primary dose #2)	N/A	N/A	N/A	N/A
Bivalent Booster Only 1 bivalent booster at least 2 months after completion of primary series	N/A	1 Bivalent Booster 0.2 ml	1 Bivalent Booster 0.3 ml	0.2 ml	0.3 ml

Pfizer (Immunocompromised)					
Age Group	6 months – 4 years	5-11 years	12 years +	Bivalent Booster 5-11 yrs	Bivalent Booster 12 years and older
Vial Cap Color/Label	Maroon Cap	Orange Cap	Gray Cap	Orange Cap Orange Border	Gray Cap
Description	Dilute Before Use 2.2 ml 10 doses per vial	Dilute Before Use 1.3 ml 10 doses per vial	Do Not Dilute 6 doses per vial	Dilute Before Use 1.3ml 10 doses per vial	Do Not Dilute 6 doses per vial Dose volume 0.3 ml
Primary Dose #1	3 mcg Dose volume 0.2 ml	10 mcg Dose volume 0.2 ml	30 mcg Dose volume 0.3 ml	N/A	N/A
Primary Dose #2 3-8 weeks from 1 st primary dose	3 mcg Dose volume 0.2 ml	10 mcg Dose volume 0.2 ml	30 mcg Dose volume 0.3 ml	N/A	N/A
Primary Dose #3	3 mcg (0.2 ml) (8 weeks from primary dose #2)	10 mcg (0.2 ml) (4 weeks from primary dose #2)	30 mcg (0.3 ml) (at least 4 weeks from primary dose #2)	N/A	N/A
Bivalent Booster Only 1 bivalent booster at least 2 months after completion of primary series	N/A	1 Bivalent Booster 0.2 ml	1 Bivalent Booster 0.3 ml	0.2 ml	0.3 ml

Moderna

Moderna (Most People)					
Age Group	6 months – 5 years	6-11 years	12-17 years	18 years +	6 years + Bivalent Booster
Vial Cap Color Label	Blue Cap	Blue Cap	Red Cap	Red Cap	Blue Cap
Vial Label	Magenta Border	Purple Border	Blue Border	Blue Border	Gray Border
Description	Do Not Dilute 10 doses per vial	Do Not Dilute 10 doses per vial	Do Not Dilute 10 doses per vial	Do Not Dilute 10 doses per vial	Do Not Dilute 5 doses per vial
Primary Dose #1	25 mcg Dose volume 0.25 ml	50 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	N/A
Primary Dose #2 4-8 weeks from 1 st primary dose	25 mcg Dose volume 0.25 ml	50 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	N/A
Bivalent Booster Only 1 bivalent booster at least 2 months after completion of primary series	N/A	1 Bivalent Booster Dose 0.25 ml	1 Bivalent Booster Dose 0.5 ml	1 Bivalent Booster Dose 0.5 ml	6-11 yrs 0.25 ml 12 yrs + 0.5 ml

Moderna (Immunocompromised)					
Age Group	6 months – 5 years	6-11 years	12-17 years	18 years +	6 years + Bivalent Booster
Vial Cap Color Label	Blue Cap	Blue Cap	Red Cap	Red Cap	Blue Cap
Vial Label	Magenta Border	Purple Border	Blue Border	Blue Border	Gray Border
Description	Do Not Dilute 10 doses per vial	Do Not Dilute 10 doses per vial	Do Not Dilute 10-11 doses per vial	Do Not Dilute 10-11 doses per vial	Do Not Dilute 5 doses per vial
Primary Dose #1	25 mcg Dose volume 0.25 ml	50 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	N/A
Primary Dose #2 4 weeks from 1 st primary dose	25 mcg Dose volume 0.25 ml	50 mcg Dose volume 0.5 ml	50 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	N/A
Primary Dose #3 4 weeks from primary #2	25 mcg Dose volume 0.25 ml	50 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	100 mcg Dose volume 0.5 ml	N/A
Bivalent Booster Only 1 bivalent booster at least 2 months after completion of primary series	N/A	1 Bivalent Booster 0.25 ml	1 Bivalent Booster 0.5 ml	1 Bivalent Booster 0.5 ml	6-11 yrs 0.25 ml 12 yrs + 0.5 ml

Novavax		Monovalent Booster
Age Group	12 years and older	18 years +
Description	Do Not Dilute 10 doses per vial	Do Not Dilute 10 Doses per vial Dose 0.5 ml
Primary Dose #1	Dose volume 0.5 ml	N/A
Primary Dose #2 3 weeks from 1st primary dose	0.5 ml	N/A
Primary Dose #3	Not Authorized	N/A
Booster At least 6 weeks from Primary #2	N/A	Only if unable or unwilling to receive bivalent mRNA booster and if not previously received a booster

NOVAVAX	
Freezer -50°C to -15°C (-58°F to 5°F)	Do Not Freeze
Refrigerator 2°C to 8°C (36°F to 46°F)	Until date of expiration
Room Temperature 8°C to 25°C (46°F to 77°F)	Up to 6 Hours
After 1st Puncture	Discard after 6 hours

Pfizer/Moderna COVID-19 VACCINE STORAGE AND HANDLING

Pfizer				
Vial Cap Color/Label	Maroon Cap	Orange Cap	Gray Cap	Bivalent Booster
Ultra Low Freezer -90°C to -60°C (-130°F to -76°F)	Up to 12 Months	Up to 12 Months	Up to 12 Months	Up to 12 Months
Freezer (-25C -15C)	Do Not Store	Do Not Store	Do Not Store	Do Not Store
Refrigerator [2°C to 8°C (35°F to 46°F)]	10 Weeks	10 Weeks	10 Weeks	10 Weeks
Room Temperature 8°C to 25°C (46°F to 77°F)	12 Hours Prior to 1 st Puncture	12 Hours Prior to 1 st Puncture	12 Hours Prior to 1 st Puncture	12 Hours Prior to 1 st Puncture
After 1st Puncture	Discard After 12 Hours	Discard After 12 Hours	Discard After 12 Hours	Discard After 12 Hours

Moderna					
Vial Cap Color	Blue Cap	Blue Cap	Red Cap	Red Cap	Blue Cap
Label	Magenta Border	Purple Border	Blue Border	Blue Border	Gray Border
Freezer -50°C to -15°C (-58°F to 5°F)	Up to date of expiration	Up to date of expiration	Up to date of expiration	Up to date of expiration	Up to date of expiration
Refrigerator 2°C to 8°C (36°F to 46°F)	Up to 30 days	Up to 30 days	Up to 30 days	Up to 30 days	Up to 30 days
Room Temperature 8°C to 25°C (46°F to 77°F)	Total of 24 hours	Total of 24 hours	Total of 24 hours	Total of 24 hours	Total of 24 hours
After 1st Puncture	Discard after 12 hours	Discard after 12 hours	Discard after 12 hours	Discard after 12 hours	Discard after 12 hours

Monkeypox Vaccination Schedule

JYNNEOS		
Age	Under 18 years	18 years and Older
Route Dose Location	Subcutaneous (SQ) 0.5 ml Upper Arm	Intradermal (ID) 0.1 ml Forearm
Primary Dose #1	0.5 ml	0.1 ml
Primary Dose #2	At least 28 days from Primary Dose #1	At least 28 days from Primary Dose #1

Storage/Handling	
Freezer -25 to -15 C/ -13 to 5 F	Until Ready to Use
Refrigerator 2 to 8 C/ 35 to 46 F	Up to 12 Hours
Room Temperature 8 to 25 C/ 46 to 77 F	Up to 8 Hours after Initial Puncture