# COVID-19 Vaccination Schedule

## Pfizer

| Pfizer (Most People) | | | | | |
|---|---|---|---|---|---|
| **Age Group** | **6 months – 4 years** | **5-11 years** | **12 years +** | **Bivalent Booster 5-11 yrs** | **Bivalent Booster 12 years and older** |
| **Vial Cap Color/Label** | **Maroon Cap** | **Orange Cap** | **Gray Cap** | **Orange Cap Orange Border** | **Gray Cap** |
| **Description** | Dilute Before Use<br>2.2 ml<br>10 doses per vial | Dilute Before Use<br>1.3 ml<br>10 doses per vial | Do Not Dilute<br>6 doses per vial | Dilute Before Use<br>1.3ml<br>10 doses per vial | Do Not Dilute<br>Dose volume<br>0.3 ml<br>6 doses per vial |
| **Primary Dose #1** | 3 mcg<br>Dose volume<br>0.2 ml | 10 mcg<br>Dose volume<br>0.2 ml | 30 mcg<br>Dose volume<br>0.3ml | N/A | N/A |
| **Primary Dose #2**<br>***3-8 weeks from 1st primary dose*** | 3 mcg<br>Dose volume<br>0.2 ml | 10 mcg<br>Dose volume<br>0.2 ml | 30 mcg<br>Dose volume<br>0.3ml | N/A | N/A |
| **Primary Dose #3 (Bivalent)** | 3 mcg *(8 weeks from primary dose #2)* | N/A | N/A | N/A | N/A |
| **Bivalent Booster**<br>***Only 1 bivalent booster at least 2 months after completion of primary series*** | 1 Bivalent Booster<br>0.2 ml | 1 Bivalent Booster<br>0.2 ml | 1 Bivalent Booster<br>0.3 ml | 0.2 ml | 0.3 ml |

| Pfizer (Immunocompromised) | | | | | |
|---|---|---|---|---|---|
| **Age Group** | **6 months – 4 years** | **5-11 years** | **12 years +** | **Bivalent Booster 5-11 yrs** | **Bivalent Booster 12 years and older** |
| **Vial Cap Color/Label** | **Maroon Cap** | **Orange Cap** | **Gray Cap** | **Orange Cap Orange Border** | **Gray Cap** |
| **Description** | Dilute Before Use<br>2.2 ml<br>10 doses per vial | Dilute Before Use<br>1.3 ml<br>10 doses per vial | Do Not Dilute<br>6 doses per vial | Dilute Before Use<br>1.3ml<br>10 doses per vial | Do Not Dilute<br>6 doses per vial<br>Dose volume 0.3 ml |
| **Primary Dose #1** | 3 mcg<br>Dose volume<br>0.2 ml | 10 mcg<br>Dose volume<br>0.2 ml | 30 mcg<br>Dose volume<br>0.3 ml | N/A | N/A |
| **Primary Dose #2**<br>***3-8 weeks from 1st primary dose*** | 3 mcg<br>Dose volume<br>0.2 ml | 10 mcg<br>Dose volume<br>0.2 ml | 30 mcg<br>Dose volume<br>0.3 ml | N/A | N/A |
| **Primary Dose #3**<br>**Bivalent** | 3 mcg (0.2 ml) *(8 weeks from primary dose #2)* | 10 mcg (0.2 ml) *(4 weeks from primary dose #2)* | 30 mcg (0.3 ml) *(at least 4 weeks from primary dose #2)* | N/A | N/A |
| **Bivalent Booster**<br>***Only 1 bivalent booster at least 2 months after completion of primary series*** | 1 Bivalent Booster<br>0.2 ml | 1 Bivalent Booster<br>0.2 ml | 1 Bivalent Booster<br>0.3 ml | 0.2 ml | 0.3 ml |

## Moderna

| Moderna (Most People) | | | | | |
|---|---|---|---|---|---|
| **Age Group** | **6 months – 5 years** | **6-11 years** | **12-17 years** | **18 years +** | **6 years + Bivalent Booster** |
| **Vial Cap Color Label** | **Blue Cap** | **Blue Cap** | **Red Cap** | **Red Cap** | **Blue Cap** |
| **Vial Label** | **Magenta Border** | **Purple Border** | **Blue Border** | **Blue Border** | **Gray Border** |
| **Description** | Do Not Dilute<br>10 doses per vial | Do Not Dilute<br>10 doses per vial | Do Not Dilute<br>10 doses per vial | Do Not Dilute<br>10 does per vial | Do Not Dilute<br>5 doses per vial |
| **Primary Dose #1** | 25 mcg<br>Dose volume<br>0.25 ml | 50 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | N/A |
| **Primary Dose #2**<br>***4-8 weeks from 1st primary dose*** | 25 mcg<br>Dose volume<br>0.25 ml | 50 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | N/A |
| **Bivalent Booster**<br>*Only 1 bivalent booster at least 2 months after completion of primary series* | 1 Bivalent Booster<br>0.2 ml | 1 Bivalent Booster<br>Dose<br>0.25 ml | 1 Bivalent Booster<br>Dose<br>0.5 ml | 1 Bivalent Booster<br>Dose<br>0.5 ml | **6-11 yrs 0.25 ml**<br>**12 yrs + 0.5 ml** |

| Moderna (Immunocompromised) | | | | | |
|---|---|---|---|---|---|
| **Age Group** | **6 months – 5 years** | **6-11 years** | **12-17 years** | **18 years +** | **6 years + Bivalent Booster** |
| **Vial Cap Color Label** | **Blue Cap** | **Blue Cap** | **Red Cap** | **Red Cap** | **Blue Cap** |
| **Vial Label** | **Magenta Border** | **Purple Border** | **Blue Border** | **Blue Border** | **Gray Border** |
| **Description** | Do Not Dilute<br>10 doses per vial | Do Not Dilute<br>10 doses per vial | Do Not Dilute<br>10-11 doses per vial | Do Not Dilute<br>10-11 doses per vial | Do Not Dilute<br>5 doses per vial |
| **Primary Dose #1** | 25 mcg<br>Dose volume<br>0.25 ml | 50 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | N/A |
| **Primary Dose #2**<br>***4 weeks from 1st primary dose*** | 25 mcg<br>Dose volume<br>0.25 ml | 50 mcg<br>Dose volume<br>0.5 ml | 50 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | N/A |
| **Primary Dose #3**<br>***4 weeks from primary #2*** | 25 mcg<br>Dose volume<br>0.25 ml | 50 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | 100 mcg<br>Dose volume<br>0.5 ml | N/A |
| **Bivalent Booster**<br>*Only 1 bivalent booster at least 2 months after completion of primary series* | 1 Bivalent Booster<br>0.2 ml | 1 Bivalent Booster<br>0.25 ml | 1 Bivalent Booster<br>0.5 ml | 1 Bivalent Booster<br>0.5 ml | **6-11 yrs 0.25 ml**<br>**12 yrs + 0.5 ml** |

| Novavax | | Monovalent Booster |
|---|---|---|
| **Age Group** | **12 years and older** | **18 years +** |
| **Description** | Do Not Dilute<br>10 doses per vial | Do Not Dilute<br>10 Doses per vial<br>Dose 0.5 ml |
| **Primary Dose #1** | Dose volume<br>0.5 ml | N/A |
| **Primary Dose #2**<br>***3 weeks from 1st primary dose*** | 0.5 ml | N/A |
| **Primary Dose #3** | Not Authorized | N/A |
| **Booster**<br>**At least 6 weeks from Primary #2** | N/A | Only if unable or unwilling to receive bivalent mRNA booster and if not previously received a booster |

| NOVAVAX | |
|---|---|
| **Freezer** -50°C to -15°C (-58°F to 5°F) | Do Not Freeze |
| **Refrigerator** 2°C to 8°C (36°F to 46°F) | Until date of expiration |
| **Room Temperature** 8°C to 25°C (46°F to 77°F) | Up to 6 Hours |
| **After 1st Puncture** | Discard after 6 hours |

## Pfizer/Moderna COVID-19 VACCINE STORAGE AND HANDLING

| Pfizer | | | | |
|---|---|---|---|---|
| **Vial Cap Color/Label** | **Maroon Cap** | **Orange Cap** | **Gray Cap** | **Bivalent Booster** |
| **Ultra Low Freezer** -90ºC to -60ºC (-130ºF to -76ºF) | Up to 12 Months | Up to 12 Months | Up to 12 Months | Up to 18 Months |
| **Freezer (-25C -15C)** | Do Not Store | Do Not Store | Do Not Store | Do Not Store |
| **Refrigerator** [2ºC to 8ºC (35ºF to 46ºF) | 10 Weeks | 10 Weeks | 10 Weeks | 10 Weeks |
| **Room Temperature** 8°C to 25°C (46°F to 77°F) | 12 Hours Prior to 1st Puncture | 12 Hours Prior to 1st Puncture | 12 Hours Prior to 1st Puncture | 12 Hours Prior to 1st Puncture |
| **After 1st Puncture** | Discard After 12 Hours | Discard After 12 Hours | Discard After 12 Hours | Discard After 12 Hours |

| Moderna | | | | | |
|---|---|---|---|---|---|
| **Vial Cap Color** | **Blue Cap** | **Blue Cap** | **Red Cap** | **Red Cap** | **Blue Cap** |
| **Label** | **Magenta Border** | **Purple Border** | **Blue Border** | **Blue Border** | **Gray Border** |
| **Freezer** -50°C to -15°C (-58°F to 5°F) | Up to date of expiration | Up to date of expiration | Up to date of expiration | Up to date of expiration | Up to date of expiration |
| **Refrigerator** 2°C to 8°C (36°F to 46°F) | Up to 30 days | Up to 30 days | Up to 30 days | Up to 30 days | Up to 30 days |
| **Room Temperature** 8°C to 25°C (46°F to 77°F) | Total of 24 hours | Total of 24 hours | Total of 24 hours | Total of 24 hours | Total of 24 hours |
| **After 1st Puncture** | Discard after 12 hours | Discard after 12 hours | Discard after 12 hours | Discard after 12 hours | Discard after 12 hours |