Administration and Storage of Novavax

Primary Series

Given to ages 18 years and older

Dose 1--- 0.5 ml*

wait at least 3 weeks

Dose 2--- 0.5 ml*

Booster Administration

NOT AUTHORIZED

DO NOT FREEZE

Refrigerator (2 to 8 C/36 to 46 F):

Store in refrigerator. No expiry printed on vial.

Punctured vials may be stored for 6 hours.

Room Temp. (8 to 25 C/46 to 77 F):

Punctured vials may be stored for 6 hours.

Regardless of storage, discard punctured vials after 6 hours.

Storage

^{*} each 0.5 ml dose contains 5 mcg SARS-CoV-2rS and 50 mcg Matrix- \mathbf{M}^{TM} adjuvant

Administration and Storage of Novavax

Special Considerations

Mixed Dosing of Primary Series

If a product other than Novavax is administered as dose 2 in primary series, it is considered an error and must be reported to VAERS.

If a product other than Novavax is administered as dose 2 in primary series <u>due to a</u>

<u>contraindication</u>, it is not considered an error.

The second dose would be given at least 4 weeks from the first dose of Novavax.

Coadministration With Other Vaccines

Novavax may be given along with other vaccines if determined appropriate by the provider.

Consider waiting 4 weeks before receiving a Covid-19 vaccination if an orthopoxvirus vaccination has been given. If a Covid-19 vaccination has been given already, there is no need to wait before receiving and orthopoxvirus vaccination.