

Smoking During Pregnancy



Tobacco Prevention
and Cessation

March 2025

Everyone deserves the opportunity to be healthy, free from the harm commercial tobacco use can cause. Use of commercial tobacco products impacts even the youngest Indiana residents, as smoking during pregnancy can harm the health of both pregnant Hoosiers and their children.

Smoking during Pregnancy in Indiana and the United States

- In 2023, the rate of smoking during pregnancy among Indiana residents was reported as **5.3%** which is significantly lower compared to 2022, where **6.6%** was reported.*
- The percentage of Indiana residents who smoked during pregnancy has historically been higher than the national average. In 2022, the most recent year of data available, the U.S. rate of smoking during pregnancy was 3.7%.**
- In 2023, among counties with stable rates, county rates of smoking during pregnancy ranged from **2.3% (Hancock County)** to **17.8% (Switzerland County)**.
- Of Indiana's 92 counties, 24 had a smoking during pregnancy rate significantly higher than the statewide rate.

Fast Facts

In 2023, **5.3%** of Indiana residents smoked during pregnancy.

There were over **4,100 babies** born to Hoosiers who smoked during pregnancy in 2023.

Risks of Smoking during Pregnancy

Smoking can impact every phase of reproduction. When trying to become pregnant, smoking can cause problems with fertility.

During Pregnancy

Smoking can increase the risk of several pregnancy complications, including:

- Low birth weight/reduced growth
- Baby born too early (premature birth)
- Ectopic pregnancy

After Pregnancy

Exposure to secondhand smoke can increase the risk for additional complications for the baby, including:

- Sudden Infant Death Syndrome (SIDS)
- Health problems due to weaker lungs

With the transition to the new platform for recording vital records data, Database for Registering Indiana's Vital Records Events (DRIVE), in 2021, new questions were added to the Indiana birth worksheet pertaining to tobacco use. The added questions focused on use of different forms of tobacco, trying to quit tobacco during pregnancy, and how soon an individual smoked cigarettes after waking up. We attempted to calculate a more complete statistic for people who had smoked during pregnancy using new variables that had been added to our data. However, due to questions around quality and completeness, we determined that these newer variables are not ready for reporting. For this reason, we have decided to revert to our method of reporting before our DRIVE transition. This means that smoking during pregnancy is calculated using variables for each trimester indicating whether the person had smoked during that trimester.

*Smoking during pregnancy estimates are among people that had a live birth.

**The national smoking during pregnancy rate in 2022 (most recent year available) is based on data from all U.S. states and the District of Columbia. Due to changes in the collection and cleaning of Indiana birth data starting in 2021, Indiana smoking during pregnancy variables should not be directly compared to the U.S. estimate.

Benefits of Quitting Smoking

It is never too late to treat nicotine dependence. Many pregnant women are tempted to cut down the number of cigarettes they smoke, but quitting is the best thing to do for their pregnancy. The benefits of quitting smoking can be seen immediately:

- After one day of not smoking, the baby will get more oxygen and the mother will have more energy and be able to breathe more easily.
- Reduces many risks for the baby, including premature birth and low birth weight.
- Reduces the risk of diseases such as heart and lung disease, stroke and cancer.

Quit Now Indiana - Pregnancy Program

Quit Now Indiana (1-800-QUIT-NOW) offers free, evidence-based nicotine dependence treatment to Hoosiers who use commercial tobacco products. For those who are planning to get pregnant, are expecting, or have recently given birth, Quit Now Indiana offers a tailored cessation program including seven coaching sessions with treatment specialists trained to support the quit journey during pregnancy and postpartum to prevent relapse. For more information, visit QuitNowIndiana.com.

Percentage of Indiana residents who smoked during pregnancy, Indiana Counties, 2023*

County	%	County	%	County	%	County	%
Adams	3.7	Franklin	5.7 (U)	Lawrence	11.4 (H)	Rush	11.9 (H)
Allen	4.5	Fulton	10.8 (H)	Madison	8.9 (H)	Scott	10.2 (H)
Bartholomew	7.4	Gibson	5.3 (U)	Marion	3.2 (L)	Shelby	7.6
Benton	8.7 (U)	Grant	15.4 (H)	Marshall	4.9	Spencer	7.0 (U)
Blackford	15.8 (U)	Greene	11.1 (H)	Martin	7.3 (U)	St. Joseph	4.2
Boone	2.1 (U)	Hamilton	0.4 (U)	Miami	9.0	Starke	9.2
Brown	8.3 (U)	Hancock	2.3 (L)	Monroe	5.9	Steuben	8.5
Carroll	7.2 (U)	Harrison	7.7	Montgomery	9.0 (H)	Sullivan	9.3
Cass	5.6	Hendricks	2.4 (L)	Morgan	9.4 (H)	Switzerland	17.8 (H)
Clark	7.1	Henry	8.8	Newton	11.4 (U)	Tippecanoe	4.5
Clay	7.0	Howard	8.0	Noble	6.2	Tipton	4.1 (U)
Clinton	6.8	Huntington	6.6	Ohio	Suppressed	Union	8.8 (U)
Crawford	12.9 (U)	Jackson	7.4	Orange	8.8 (U)	Vanderburgh	6.6
Daviess	5.1	Jasper	8.5	Owen	13.2 (H)	Vermillion	9.2 (U)
Dearborn	8.0	Jay	7.7	Parke	5.4 (U)	Vigo	10.1 (H)
Decatur	11.5 (H)	Jefferson	13.0 (H)	Perry	11.8 (H)	Wabash	7.2
DeKalb	10.0 (H)	Jennings	9.5 (H)	Pike	Suppressed	Warren	8.9 (U)
Delaware	8.3 (H)	Johnson	3.9	Porter	4.0	Warrick	2.0 (U)
Dubois	3.3 (U)	Knox	10.3 (H)	Posey	6.5 (U)	Washington	9.6 (H)
Elkhart	3.1 (L)	Kosciusko	6.7	Pulaski	8.0 (U)	Wayne	9.1 (H)
Fayette	11.1 (H)	LaGrange	1.4 (U)	Putnam	7.2	Wells	7.4
Floyd	7.6	Lake	2.5 (L)	Randolph	10.0	White	9.9 (H)
Fountain	5.6 (U)	LaPorte	8.2 (H)	Ripley	8.8	Whitley	5.9

H=Significantly higher than the statewide rate, L=Significantly lower than the statewide rate, U=Rates based on counts less than 20 are considered unstable and should be interpreted with caution. Rates are suppressed if based on counts less than 5.

*Smoking during pregnancy estimates are among people that had a live birth.