



## INDIANA DEPARTMENT OF ENVIRONMENTAL MANAGEMENT

*We Protect Hoosiers and Our Environment.*

*Mitchell E. Daniels Jr.*  
Governor

*Thomas W. Easterly*  
Commissioner

100 North Senate Avenue  
Indianapolis, Indiana 46204  
(317) 232-8603  
Toll Free (800) 451-6027  
[www.idem.IN.gov](http://www.idem.IN.gov)

May 26, 2011

Ms. Susan Hedman  
Regional Administrator  
U.S. Environmental Protection Agency  
Region 5  
77 West Jackson Boulevard  
Chicago, IL 60604-3950

Dear Ms. Hedman:

Re: Request for Redesignation Petition and  
Maintenance Plan for the Indiana Portion  
(Lake and Porter Counties) of the Chicago-  
Gary-Lake IL-IN Nonattainment Area for  
Fine Particles

The Indiana Department of Environmental Management (IDEM) submits a Redesignation Petition and Maintenance Plan for the Indiana Portion (Lake and Porter counties) of the Chicago-Lake IL-IN Nonattainment Area, which was designated as nonattainment of the annual standard for fine particles on April 5, 2005. IDEM conducted a public hearing concerning the Redesignation Petition and Maintenance Plan on May 18, 2011, and the public comment period concluded on May 20, 2011.

This submittal documents the public review process, including a detailed summary of and response to substantive comments.

The attached document consists of the following:

### ***Redesignation Petition and Maintenance Plan***

- A formal request that the Indiana Portion (Lake and Porter counties) of the Chicago-Lake IL-IN Nonattainment Area be redesignated to attainment and reclassified as maintenance. It contains and meets the requirements set forth in Section 107 of the Clean Air Act and in United States Environmental Protection Agency (U.S. EPA) Redesignation Guidance.
- A maintenance year of 2025 is established and 2015 and 2020 are analyzed as interim years.
- The appendices of the document contain historical air quality trend data, projected emission inventory data, and thorough documentation of the mobile emissions analysis.

**Motor Vehicle Emissions Budgets**

- Contained in the Redesignation Petition is a new Motor Vehicle Emissions Budget for 2016 and 2025. The Northwestern Indiana Regional Planning Commission's (NIRPC's) travel demand forecasting model and U.S. EPA's software program referred to as MOBILE6.2 were used to determine emissions for the annual fine particle nonattainment area.
- A conservative margin of safety was applied to the 2016 and 2025 projected emissions.
- The travel demand model was updated with the best available assumptions.
- Vehicle registration data gathered from the Indiana Bureau of Motor Vehicles were used to replace the MOBILE6.2 default vehicle age distribution.

IDEM requests that U.S. EPA proceed with review and approval of this submittal. If you have any questions or need additional information, please contact Scott Deloney, Chief, Air Programs Branch, at (317) 233-5694.

Sincerely,



Keith Baugues  
Assistant Commissioner  
Office of Air Quality

KB/sad/ghf

Enclosure:

Request for Redesignation Petition and Maintenance Plan Under the Annual National Ambient Air Quality Standard for Fine Particles for the Indiana Portion of the Chicago-Gary-Lake County, IL-IN Nonattainment Area

Cc: John Summerhays, U.S. EPA Region 5 (w/ enclosures)  
Matt Rau, U.S. EPA Region 5 (w/ enclosures)  
John Mooney, U.S. EPA Region 5 (no enclosures)  
Cheryl Newton, U.S. EPA Region 5 (no enclosures)  
Pat Morris, U.S. EPA Region 5 (no enclosures)  
Bill Brown, NIRPC (w/ enclosures)  
Rob Kaleel, Illinois EPA (w/ enclosures)  
Scott Deloney, IDEM (no enclosures)  
Christine Pedersen, IDEM (no enclosures)  
Gale Ferris, IDEM (w/ enclosures)

REQUEST FOR REDESIGNATION AND  
MAINTENANCE PLAN  
UNDER THE ANNUAL NATIONAL  
AMBIENT AIR QUALITY  
STANDARD FOR FINE PARTICLES

For the Indiana Portion  
of the

Chicago-Gary-Lake County, IL-IN  
Nonattainment Area for Fine Particles

**Lake and Porter Counties, Indiana**

Prepared By:  
The Indiana Department of Environmental Management

May 2011

**This page intentionally left blank**

## TABLE OF CONTENTS

<b>1.0 INTRODUCTION</b>	<b>1</b>
1.1 Background	1
1.2 Geographical Description	3
1.3 Status of Air Quality	3
<b>2.0 REQUIREMENTS FOR REDESIGNATION</b>	<b>3</b>
2.1 General	3
2.2 Fine Particles Monitoring	3
2.3 Emission Inventory	4
2.4 Modeling Demonstration	4
2.5 Controls and Regulations	4
2.6 Corrective Actions for Potential Future Violations of the Fine Particle Standard	5
<b>3.0 FINE PARTICLES MONITORING</b>	<b>5</b>
3.1 Fine Particle Monitoring Network	5
3.2 Ambient Fine Particles Monitoring Data	6
3.3 Quality Assurance	14
3.4 Continued Monitoring	14
<b>4.0 EMISSION INVENTORY</b>	<b>14</b>
4.1 Emission Trends	14
4.2 Base Year Inventory	25
4.3 Emission Projections	26
4.4 Demonstration of Maintenance	32
4.5 Permanent and Enforceable Emission Reductions	33
4.6 Provisions for Future Updates	33
<b>5.0 TRANSPORTATION CONFORMITY BUDGETS</b>	<b>33</b>
5.1 On-Road Emission Estimates	33
5.2 Overview	33
5.3 Analysis Years	34
5.4 Emission Estimates	34
5.5 Motor Vehicle Emission Budget	35
5.6 Commitment to Amend Motor Vehicle Emission Budgets Using Motor Vehicle Emission Simulator (MOVES)	35

<b>6.0 CONTROL MEASURES AND REGULATIONS .....</b>	<b>36</b>
6.1 Reasonably Available Control Technology (RACT) .....	36
6.2 Implementation of Past State Implementation Plan (SIP) Revisions .....	36
6.3 Nitrogen Oxides (NO <sub>x</sub> ) Rule .....	36
6.4 Measures Beyond Clean Air Act SIP Requirements .....	37
6.5 Controls to Remain in Effect .....	41
6.6 New Source Review Provisions .....	41
<b>7.0 MODELING AND METEOROLOGY .....</b>	<b>42</b>
7.1 Summary of Modeling Results to Support Rulemakings .....	42
7.2 Lake Michigan Air Directors Consortium (LADCO) Round 5 Speciated Modeled Attainment Test Results .....	46
7.3 LADCO Round 5 Particle Source Apportionment Results .....	47
7.4 Summary of Existing Modeling Results .....	56
7.5 Meteorological Analysis for Lake and Porter Counties, Indiana .....	56
7.6 Surface Air Conditions Present During High Fine Particle Concentration Days .....	56
7.7 Upper Air Conditions Present During High Fine Particle Concentration Days .....	56
7.8 Analyses of Atmospheric Conditions During High Fine Particle Concentration Days .....	57
7.9 Summary of Air Quality Index Days in Lake and Porter Counties, Indiana .....	57
7.10 Summary of Meteorological Analysis for Lake and Porter Counties, Indiana .....	59
<b>8.0 CORRECTIVE ACTIONS .....</b>	<b>59</b>
8.1 Commitment to Revise Plan .....	59
8.2 Commitment for Contingency Measures .....	60
8.3 Contingency Measures .....	60
<b>9.0 PUBLIC PARTICIPATION .....</b>	<b>61</b>
<b>10.0 CONCLUSIONS .....</b>	<b>62</b>

## **FIGURES**

Figure 3.1 Chicago Nonattainment Area .....	6
---	---

## **TABLES**

Table 1.1 National Ambient Air Quality Standards for Fine Particles (PM <sub>2.5</sub> ) .....	2
Table 3.1 Monitoring Data for Indiana's Portion of the Chicago Area .....	7
Table 3.2 Monitoring Data for Illinois' Portion of the Chicago Area .....	8
Table 4.1 Comparison of 2008 Estimated and 2025 Projected Emission Estimates, Lake and Porter Counties, Indiana .....	32
Table 4.2 Comparison of 2008 Estimated and 2025 Projected Emission Estimates, Entire Chicago Area .....	32



Table 5.1	Emission Estimates for On-Road Mobile Sources .....	34
Table 5.2	Motor Vehicle Emission Budgets for Lake and Porter Counties .....	35
Table 6.1	Trends in Electric Generating Unit (EGU) NO <sub>x</sub> Emissions Statewide in Indiana .....	37
Table 7.1	Clean Air Transport Rule Modeling Results from U.S. EPA - 2010 .....	43
Table 7.2	LADCO Round 6 Modeling Results for Annual PM <sub>2.5</sub> – Without the Clean Air Interstate Rule Emission Reductions.....	45
Table 7.3	LADCO Round 5 Speciated Modeled Attainment Test Modeling Results for Lake and Porter Counties, Indiana .....	47
Table 7.4	Ranking and Percentage of Highest Number of Days at Air Quality Index Levels of Health Concern.....	59

## **GRAPHS**

Graph 3.1	Fine Particle Design Values for Indiana’s Portion of the Chicago Area, 2006 through 2010.....	10
Graph 3.2	Fine Particle Design Values for Illinois’ Portion of the Chicago Area, 2006 through 2010.....	11
Graph 3.3	Annual Fine Particle Trends for Indiana’s Portion of the Chicago Area, 2006 through 2010.....	12
Graph 3.4	Annual Fine Particle Trends for Illinois’ Portion of the Chicago Area, 2006 through 2010.....	13
Graph 4.1	NO <sub>x</sub> Point Source Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008 .....	15
Graph 4.2	Sulfur Dioxide (SO <sub>2</sub> ) Point Source Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008 .....	16
Graph 4.3	Direct PM <sub>2.5</sub> Point Source Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008 .....	16
Graph 4.4	NO <sub>x</sub> Point Source Emission Trends, Entire Chicago Area, 2002, 2005, and 2008.....	17
Graph 4.5	SO <sub>2</sub> Point Source Emission Trends, Entire Chicago Area, 2002, 2005, and 2008 .....	17
Graph 4.6	Direct PM <sub>2.5</sub> Point Source Emission Trends, Entire Chicago Area, 2002, 2005, and 2008 .....	18
Graph 4.7	NO <sub>x</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008 ...	19
Graph 4.8	SO <sub>2</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008 ....	20
Graph 4.9	Direct PM <sub>2.5</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008 .....	20
Graph 4.10	NO <sub>x</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008 .....	21
Graph 4.11	SO <sub>2</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008 .....	21
Graph 4.12	Direct PM <sub>2.5</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008.....	22
Graph 4.13	NO <sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana, 1999 to 2009 .....	23
Graph 4.14	Indiana Statewide NO <sub>x</sub> Emissions from EGUs, 1999 to 2009 .....	24
Graph 4.15	SO <sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana, 1999 to 2009 .....	24
Graph 4.16	Indiana Statewide SO <sub>2</sub> Emissions from EGUs, 1999 to 2009.....	25
Graph 4.17	Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO <sub>x</sub> Emissions, Lake and Porter Counties, Indiana.....	27
Graph 4.18	Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected SO <sub>2</sub> Emissions, Lake and Porter Counties, Indiana .....	28

Graph 4.19	Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected Direct PM <sub>2.5</sub> Emissions, Lake and Porter Counties, Indiana.....	28
Graph 4.20	Comparison of 2005 and 2008 Estimated and 2020, 2015, and 2025 Projected NO <sub>x</sub> , SO <sub>2</sub> , and Direct PM <sub>2.5</sub> Emissions, Lake and Porter Counties, Indiana.....	29
Graph 4.21	Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO <sub>x</sub> Emissions, Entire Chicago Area.....	29
Graph 4.22	Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected SO <sub>2</sub> Emissions, Entire Chicago Area.....	30
Graph 4.23	Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected Direct PM <sub>2.5</sub> Emissions, Entire Chicago Area.....	30
Graph 4.24	Comparison of 2005 and 2008 Estimated and 2015 and 2025 Projected NO <sub>x</sub> , SO <sub>2</sub> , and Direct PM <sub>2.5</sub> Emissions, Entire Chicago Area .....	31
Graph 7.1	PM <sub>2.5</sub> Design Value Trends for Lake and Porter Counties, Indiana, 2000 to 2010 ...	44
Graph 7.2	LADCO Round 6 Modeling Results for Lake and Porter Counties, Indiana for 2012 and 2018 .....	46

## **CHARTS**

Chart 7.1	Regional/Emission Sector Particulate Source Apportionment Technology (PSAT) Results.....	48
Chart 7.2	Modeled Contributions by Specie at the East Chicago – Franklin School PM <sub>2.5</sub> Monitor .....	52
Chart 7.3	Modeled Contributions by Specie at the Griffith PM <sub>2.5</sub> Monitor .....	52
Chart 7.4	Modeled Contributions by Specie at the Gary – Madison Street PM <sub>2.5</sub> Monitor .....	53
Chart 7.5	Modeled Contributions by Specie at the Gary – Ivanhoe School PM <sub>2.5</sub> Monitor .....	53
Chart 7.6	Modeled Contributions by Specie at the Hammond – Purdue PM <sub>2.5</sub> Monitor .....	54
Chart 7.7	Modeled Contributions by Specie at the Hammond – Clark High School PM <sub>2.5</sub> Monitor .....	54
Chart 7.8	Modeled Contributions by Specie at the Dunes National Lakeshore PM <sub>2.5</sub> Monitor .....	55
Chart 7.9	Modeled Contributions by Specie at the Ogden Dunes PM <sub>2.5</sub> Monitor .....	55
Chart 7.10	Distribution of PM <sub>2.5</sub> Concentration Days on the AQI Levels of Health Concern, Lake and Porter Counties, Indiana .....	58



## **APPENDICES**

- A Air Quality System (AQS) and Indiana Department of Environmental Management (IDEM) Fine Particle Monitor Data Values for the Chicago Area (2000 –2010)
- B NO<sub>x</sub>, SO<sub>2</sub>, and Direct PM<sub>2.5</sub> Point Source Emissions (2002, 2005, and 2008), Chicago Area
- C NO<sub>x</sub>, SO<sub>2</sub>, and Direct PM<sub>2.5</sub> Emission Trends, All Sources (2002, 2005, and 2008), Chicago Area
- D NO<sub>x</sub> and SO<sub>2</sub> Emission Trends from EGUs, Lake and Porter Counties, Indiana (1999 to 2009)
- E 2005 and 2008 Base Year Emission Inventories and 2015, 2020, and 2025 Projected Emission Inventories for NO<sub>x</sub>, SO<sub>2</sub>, and Direct PM<sub>2.5</sub>, Chicago Area
- F Example Mobile Source Input and Output Calculation Files for Lake and Porter Counties, Indiana
- G IDEM - Area Source Inventory Standard Operating Procedure
- H LADCO Emission Estimates Technical Support Document
- I LADCO Round 5 Modeling Technical Support Document (Round 5 Photochemical Modeling Based on “Base M” Emission Inventory, revised version of “Base K”)
- J Modeling Summary
- K Public Participation Process Documents

**This page intentionally left blank**

**REQUEST FOR REDESIGNATION AND MAINTENANCE PLAN  
UNDER THE ANNUAL NATIONAL AMBIENT AIR  
QUALITY STANDARD FOR FINE PARTICLES**

**CHICAGO-GARY-LAKE COUNTY IL-IN AREA**

**1.0 INTRODUCTION**

This document supports Indiana's request that Lake and Porter counties, Indiana, which are part of the Chicago-Gary-Lake County IL-IN nonattainment area for fine particles (herein referred to as the "Chicago Area"), be redesignated from nonattainment to attainment of the 1997 annual standard for fine particles. All monitors for fine particles in the Chicago Area have recorded three years of quality assured ambient air quality monitoring data for the years 2008 through 2010, demonstrating attainment with the annual standard for fine particles, therefore; Indiana's portion of the Chicago Area is eligible for redesignation.

Section 107 of the Clean Air Act (CAA) establishes specific requirements to be met in order for an area to be considered for redesignation, including:

- (a) A determination that the area has attained the annual standard for fine particles.
- (b) A State Implementation Plan (SIP) for the area under Section 110(k) is fully approved.
- (c) A determination that the improvement in air quality is due to permanent and enforceable reductions in emissions resulting from implementation of the SIP and other federal requirements.
- (d) A maintenance plan under Section 175A is fully approved.
- (e) A determination that all Section 110 and Part D requirements have been met.

A maintenance plan provides for the continued attainment of the air quality standard by an area for a period of ten years after the United States Environmental Protection Agency (U.S. EPA) has formally redesignated the area to attainment. The plan also provides assurances that even if there is a subsequent exceedance of the air quality standard, then measures in the maintenance plan will prevent any future occurrences through contingency measures that would be triggered.

This document addresses each of these requirements, and provides additional information to support continued compliance with the annual standard for fine particles.

**1.1 Background**

The CAA requires states with areas designated nonattainment of the applicable National Ambient Air Quality Standard (NAAQS) for fine particles to develop SIPs to expeditiously attain and maintain the standard. In 1997, U.S. EPA set daily and annual air quality standards for fine particles (PM<sub>2.5</sub>), as shown in Table 1.1. The terms "fine particles" and "PM<sub>2.5</sub>" are used synonymously throughout this document. The PM<sub>2.5</sub> standards were legally challenged and upheld by the U.S. Supreme Court in February of 2001. In 1999, the Indiana Department of

Environmental Management (IDEM) began monitoring for fine particle concentrations. U.S. EPA designated areas in Indiana under the annual fine particles standard on December 17, 2004, as attainment, nonattainment or unclassifiable, with an effective date of April 5, 2005.

**Table 1.1**  
**National Ambient Air Quality Standards for Fine Particles**

	<b>Annual</b>	<b>24-Hour</b>
1997 PM <sub>2.5</sub> Standards	<b>15 µg/m<sup>3</sup></b> Annual arithmetic mean, averaged over three years	<b>65 µg/m<sup>3</sup></b> 24-hour average, 98 <sup>th</sup> percentile, averaged over three years
2006 PM <sub>2.5</sub> Standards	<b>15 µg/m<sup>3</sup></b> Annual arithmetic mean, averaged over three years	<b>35 µg/m<sup>3</sup></b> 24-hour average, 98 <sup>th</sup> percentile, averaged over three years

Note: The Chicago Area meets the 1997 and 2006 24-hour NAAQS for fine particles. Since this area is solely designated nonattainment under the 1997 annual standard for fine particles, this document only addresses the annual standard.

On December 17, 2004, based on 2001 through 2003 monitoring data, U.S. EPA designated the Chicago Area as nonattainment of the annual standard for fine particles (40 CFR 81.315). The Chicago Area was subject to Section 172 of the CAA, including the development of a plan to reduce nitrogen oxides (NO<sub>x</sub>), sulfur dioxide (SO<sub>2</sub>), and direct fine particle (PM<sub>2.5</sub>) emissions and a demonstration that the area will meet the annual standard for fine particles by April 5, 2010. In order to satisfy these requirements, Indiana submitted a redesignation petition and maintenance plan to U.S. EPA on April 3, 2008, demonstrating that Lake and Porter counties, Indiana met the NAAQS for fine particles by April 5, 2010, with an ample margin of safety. The Chicago Area monitors have continued to meet the annual NAAQS for fine particles since the end of 2007.

There were no fine particle monitors in the Chicago Area that violated the 1997 24-hour standard for fine particles and none that currently violate the 2006 24-hour standard for fine particles. As a result, the Chicago Area was designated nonattainment for fine particles solely under the 1997 annual standard. Therefore, this document pertains only to the 1997 annual standard for fine particles.

The Chicago nonattainment area for fine particles, as defined in Section 1.2, has not previously been subject to nonattainment area rulemakings for fine particles. However, the Chicago Area had been subject to nonattainment area rulemakings under the 1-hour ozone standard. The 1-hour ozone standard was revoked on June 15, 2005. Lake and Porter counties had also been subject to nonattainment rulemakings under the 8-hour ozone standard and were redesignated to attainment and classified as maintenance under the 8-hour ozone standard on May 11, 2010. The Illinois portion of the Chicago Area 8-hour ozone nonattainment area remains designated as “nonattainment” under the 8-hour ozone standard.

## 1.2 Geographical Description

The Chicago nonattainment area for fine particles consists of Lake and Porter counties, Indiana and Cook, DuPage, Kane, Lake, McHenry, and Will counties, and portions of Grundy (Aux Sable and Goose Lake Townships) and Kendall (Oswego Township) counties, Illinois. The Chicago Area contains such cities as Gary, Hammond, East Chicago, Portage, and Valparaiso in Indiana and Chicago, Elgin, Aurora, and Joliet in Illinois. This area is depicted in Figure 3.1.

IDEM and Illinois Environmental Protection Agency (IEPA) are responsible for assuring the nonattainment area for fine particles complies with the CAA requirements. IDEM is responsible for Lake and Porter counties, Indiana, while IEPA is responsible for Cook, DuPage, Kane, Lake, McHenry, and Will Counties, and portions of Grundy (Aux Sable and Goose Lake Townships) and Kendall (Oswego Township) counties, Illinois. IDEM and IEPA have worked cooperatively with U.S. EPA Region V to address attainment planning issues.

Although the agencies have worked together on a comprehensive plan for the multi-state nonattainment areas, the State of Illinois is required to make a separate submittal for its portion of the planning components to U.S. EPA. As such, this submittal only covers Lake and Porter counties, Indiana.

## 1.3 Status of Air Quality

Monitoring data for fine particles for the three years, 2008 through 2010, demonstrates that air quality has met the annual NAAQS for fine particles in the Chicago Area. This fact, accompanied by the permanent and enforceable reductions in emission levels discussed in Section 4.0, justifies a redesignation to attainment for the area based on Section 107(d)(3)(E) of the CAA.

# 2.0 REQUIREMENTS FOR REDESIGNATION

## 2.1 General

Section 110 and Part D of the CAA list a number of requirements that must be met by nonattainment areas prior to consideration for redesignation to attainment. In addition, U.S. EPA has published detailed guidance in a document entitled "Procedures for Processing Requests to Redesignate Areas to Attainment", issued September 4, 1992, to Regional Air Directors. This document is hereafter referred to as "Redesignation Guidance". This Request for Redesignation and Maintenance Plan is based on the Redesignation Guidance, supplemented with additional guidance received from staff of the Attainment Planning and Maintenance Section of U.S. EPA Region V. The specific requirements for redesignation are listed below.

## 2.2 Fine Particles Monitoring

- 1) A demonstration that the annual standard for fine particles, as published in 40 Code of Federal Regulations (CFR) 50.13, has been attained. Fine particle

monitoring data must show that violations of the annual ambient standard are no longer occurring.

- 2) Ambient monitoring data quality assured in accordance with 40 CFR 58.15, recorded in the U.S. EPA Air Quality System (AQS) database, and available for public view.
- 3) A showing that the three-year average of annual values, based on data from all monitoring sites in the area or its affected downwind environs, do not exceed 15.0 micrograms per cubic meter ( $\mu\text{g}/\text{m}^3$ ). This showing must rely on three complete, consecutive calendar years of quality assured data.
- 4) A commitment that, once redesignated, the state will continue to operate an appropriate monitoring network to verify the area is in compliance (maintenance) with the standard.

### 2.3 Emission Inventory

- 1) A comprehensive emission inventory of the precursors of fine particles (direct  $\text{PM}_{2.5}$ ,  $\text{NO}_x$ , and  $\text{SO}_2$  completed for the base year (2008 in this case).
- 2) A projection of the emission inventory to a year at least ten years following redesignation.
- 3) A demonstration that the projected level of emissions is sufficient to maintain the annual standard for fine particles.
- 4) A demonstration that improvement in air quality between the year violations occurred and the year attainment was achieved is based on permanent and enforceable emission reductions and not on temporary adverse economic conditions or unusually favorable meteorology.
- 5) Provisions for future updates of the inventory to enable tracking of the emission levels, including emission inventory statements from emission sources.

### 2.4 Modeling Demonstration

While no modeling is required for redesignating nonattainment areas, IDEM has evaluated the results of federal control-case modeling to demonstrate that compliance with the standard will be maintained.

### 2.5 Controls and Regulations

- 1) A U.S. EPA approved SIP control strategy that includes Reasonably Available Control Technology (RACT) requirements for existing stationary sources covered

by Control Technology Guidelines (CTG) and non-CTG RACT for all major sources.

- 2) Evidence that control measures required in past SIP revisions have been fully implemented.
- 3) Acceptable provisions to provide for new source review.
- 4) Assurances that existing controls will remain in effect after redesignation, unless the state demonstrates through photochemical modeling that the standard can be maintained without one or more controls.
- 5) If appropriate, a commitment to adopt a requirement that all transportation plans conform with, and are consistent with, the SIP.

## 2.6 Corrective Actions for Potential Future Violations of the Fine Particle Standard

- 1) A commitment to submit a revised plan eight years after redesignation.
- 2) A commitment to expeditiously enact and implement additional contingency control measures in response to exceeding specified predetermined levels (triggers) or in the event that future violations of the ambient standard occur.
- 3) A list of potential contingency measures that would be implemented in such an event.
- 4) A list of NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub> sources potentially subject to future controls.

## 3.0 FINE PARTICLES MONITORING

### 3.1 Fine Particle Monitoring Network

There are currently twenty-three Federal Reference Method monitors measuring fine particle concentrations in this nonattainment area. Six monitors are located in Indiana's portion of the nonattainment area and are operated by IDEM's Office of Air Quality (OAQ). Seventeen monitors are located in Illinois's portion of the nonattainment and are operated by IEPA's Bureau of Air; Air Monitoring Section (AMS). A listing of the monitor readings from 2008 through 2010 are shown in Tables 3.1, 3.2, and Appendix A and was retrieved from U.S. EPA's AQS database. The locations of the monitoring sites for this nonattainment area are shown in Figure 3.1.


There are also four source oriented monitors located in the nonattainment area that collect background fine particle concentrations. Two monitors are located in Lake County, Indiana and are operated by IDEM's OAQ (Gary – IITRI (18-089-0022) and Gary – Burr Street (18-089-0026)). Two monitors are located in Cook County, Illinois and are operated by IEPA's Bureau



of Air AMS (McCook (17-031-1016) and Schiller Park (17-031-3103)). While the source oriented monitors are not used to determine attainment with the annual standard for fine particles, the monitoring locations and data are included as supporting material.

Three monitors in Indiana's portion of the nonattainment area have been discontinued. The Gary - Federal Building monitor (18-089-1016) in Lake, County, Indiana was discontinued in September 2002. The Dunes National Lakeshore monitor (18-127-0020) in Porter County, Indiana, and the Gary - Ivanhoe School monitor (18-089-1003) in Lake County, Indiana were discontinued on December 31, 2007.

**Figure 3.1**  
**Chicago Nonattainment Area**



### 3.2 Ambient Fine Particles Monitoring Data

The following information summarizes U.S. EPA's "Guideline on Data Handling Conventions for the PM NAAQS," U.S. EPA-454/R-99-008, April 1999. Three complete years of fine particle monitoring data are required to demonstrate attainment at a monitoring site. The annual ambient air quality standard for fine particles is met at an ambient air quality monitoring site

when the three-year average of the annual average of fine particle concentrations is less than or equal to  $15.0 \mu\text{g}/\text{m}^3$ . When this occurs, the site is said to be in attainment. While calculating design values, three significant digits must be carried in the computations, with final values rounded to the nearest  $0.1 \mu\text{g}/\text{m}^3$ . Decimals 0.05 or greater are rounded up, and those less than 0.05 are rounded down, so that  $15.049 \mu\text{g}/\text{m}^3$  is the largest concentration that is less than or equal to  $15.0 \mu\text{g}/\text{m}^3$ . Values at or below  $15.0 \mu\text{g}/\text{m}^3$  meet the standard. Values equal to or greater than  $15.1 \mu\text{g}/\text{m}^3$  exceed the standard.

Data handling procedures are applied on an individual basis at each monitor in the area. An individual site's three-year average of the annual average fine particles concentration is also called the site's *design value*. An area is in compliance with the annual NAAQS for fine particles only if all monitoring sites meet the NAAQS. The air quality design value for the area is the highest design value among all sites in the area. Table 3.1 outlines the annual fine particle values by site and the 2008 through 2010 design values for the six active fine particle monitoring sites in Indiana's portion of the Chicago Area. Table 3.2 outlines the annual fine particle values by site and the 2008 through 2010 design values for the seventeen fine particle monitoring sites in Illinois' portion of the Chicago Area. Appendix A contains the complete monitoring data monitoring summary from 2000 to 2010 for all of the Chicago Area monitors which includes the twenty-three active, the four source oriented, and the three discontinued fine particles monitoring sites.

**Table 3.1**  
**Monitoring Data for Indiana's Portion of the Chicago Area**  
**(Annual Average and 2008 - 2010 Design Values)**

<b>SITE ID</b>	<b>COUNTY</b>	<b>SITE NAME</b>	<b>YEAR</b>	<b>Annual Average (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>2008-2010 Design Value (<math>\mu\text{g}/\text{m}^3</math>)</b>
18-089-0006	Lake	East Chicago – Franklin School	2008	11.95	<b>11.9</b>
			2009	11.34	
			2010	12.48	
18-089-0027	Lake	Griffith	2008	11.69	<b>11.7</b>
			2009	11.00	
			2010	12.39	
18-089-2004	Lake	Hammond – Purdue	2008	11.66	<b>N/A*</b>
			2009	N/A*	
			2010	12.30	
18-089-2010	Lake	Hammond – Clark High School	2008	12.42	<b>11.7</b>
			2009	10.80	
			2010	11.90	
18-089-0031	Lake	Gary – Madison Street	2008	12.27	<b>12.4</b>
			2009	12.12	
			2010	12.90	

18-127-0024	Porter	Ogden Dunes	2008	10.89	<b>11.2</b>
			2009	11.29	
			2008	11.56	

\*The Hammond – Purdue monitoring site was physically removed to accommodate heating and cooling building repairs on the roof which were conducted from February 12, 2009 through November 18, 2009. As a result, the 2009 monitoring data and 2008 through 2010 three year design value are considered incomplete.

**Table 3.2**  
**Monitoring Data for Illinois' Portion of the Chicago Area**  
**(Annual Average and 2008 through 2010 Design Values)**

<b>SITE ID</b>	<b>COUNTY</b>	<b>SITE NAME</b>	<b>YEAR</b>	<b>Annual Average (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>2008-2010 Design Value (<math>\mu\text{g}/\text{m}^3</math>)</b>
17-031-0022	Cook	Chicago – Washington High School	2008	12.54	<b>12.7</b>
			2009	11.62	
			2010	14.04	
17-031-0050	Cook	Chicago – Southeast Police Station	2008	11.80	<b>11.8</b>
			2009	10.99	
			2010	12.47	
17-031-0052	Cook	Chicago – Mayfair Pump Station	2008	12.18	<b>12.5</b>
			2009	12.69	
			2010	12.57	
17-031-0057	Cook	Chicago – Springfield Pump Station	2008	12.03	<b>11.8</b>
			2009	11.33	
			2010	12.03*	
17-031-0076	Cook	Chicago – Com Ed Maintenance Building	2008	11.89	<b>11.8</b>
			2009	11.12	
			2010	12.25	
17-031-2001	Cook	Blue Island	2008	12.50	<b>11.9</b>
			2009	11.68	
			2010	11.59	
17-031-3301	Cook	Summit	2008	12.03	<b>12.0</b>
			2009	11.62	
			2010	12.23	
17-031-4007	Cook	Des Plaines	2008	11.35	<b>11.0</b>
			2009	11.02	
			2010	10.60	
17-031-4201	Cook	Northbrook	2008	10.09	<b>9.6</b>
			2009	9.33	
			2010	9.34	
17-031-6005	Cook	Cicero	2008	13.25*	<b>12.4</b>
			2009	11.98*	
			2010	11.90	


17-043-4002	DuPage	Naperville	2008	11.28	<b>10.9</b>
			2009	9.83	
			2010	11.67	
17-089-0003	Kane	Elgin	2008	10.79	<b>10.6</b>
			2009	9.61	
			2010	11.29	
17-089-0007	Kane	Aurora	2008	10.34	<b>10.6</b>
			2009	10.01	
			2010	11.44	
17-097-1007	Lake	Zion	2008	9.34	<b>9.3</b>
			2009	8.83	
			2010	9.66	
17-111-0001	McHenry	Cary	2008	10.10	<b>10.0</b>
			2009	9.65	
			2010	10.24	
17-197-1002	Will	Joliet	2008	11.66	<b>11.3</b>
			2009	10.52	
			2010	11.83	
17-197-1011	Will	Braidwood	2008	10.31	<b>9.7</b>
			2009	8.73	
			2010	10.02	

\*Indicates the mean does not satisfy summary criteria.

Red Text Indicates Incomplete Data

Graph 3.1 visually demonstrates the 2006 through 2010 design values for Indiana's portion of the Chicago Area.

**Graph 3.1**  
**Fine Particle Design Values for Indiana's Portion of the Chicago Area,**  
**2006 through 2010**




Red line represents the annual standard for fine particles of 15  $\mu\text{g}/\text{m}^3$

The Hammond - Purdue monitoring site was physically removed to accommodate heating and cooling building repairs on the roof which were conducted from February 12, 2009 through November 18, 2009. As a result, the 2007 through 2009 and 2008 through 2010 three year design values are considered incomplete and are not included in Graph 3.1.


Graph 3.2 visually demonstrates the 2006 through 2010 design values for Illinois' portion of the Chicago Area.

**Graph 3.2**  
**Fine Particle Design Values for Illinois' Portion of the Chicago Area,**  
**2006 through 2010**



Red line represents the annual standard for fine particles of  $15 \mu\text{g}/\text{m}^3$ .

**Graph 3.3**  
**Annual Fine Particle Trends for Indiana's Portion of the Chicago Area,**  
**2006 through 2010**




Red line represents the annual standard for fine particles of 15  $\mu\text{g}/\text{m}^3$

The Hammond - Purdue monitoring site was physically removed to accommodate heating and cooling building repairs on the roof which were conducted from February 12, 2009 through November 18, 2009. As a result, the 2009 monitoring data is considered incomplete and has not been included in Graph 3.3.



**Graph 3.4**  
**Annual Fine Particle Trends for Illinois' Portion of the Chicago Area,**  
**2006 through 2010**



Red line represents the annual standard for fine particles of  $15 \mu\text{g}/\text{m}^3$

The design values for Lake and Porter counties, Indiana, along with the nonattainment area in its entirety, demonstrate that the annual NAAQS for fine particles has been attained. Appendix A contains the complete fine particles monitoring data summary for the years 2000 through 2010.

Graphs 3.1 and 3.2 show the trends in design values, while Graphs 3.3 and 3.4 show the trend for annual fine particles. A comprehensive list of the design values for the Chicago Area fine particles monitoring sites over the 2000 through 2010 monitoring period is outlined in Appendix A. The area's design values have recently trended downward, as emissions have declined due to programs such as the Acid Rain program and cleaner automobiles and fuels both regionally and locally. U.S. EPA's rule to control nitrogen oxides from specific source categories (40 CFR Parts 51, 72, 75, and 96, published on October 17, 1998 and referred to as the "NO<sub>x</sub> SIP Call") has significantly reduced emissions from large electric generating units (EGUs), industrial boilers, and cement kilns. Indiana's NO<sub>x</sub> SIP Call Rule was adopted into the Indiana Administrative Code (IAC) on June 6, 2001 at 326 IAC 10-3 and 326 ICA 10-4. The elevated

fine particle values for 2005 are considered an abnormal occurrence. An analysis of meteorological conditions and monitoring values is included in Section 7.0 and supports the conclusion that attainment of the standard as of 2010 is not the result of unusually favorable meteorological conditions. It is expected that this downward trend will persist as the above programs continue and U.S. EPA's proposed Clean Air Transport Rule is implemented.

### 3.3 Quality Assurance

Indiana and Illinois have quality assured all data shown in Appendix A in accordance with 40 CFR 58.10 and recorded the data in the AQS database and, thus, the data is available to the public.

### 3.4 Continued Monitoring

Indiana and Illinois commit to continue monitoring fine particle concentrations at the active sites indicated in Tables 3.1, 3.2, and Appendix A. IDEM will consult with U.S. EPA Region V staff prior to making changes to the existing Indiana monitoring network through the annual network review should changes become necessary in the future. IDEM will continue to quality assure the Indiana monitoring data to meet the requirements of 40 CFR 58. IDEM will enter all data into AQS on a timely basis in accordance with federal guidelines.

## **4.0 EMISSION INVENTORY**

U.S. EPA's Redesignation Guidance and Implementation Rules require the submittal of a comprehensive inventory of precursor emissions for fine particles ( $\text{NO}_x$ ,  $\text{SO}_2$ , and direct  $\text{PM}_{2.5}$ ) representative of the year when the area achieves attainment of the annual NAAQS for fine particles (base year). IDEM is using 2008 as the base year. IDEM must also demonstrate that the improvement in air quality between the year that violations occurred and the year that attainment was achieved is based on permanent and enforceable emission reductions. Other requirements related to the emission inventory include: a projection of the emission inventory to a year at least ten years following redesignation; a demonstration that the projected level of emissions is sufficient to maintain the annual standard for fine particles; and, a commitment to provide future updates of the inventory to enable tracking of emission levels during the ten year maintenance period. Consistent with the federal implementation rule for fine particles, IDEM and U.S. EPA do not consider volatile organic compounds (VOCs) or ammonia ( $\text{NH}_3$ ) to be significant contributors to fine particles. The following subsections address each of these requirements.

### 4.1 Emission Trends


#### Point Sources

The point source data are obtained from Indiana's emissions reporting program and are based on county point source totals. Lake and Porter counties, Indiana, had a 34.31% reduction in  $\text{NO}_x$  point source emissions, a 7.34% reduction in  $\text{SO}_2$  point source emissions, and a 8.71% reduction


in direct PM<sub>2.5</sub> point source emissions from 2002 to 2008. Lake and Porter counties, Indiana, had a 4.77% reduction in NO<sub>x</sub> point source emissions, a slight increase (14.08%) in SO<sub>2</sub> point source emissions, and a slight increase (3.49%) in direct PM<sub>2.5</sub> point source emissions from 2005 to 2008. A moderate increase in Lake and Porter counties, Indiana, in direct PM<sub>2.5</sub> point source emissions from 2005 to 2008 is noted, however, this increase in direct PM<sub>2.5</sub> emissions is due to the fact that most companies did not submit their direct PM<sub>2.5</sub> emissions data in 2005 but did submit direct PM<sub>2.5</sub> data in the 2009 emission inventory, from which the 2008 inventory is extrapolated. Graphs 4.1, 4.2, and 4.3 demonstrate the trend in point source emissions of NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub> for the Lake and Porter counties, Indiana that generally correspond to the years of monitored values used in this redesignation petition.

Point source data for Illinois' portion of the Chicago Area is based on county point source. Point source data for the years 2002 and 2008 were taken directly from Illinois EPA's document entitled *Maintenance Plan for the Chicago Nonattainment for the 1997 PM<sub>2.5</sub> National Ambient Air Quality Standard*, prepared September 7, 2010. This document is hereafter referred to as the "Chicago Maintenance Plan". The 2005 point source data for Illinois' portion of the Chicago nonattainment area were interpolated from the 2002 and 2008 emissions data. The Chicago Area had a 33.93% reduction in NO<sub>x</sub> point source emissions, a 19.48% reduction in SO<sub>2</sub> point source emissions, and a slight increase (4.61%) in direct PM<sub>2.5</sub> point source emissions from 2002 to 2008. The Chicago Area had a 12.72% reduction in NO<sub>x</sub> point source emissions, a 5.62% reduction in SO<sub>2</sub> point source emissions, and a slight increase (7.95%) in direct PM<sub>2.5</sub> point source emissions from 2005 to 2008. The moderate increase in direct PM<sub>2.5</sub> emissions is far outweighed by the significant reductions in NO<sub>x</sub> and SO<sub>2</sub> that have occurred and will continue to occur in the future. Graphs 4.4, 4.5, and 4.6 demonstrate the trend in point source emissions of NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub> emissions for the entire Chicago Area. Graphs and data tables of emissions for the point source category can be found in Appendix B.


**Graph 4.1**  
**NO<sub>x</sub> Point Source Emission Trends, Lake and Porter Counties,**  
**Indiana, 2002, 2005, and 2008**




**Graph 4.2**  
**SO<sub>2</sub> Point Source Emission Trends, Lake and Porter Counties,**  
**Indiana, 2002, 2005, and 2008**




**Graph 4.3**  
**Direct PM<sub>2.5</sub> Point Source Emission Trends, Lake and Porter Counties,**  
**Indiana, 2002, 2005, and 2008**




**Graph 4.4**  
**NO<sub>x</sub> Point Source Emission Trends, Entire Chicago Area, 2002, 2005, and 2008**



**Graph 4.5**  
**SO<sub>2</sub> Point Source Emission Trends Entire Chicago Area, 2002, 2005, and 2008**



**Graph 4.6**  
**Direct PM<sub>2.5</sub> Point Source Emission Trends, Entire Chicago Area, 2002, 2005, and 2008**



#### All Anthropogenic Sources


Periodic inventories, which include emissions from all sectors (mobile, area, nonroad, and point source), were prepared for 2002, 2005, and 2008. The 2008 data for Lake and Porter counties, Indiana were extrapolated from the 2005 emission inventory. The 2002 and 2008 data for Illinois' portion of the Chicago Area were taken directly from the Chicago Maintenance Plan. The 2005 data were interpolated from the 2002 and 2008 emissions data. Regional NO<sub>x</sub> emission reductions affect fine particle levels in the Chicago Area far more so than NO<sub>x</sub> emission reductions within the nonattainment area itself. These emission trends roughly follow the years of monitored trends discussed in Section 3.0. There is a downward trend in NO<sub>x</sub> emissions from 2002 to 2005 and a further decrease through 2008. The decrease in NO<sub>x</sub> can be largely attributed to the impact of the NO<sub>x</sub> SIP Call. There is a general downward trend in direct PM<sub>2.5</sub> and SO<sub>2</sub> emissions from 2002 to 2008, as well. As noted previously, there is an overall increase in direct PM<sub>2.5</sub> anthropogenic emissions in Lake and Porter counties, Indiana, but an overall decrease in the entire nonattainment area. Graphs and data tables of emissions from each source category are available in Appendix C.

Mobile emission inventories and projections for Lake and Porter counties, Indiana were prepared by the Northwestern Indiana Regional Planning Commission (NIRPC), the Indiana Department of Transportation, and IDEM and are explained in further detail in Section 5.0. All 2005 data for Lake and Porter counties, Indiana are from the 2005 periodic inventory which has been identified as one of the preferred databases for SIP development. All 2008 emissions were extrapolated

from the 2005 Lake Michigan Air Directors Consortium's (LADCO's) modeling inventory, using LADCO's growth factors, for all sectors except point sources (electrical generating units and non-electrical generating units). Point source emissions for 2008 were compiled from Indiana's annual emissions inventory database.


Graphs 4.7, 4.8, and 4.9 show the trend in anthropogenic emissions for Lake and Porter counties, Indiana. Graphs 4.10, 4.11, and 4.12 show the trend in anthropogenic emissions for the entire Chicago Area. The emission inventory development and emissions projection discussion below, with the exception of the mobile emission inventory and projections, identify procedures used by IDEM and LADCO regarding emissions for the Chicago Area.

**Graph 4.7**  
**NO<sub>x</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008**







**Graph 4.8**  
**SO<sub>2</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008**




**Graph 4.9**  
**Direct PM<sub>2.5</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008**




**Graph 4.10**  
**NO<sub>x</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008**



**Graph 4.11**  
**SO<sub>2</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008**



**Graph 4.12**  
**Direct PM<sub>2.5</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008**



#### EGU Sources

Both NO<sub>x</sub> and SO<sub>2</sub> emissions are decreasing substantially in Lake and Porter counties, Indiana, as well as statewide in response to national programs affecting all EGUs, such as the Acid Rain program and the NO<sub>x</sub> SIP Call. Other sectors of the inventory also impact the formation of fine particles, but large regional sources, such as EGUs, have a substantial impact on the formation of fine particles. The data was taken from U.S. EPA's Clean Air Markets database.<sup>1</sup>

As part of the NO<sub>x</sub> SIP Call, the states were required to adopt into their rules a budget for all large EGUs. Indiana's budget is referenced in 326 IAC 10-4. The budget represents a statewide cap on NO<sub>x</sub> emissions. Although each unit is allocated emissions based upon historic heat input, utilities can meet this budget by over-controlling certain units or purchasing credits from the market to account for overages at other units. To summarize, NO<sub>x</sub> emissions have dramatically decreased over the years represented on these graphs.


These emissions, capped by the state rule, are expected to remain near these levels throughout the maintenance period covered by this request. The state cap for the NO<sub>x</sub> SIP Call remained in place through 2008, at which time the Clean Air Interstate Rule (CAIR) program superseded it. CAIR, issued in March 2005, adopted by the Indiana Air Pollution Control Board on November 1, 2006, and implemented beginning in 2010, will continue to reduce regional EGU NO<sub>x</sub>

<sup>1</sup> <http://www.epa.gov/airmarkets>


emissions statewide by approximately another 17% by 2015 and 57% for EGU SO<sub>2</sub> emissions by 2015. The D.C. Circuit court's vacatur of CAIR in July of 2008 and subsequent remand without vacatur of CAIR in December of 2008 directs U.S. EPA to revise the CAIR rule in the future. The proposed Clean Air Transport Rule (CAIR's replacement rule) will result in similar or greater emission reductions than assumed within the current emission inventories once it is implemented.

There are three EGUs located in Indiana's portion of the nonattainment area: The Northern Indiana Public Service Company (NIPSCO) – Bailly Generating Station, Dominion Resources – State Line Generating Plant, and British Petroleum – Whiting Clean Energy Power Plant. Graphs 4.13 and 4.15 depict the trends in NO<sub>x</sub> and SO<sub>2</sub> emissions from EGUs in Lake and Porter counties, Indiana for the years 1999 through 2009. As the result of a recent settlement with U.S. EPA to resolve violations of the CAA's new source review requirements, NIPSCO has agreed to upgrade existing pollution control equipment, as well as install additional pollution control equipment at three of its four coal-fired power plants located in Northwest Indiana (i.e. Bailly, Michigan City, and R.H. Schafer Generating Stations) to comply with stringent NO<sub>x</sub> and SO<sub>2</sub> emission rates and annual tonnage limitations. The settlement also requires NIPSCO to optimize existing PM pollution control equipment to meet unit-specific emissions and permanently retire its fourth facility, the Dean H. Mitchell Generating Station. The Dean H. Mitchell Generating Station has been out of operation since 2002 and its permanent retirement will ensure that the facility does not restart without proper permitting under the CAA. Graphs 4.13 and 4.15 depict the trends in NO<sub>x</sub> and SO<sub>2</sub> EGU emissions in Lake and Porter counties, Indiana for the years 1999 through 2009. Graphs 4.14 and 4.16 depict the trends in statewide NO<sub>x</sub> and SO<sub>2</sub> EGU emissions for the years 1999 through 2009. Graphs and data tables of emissions from the EGU source category can be found in Appendix D.


**Graph 4.13**  
**NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana, 1999 to 2009**




**Graph 4.14**  
**Indiana Statewide NO<sub>x</sub> Emissions from EGUs, 1999 to 2009**



**Graph 4.15**  
**SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana, 1999 to 2009**



**Graph 4.16**  
**Indiana Statewide SO<sub>2</sub> Emissions from EGUs, 1999 to 2009**



#### 4.2 Base Year Inventory

IDEM prepared a comprehensive inventory for the Chicago Area, including area, mobile, nonroad, and point sources for direct PM<sub>2.5</sub>, NO<sub>x</sub>, and SO<sub>2</sub> for 2005 and 2008 (the years with the most complete emissions inventory available at this time). The 2008 emission inventory represents a base year for maintenance purposes. The 2007 implementation rule for the annual standard for fine particles states that NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub> are the regulated precursors of fine particles. Ammonia and VOCs are not required to be addressed unless the state or U.S. EPA make a technical demonstration that emissions of these pollutants from sources in the state significantly contribute to PM<sub>2.5</sub> concentrations in a given nonattainment area. U.S. EPA and IDEM have not determined ammonia or VOCs are significant contributors to fine particles formation in Indiana. IDEM's 2008 base year inventory was developed as follows:

- Area sources were extrapolated from the Indiana 2005 periodic inventory submitted to U.S. EPA.
- Mobile source emissions were calculated from MOBILE6.2 model-produced emission factors and data extracted from the region's travel-demand model. These emissions were then interpolated as needed to determine 2008 base year values.
- Point source information was compiled from IDEM's emissions inventory database and U.S. EPA's Clean Air Markets acid rain database.
- Biogenic emissions are not specifically included in these summaries, but are included in the photochemical modeling results presented in Section 7.0.

- Nonroad emissions were extrapolated from the 2005 National Emissions Inventory (NEI). To address concerns about the accuracy of some of the categories in U.S. EPA's nonroad emissions model, Lake Michigan Air Directors Consortium (LADCO) contracted with two companies to review the base data and make recommendations. One of the contractors also estimated emissions for two nonroad categories not included in U.S. EPA's nonroad model. Emissions were estimated for commercial marine vessels and railroads. The recreational motorboat population and spatial surrogates (used to assign emissions to each) were also significantly updated. The populations for the construction equipment category were also reviewed and updated based upon surveys completed in the Midwest and the temporal allocation for agricultural sources was also updated. A new nonroad estimation model was provided by U.S. EPA for the 2005 analysis.
- Area, nonroad, mobile, and point source emissions data referenced for Illinois' portion of the Chicago Area were taken directly from the Chicago Maintenance Plan. This inventory was prepared using similar methodologies. Emissions for the year 2005 were interpolated from the 2002 and 2008 emissions data.

Appendix C contains data tables and graphs of these emissions.

#### 4.3 Emission Projections

In consultation with U.S. EPA and other stakeholders, IDEM selected the year 2025 as the maintenance year for this redesignation request. This document contains projected emission inventories for 2015, 2020, and 2025. These emission projections were prepared by IDEM, with assistance from LADCO, Illinois EPA, and the Northwest Indiana Regional Planning Commission.

The detailed 2015, 2020, and 2025 emission inventory for the Chicago Area can be found in Appendix E. Emission trends are an important gauge for continued compliance with the annual standard for fine particles. Therefore, IDEM performed an initial comparison of the inventories for the base year of 2008, interim years of 2015 and 2020, and maintenance year of 2025 for Lake and Porter counties, Indiana, as well as the entire Chicago Area.


The 2005 LADCO modeling inventory was used as the basis for estimated emissions for Lake and Porter counties, Indiana for the years 2008, 2015, 2020, and 2025, using LADCO's growth factors, for all sectors except point sources (electric generating units and non-electric generating units). Point source emissions for 2005 and 2008 were compiled from Indiana's annual emission inventory database. The 2015 interim year emissions were interpolated based on the 2009 and 2018 LADCO modeling inventory, using LADCO's growth factors, for all sectors. The 2020 interim year emissions and the 2025 maintenance year emissions were extrapolated from the 2009 and 2018 LADCO modeling inventory.

The projected emission inventories for 2015, 2020, and 2025 for Illinois' portion of the Chicago Area were taken directly from the Chicago Maintenance Plan.

Graphs 4.17, 4.18, 4.19, and 4.20 visually compare 2005 and 2008 (base year) NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub> county total estimated emissions with the 2015, 2020, and 2025 projected emissions for Lake and Porter counties, Indiana. Graphs 4.21, 4.22, 4.23, and 4.24 visually compare the 2005 and 2008 (base year) NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub> county total estimated emissions with the 2015, 2020, and 2025 projected emissions for the entire Chicago Area. Mobile source emission inventories are further described in Section 5.0. In addition to LADCO's estimates, point source emissions were projected based upon the statewide EGU NO<sub>x</sub> budgets from Indiana's NO<sub>x</sub> rule.


EGU emission estimates for 2015, 2020, and 2025 were projected using the budgets from the proposed Transport Rule and Options 1 and 2 from the recent Notice of Data Availability. Since it is likely that the finalized rule will include a budget approximating one of these, IDEM used the option with the highest allotment each for the NIPSCO – Bailly Generating Station, Dominion Resources – State Line Generating Plant, and British Petroleum – Whiting Clean Energy Power Plant to develop conservative projections. These allowances cover 2014 forward, so the same projections were used for all three years. Further, the consent decree provides overall caps for all NIPSCO facilities in Northwest Indiana. The emissions allowed under these caps are similar to those in the Transport Rule. Therefore, IDEM is confident that emissions in these three years will be close to or below the projections. Graphs and data tables of emissions from the EGU source category can be found in Appendix D.

**Graph 4.17**  
**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO<sub>x</sub>**  
**Emissions, Lake and Porter Counties, Indiana**







**Graph 4.18**  
**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected SO<sub>2</sub>**  
**Emissions, Lake and Porter Counties, Indiana**




**Graph 4.19**  
**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected Direct PM<sub>2.5</sub>**  
**Emissions, Lake and Porter Counties, Indiana**




**Graph 4.20**  
**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO<sub>x</sub>, SO<sub>2</sub>, and Direct PM<sub>2.5</sub> Emissions, Lake and Porter Counties, Indiana**




**Graph 4.21**  
**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO<sub>x</sub> Emissions, Entire Chicago Area**




**Graph 4.22**  
**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected SO<sub>2</sub> Emissions, Entire Chicago Area**



**Graph 4.23**  
**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected Direct PM<sub>2.5</sub> Emissions, Entire Chicago Area**



**Graph 4.24**  
**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO<sub>x</sub>, SO<sub>2</sub>,  
and Direct PM<sub>2.5</sub> Emissions, Entire Chicago Area**



NO<sub>x</sub> emissions within Lake and Porter counties, Indiana are projected to decline by 41.23% between 2008 and 2025. NO<sub>x</sub> emissions within the Chicago Area are projected to decline by 46.46% between 2008 and 2025. Emission reduction benefits from U.S. EPA rules covering the NO<sub>x</sub> SIP Call, Tier 2 Motor Vehicle Emission Standards and Gasoline Sulfur Control Requirements, Heavy-Duty Highway Engine Rule, and the Nonroad Diesel Engine Rule are factored into the changes. Additionally, due to implementation of the NO<sub>x</sub> SIP Call across the eastern United States, NO<sub>x</sub> and fine particle levels entering the Chicago Area will also be decreased. SO<sub>2</sub> emissions within Lake and Porter counties, Indiana are projected to decrease by 35.54% between 2008 and 2025. SO<sub>2</sub> emissions within the Chicago Area are projected to decline by 34.73% between 2008 and 2025. Direct PM<sub>2.5</sub> emissions within Lake and Porter counties, Indiana are projected to decrease by 6.74% between 2008 and 2025. Direct PM<sub>2.5</sub> emissions within the Chicago Area are projected to decline by 2.69% between 2008 and 2025.

**Table 4.1**  
**Comparison of 2008 Estimated and 2025 Projected Emission Estimates,**  
**Lake and Porter Counties, Indiana (Tons per Year)**

	<b>2008</b>	<b>2025</b>	<b>Change</b>	<b>% Change</b>
NO <sub>x</sub>	59,581.74	35,013.79	-24,567.95	41.23% decrease
SO <sub>2</sub>	56,198.68	36,787.24	-19,411.44	35.54% decrease
Direct PM <sub>2.5</sub>	7,250.93	6,762.10	-488.83	6.74% decrease

**Table 4.2**  
**Comparison of 2008 Estimated and 2025 Projected Emission Estimates,**  
**Entire Chicago Area (Tons per Year)**

	<b>2008</b>	<b>2025</b>	<b>Change</b>	<b>% Change</b>
NO <sub>x</sub>	278,649.74	149,198.79	-129,450.95	46.46% decrease
SO <sub>2</sub>	152,367.68	99,453.24	-52,914.44	34.73% decrease
Direct PM <sub>2.5</sub>	25,767.93	25,074.10	-693.83	2.69% decrease

#### 4.4 Demonstration of Maintenance

Quality assured ambient air quality data from all the monitoring sites indicate that air quality in Indiana's portion of the nonattainment area met the annual standard for fine particles for the three-year period ending in 2010. U.S. EPA's Redesignation Guidance states, "A state may generally demonstrate maintenance of the NAAQS by either showing that future emissions of a pollutant or its precursors will not exceed the level of the attainment inventory, or by modeling to show that the future mix of sources and emission rates will not cause a violation of the NAAQS." Section 3.0 of this document shows that the Chicago Area has in fact measured attainment for fine particles for the three consecutive periods ending in 2008, 2009, and 2010. Additionally, emission projections outlined in Section 4.0 of this document clearly illustrate that regional NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub> emissions in the area will continue to decline leading to local reductions between 2008 (base year) and 2025 (maintenance plan horizon). Section 7.0 further discusses the implications of these emission trends and provides an analysis to support these conclusions.

In Indiana, major point sources in all counties are required to submit air emissions information once every three years, or annually, if the NO<sub>x</sub> or SO<sub>2</sub> potential to emit is greater than 2,500 tons per year in accordance with the Emission Reporting Rule, 326 IAC 2-6. IDEM prepares a new periodic inventory for all precursor emission sectors every three years. These precursor emission inventories will be prepared for 2011, 2014, and 2017, as necessary, to comply with the inventory reporting requirements established in the CAA. Emissions information will be compared to the 2008 base year and the 2025 projected maintenance year inventories to assess emission trends, as necessary, to assure continued compliance with the annual standard for fine particles.

#### 4.5 Permanent and Enforceable Emission Reductions

Permanent and enforceable reductions of NO<sub>x</sub> and SO<sub>2</sub> have contributed to the attainment of the annual standard for fine particles. Some of these reductions were due to the implementation of the NO<sub>x</sub> SIP Call rule and some were due to the application of tighter federal standards on motor vehicles and fuels.

Section 6.0 identifies the emission control measures specific to Lake and Porter counties, Indiana, as well as the implementation status of each measure.

#### 4.6 Provisions for Future Updates

As required by Section 175A(b) of the CAA, Indiana commits to submit to the Administrator, eight years after redesignation, an additional revision of this SIP. The revision will contain Indiana's plan for maintaining the national primary fine particles air quality standard for ten years beyond the first ten-year period after redesignation.

### **5.0 TRANSPORTATION CONFORMITY BUDGETS**

U.S. EPA requirements outlined in 40 CFR 93.118(e)(4) stipulate that motor vehicle emission budgets (MVEBs) for direct PM<sub>2.5</sub> and NO<sub>x</sub> be established as part of a SIP. The MVEBs are necessary to demonstrate conformance of transportation plans and improvement programs with the SIP.

#### 5.1 On-Road Emission Estimates

The Northwest Indiana Regional Planning Commission (NIRPC) is the Metropolitan Planning Organization (MPO) for the area that includes Lake, Porter, and LaPorte counties. This organization maintains a travel demand forecast model that is used to simulate the traffic in the area and are used to predict what the traffic would be like in future years given growth expectations. They are also used mostly to identify where travel capacity will be needed and to determine the infrastructure requirements necessary to meet that need. It is also used to support the calculation of mobile source emissions. The travel demand forecast model is used to predict the total daily vehicle miles traveled (VMT) and the U.S. EPA software program, MOBILE6.2, is used to calculate per mile emissions. The product of these two outputs, once combined, is the estimated total amount of pollution emitted by the on-road vehicles for the area analyzed.

#### 5.2 Overview

Broadly described, MOBILE6.2 is used to determine “emission factors”, which are the average emissions per mile (grams/mile) for fine particles and fine particle precursors (NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub>). There are numerous variables that can affect the emission factors. The vehicle-fleet (vehicles on the road) age and the vehicle types have a major effect on the emission factors. The facility-type on which vehicles are traveling (MOBILE6.2 facility types are Freeway, Arterial, Local, and Ramp) and the vehicle speeds also affect the emission factor values.

Meteorological factors such as air temperature and humidity affect emission factors, as does fuel type, such as low Reid Vapor Pressure gasoline. These data are estimated using the *best available data* to create emission factors for PM<sub>2.5</sub> precursors (NO<sub>x</sub>, SO<sub>2</sub>, and direct PM<sub>2.5</sub>). After emission factors are determined, they must be multiplied by the VMT to determine the quantity of vehicle-related emissions. This information is derived from the travel demand model.

It should be noted that each year analyzed will have different emission factors, volumes, speeds, and likely results in additional modeling. MOBILE6.2 input and output files are found in Appendix F.

### 5.3 Analysis Years

The travel demand model contains road networks that are time specific. The NIRPC has modeled the years 2005, 2010, 2020, and 2030. Information, including emissions, has also been interpolated from 2005 and 2010 for the year 2008, from 2010 and 2020 for the years 2015 and 2016, and from 2020 and 2030 for the year 2025. This Redesignation Petition provides emission inventory estimates for 2002, 2005, 2008, 2015, 2020, and 2025 to meet the requirements specified by the CAA and U.S. EPA. The emission estimates outlined in Section 4.0 of this document reference the mobile source emissions data contained in Table 5.1.

### 5.4 Emission Estimates

Table 5.1 outlines on-road emission estimates for the entire nonattainment area for the years 2005, 2008 (base/attainment year), 2015, 2016, 2020 (interim years), and 2025 (horizon year).

**Table 5.1**  
**Emission Estimates for On-Road Mobile Sources**

<b>Chicago Area</b>	<b>2005</b>	<b>2008</b>	<b>2015</b>	<b>2016</b>	<b>2020</b>	<b>2025</b>
Direct PM <sub>2.5</sub> (tons/year)	2,443.50	1,816.00	1,231.00	1,137.00	1,043.00	1,064.00
NO <sub>x</sub> (tons/year)	133,623.50	99,627.00	49,746.00	39,424.00	29,102.00	23,687.00
<b>Lake and Porter Counties, IN subtotal</b>						
Direct PM <sub>2.5</sub> (tons/year)	229.39	187.09	136.61	132.15	114.32	115.39
NO <sub>x</sub> (tons/year)	14,095.55	10,703.81	5,723.67	5,179.87	3,004.68	2,534.95
<b>Lake and Porter Counties, IN subtotal%</b>						
Direct PM <sub>2.5</sub> (tons/year)	9.39%	10.30%	11.10%	11.62%	10.96%	10.84%
NO <sub>x</sub> (tons/year)	10.55%	10.74%	11.51%	13.14%	10.32%	10.70%

## 5.5 Motor Vehicle Emission Budget

Table 5.2 contains the motor vehicle emission budget for the Indiana portion of the nonattainment area for the years 2016 and 2025. These MVEBs are intended to be direct replacements of those provided in Indiana's previous submittal.

**Table 5.2**  
**Motor Vehicle Emission Budgets for Lake and Porter Counties**

	2016	2025
<b>PM<sub>2.5</sub> (tons/year)</b>	145.37	132.70
<b>NO<sub>x</sub> (tons/year)</b>	5,697.86	2,915.19

Consistent with the federal implementation rule for fine particles, Indiana does not consider mobile source SO<sub>2</sub> to be a significant contributor to fine particles for this nonattainment area, as mobile source SO<sub>2</sub> constitutes less than one percent (<1%) of the area's total anthropogenic emissions.

These budgets include emission estimates calculated for 2016 and 2025. The emission estimates are derived from the MPO's travel demand models and MOBILE6.2, as described above. Through the interagency consultation process, it was determined that an interim budget year of 2016, in addition to the horizon year budget of 2025, would be appropriate. The interagency consultation group approved a ten percent (10%) safety margin for PM<sub>2.5</sub> and NO<sub>x</sub> mobile source emissions estimates for the year 2016 and a fifteen percent (15%) safety margin for PM<sub>2.5</sub> and NO<sub>x</sub> mobile source emissions estimates for the year 2025. Margins of safety are used to accommodate the wide array of assumptions that are factored into the calculation process. Since assumptions change over time, it is necessary to have a margin of safety that will accommodate the impact of refined assumptions in the process.

The rule at 40 CFR 93.101 defines safety margin as the amount by which the total projected emissions from all sources of a given pollutant are less than the total emissions that would satisfy the applicable requirement for reasonable further progress, attainment, or maintenance. When compared to the overall safety margin as defined by 40 CFR 93.101, it is evident this allocation to mobile sources is significantly below the total safety margin for all sources in Lake and Porter counties, Indiana as detailed in Table 4.1.

## 5.6 Commitment to Amend Motor Vehicle Emission Budgets Using Motor Vehicle Emission Simulator (MOVES)

On March 2, 2010, U.S. EPA published a Notice of Availability for the MOVES model. IDEM is committed to submitting a revision to the MOBILE6.2 developed MVEBs using the MOVES model as soon as possible and well in advance of the March 2, 2012, expiration of the transportation conformity "grace period." IDEM recognizes that U.S. EPA will allow the MOBILE6.2 budgets to be replaced through an adequacy notice in place of a full publication to the Federal Register.



All methodologies, latest planning assumptions, margins of safety, and MOVES model commitments were determined appropriate through the interagency consultation process.

## **6.0 CONTROL MEASURES AND REGULATIONS**

This section provides specific information on the control measures that have been or will be implemented in Lake and Porter counties, Indiana, including CAA requirements and additional state or local measures implemented beyond CAA requirements.

### **6.1 Reasonably Available Control Technology (RACT)**

As required by Section 172 of the CAA, in the mid-1990s, Indiana promulgated rules requiring RACT for emissions of VOCs. There were no specific rules required by the CAA, such as RACT for existing sources, beyond statewide rules. Statewide RACT rules have applied to all new sources locating in Indiana since that time. The Indiana rules are found in 326 IAC 8. The following is a listing of applicable rules:

326 IAC 8-1-6	BACT for Non-Specific Sources
326 IAC 8-2	Surface Coating Emission Limitations
326 IAC 8-3	Solvent Degreasing Operations
326 IAC 8-4	Petroleum Sources
326 IAC 8-5	Miscellaneous Operation
326 IAC 8-6	Organic Solvent Emission Limitations

Since Lake and Porter counties, Indiana attained the annual standard for fine particles prior to an Attainment SIP or RACT SIP being due, and since the Implementation Rule for fine particles stipulates that states are only required to draft and implement RACT rules for the precursor emission reductions necessary to attain the standard, no further RACT rules are required for this area. These Indiana rules are CAA requirements already in the SIP and provide secondary benefits for PM<sub>2.5</sub>.

### **6.2 Implementation of Past SIP Revisions**

The Chicago Area was designated nonattainment for the annual standard for fine particles in 2003 and the area has now attained the standard well in advance of its attainment deadline of 2010. As a result, Indiana is no longer required to develop and submit an Attainment SIP or RACT SIP for this area under the annual NAAQS for fine particles.

### **6.3 Nitrogen Oxides Rule**

The U.S. EPA NO<sub>x</sub> SIP Call required twenty-two states to adopt rules that would result in significant emission reductions from large EGUs, industrial boilers, and cement kilns in the eastern United States. The Indiana rule was adopted in 2001. Beginning in 2004, this rule

accounts for a reduction of approximately 31% of all NO<sub>x</sub> emissions statewide compared to previous uncontrolled years.

Twenty-one other states have also adopted these rules. The result is that significant reductions have occurred regionally and within the nonattainment area because of the number of affected units within the region. From Graphs 4.13 and 4.14, it can be seen that emissions covered by this program have been trending downward since 1999. Table 6.1, compiled from data taken from the U.S. EPA Clean Air Markets web site, quantifies the gradual NO<sub>x</sub> reductions that have occurred in Indiana as a result of Title IV (Acid Rain) of the CAA and the NO<sub>x</sub> SIP Call Rule. The NO<sub>x</sub> SIP Call cap stayed in place through 2008, at which time the caps in the CAIR program superseded it. Since CAIR is a regional cap and trade program, it cannot be predicted at this time what affect this will have on EGUs located in the nonattainment area or other upwind counties.

Further, U.S. EPA published Phase II of the NO<sub>x</sub> SIP Call that established a budget for large (emissions of greater than 1 ton per day) stationary internal combustion engines. In Indiana, the rule decreases emissions statewide from natural gas compressor stations by 4,263 tons during the ozone season (April through September). The Indiana Phase II NO<sub>x</sub> SIP Call Rule became effective February 26, 2006 and implementation began in 2007.

**Table 6.1**  
**Trends in EGU NO<sub>x</sub> Emissions Statewide in Indiana**

<b>Year</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
1999	347,216.5
2000	334,522.1
2001	315,419.7
2002	281,146.1
2003	260,980.0
2004	224,311.3
2005	207,981.6
2006	202,728.0
2007	196,553.1
2008	196,134.5
2009	110,968.9
Budget 2009-2014	108,935
Budget 2015 and later	90,779

#### 6.4 Measures Beyond Clean Air Act SIP Requirements

Reductions in fine particle precursor emissions have occurred, or are anticipated to occur, as a result of local and federal programs. These additional control measures include those listed in this section.

### Tier 2 Vehicle Standards<sup>2</sup>

Federal Tier 2 motor vehicle standards require all passenger vehicles in a manufacturer's fleet, including light-duty trucks and sport utility vehicles (SUVs), to meet an average standard of 0.07 grams of NO<sub>x</sub> per mile. Implementation began in 2004 and was completed in 2007. The Tier 2 standards also cover passenger vehicles over 8,500 pounds gross vehicle weight rating (larger pickup trucks and SUVs), which are not covered by the current Tier 1 standards. For these vehicles, the standards were phased in beginning in 2008, with full compliance in 2009. The new standards require vehicles to be 77% to 95% cleaner than those on the road prior to the program. The Tier 2 standards also reduced the sulfur content of gasoline to 30 parts per million (ppm) beginning in January 2006. Most gasoline sold in Indiana prior to January 2006 had a sulfur content of about 500 ppm. Sulfur occurs naturally in gasoline, but interferes with the operation of catalytic converters on vehicles resulting in higher NO<sub>x</sub> emissions. Lower sulfur gasoline is necessary to achieve the Tier 2 vehicle emission standards.

### Heavy-Duty Gasoline and Diesel Highway Vehicle Standards<sup>3</sup>

New U.S. EPA standards designed to reduce NO<sub>x</sub> and VOC emissions from heavy-duty gasoline and diesel highway vehicles took effect in 2004. A second phase of standards and testing procedures that began in 2007, reduced fine particles from heavy-duty highway engines and also reduced highway diesel fuel sulfur content to 15 ppm since the sulfur can damage emission control devices. The total program is expected to achieve a 90% reduction in direct PM<sub>2.5</sub> emissions and a 95% reduction in NO<sub>x</sub> emissions for these new engines using low sulfur diesel, compared to existing engines using higher sulfur content diesel. There will also be SO<sub>2</sub> reductions from these rules. U.S. EPA has not quantified the expected reductions.

### Large Nonroad Diesel Engine Standards<sup>4</sup>

In May 2004, U.S. EPA promulgated new rules for large nonroad diesel engines, such as those used in construction, agricultural, and industrial equipment, to be phased in between 2008 and 2014. The nonroad diesel rules also reduce the allowable sulfur in nonroad diesel fuel by over 99%. Prior to 2006, nonroad diesel fuel averaged approximately 3,400 ppm sulfur. This rule limited nonroad diesel sulfur content to 500 ppm by 2006 with a further reduction to 15 ppm by 2010. The combined engine and fuel rules will reduce NO<sub>x</sub> and PM emissions from large nonroad diesel engines by over 90%, compared to current nonroad engines using higher sulfur content diesel.

### Nonroad Spark-Ignition Engines and Recreational Engine Standards

This new standard, effective in July 2003, regulates NO<sub>x</sub>, VOCs, and carbon monoxide (CO) for groups of previously unregulated nonroad engines. The new standard applies to all new engines sold in the United States and imported after the standards went into effect. The standard applies to large spark-ignition engines (forklifts and airport ground service equipment), recreational

---

<sup>2</sup> <http://www.epa.gov/fedrgstr/EPA-AIR/2000/February/Day-10/a19a.htm>

<sup>3</sup> <http://www.epa.gov/fedrgstr/EPA-AIR/1997/October/Day-21/a27494.htm>

<sup>4</sup> <http://www.epa.gov/fedrgstr/EPA-AIR/1998/October/Day-23/a24836.htm>

vehicles (off-highway motorcycles and all-terrain vehicles), and recreational marine diesel engines. The regulation varies based upon the type of engine and vehicle.

The emissions from large spark-ignition engines contribute to ozone formation and ambient CO and PM<sub>2.5</sub> levels in urban areas. Tier 1 of this standard was implemented in 2004 and Tier 2 started in 2007. Like the large spark-ignition engines, emissions from recreational vehicles contribute to ozone and fine particles formation and ambient CO and PM<sub>2.5</sub> levels. For Model Year 2006 off-highway motorcycles and all-terrain vehicles, at least 50% of a manufacturer's fleet was required to meet the new exhaust emissions standard and 100% of the fleet was required to meet the standards in 2007. Recreational marine diesel engines over 37 kilowatts are used in yachts, cruisers, and other types of pleasure craft. Emissions from recreational marine engines contribute to ozone formation and PM levels, especially surrounding marinas.

When all of the nonroad spark-ignition engines and recreational engine standards are fully implemented, an overall 72% reduction in VOC, 80% reduction in NO<sub>x</sub>, and 56% reduction in CO emissions are expected by 2020.

#### *Reciprocating Internal Combustion Engine Standards*<sup>5</sup>

This new standard, effective in May 2010, regulates emissions of air toxics from existing diesel powered stationary reciprocating internal combustion engines that meet specific site rating, age, and size criteria. These engines are typically used at industrial facilities (e.g. power, chemical, and manufacturing plants) to generate electricity for compressors and pumps and to produce electricity to pump water for flood and fire control during emergencies.

The standard applies to stationary diesel engines: (1) used at area sources of air toxics and constructed or reconstructed before June 12, 2006; (2) used at major sources of air toxics, having a site rating of less than or equal to 500 horsepower, and constructed or reconstructed before June 12, 2006; and, (3) used at major sources of air toxics for non-emergency purposes, having a site rating of greater than 500 horsepower, and constructed or reconstructed before December 19, 2002.

Operators of existing engines will be required to: (1) install emissions control equipment that would limit air toxics up to 70% for stationary non-emergency engines with a site rating greater than 300 horsepower; (2) perform emission tests to demonstrate engine performance and compliance with rule requirements; and, (3) burn ultra-low sulfur fuel in stationary non-emergency engines with a site rating greater than 300 horsepower.

When all of the reciprocating internal combustion engine standards are fully implemented in 2013, U.S. EPA estimates that emissions from these engines will reduce air toxics by approximately 1,000 tons per year (tpy), PM<sub>2.5</sub> by 2,800 tpy, CO by 14,000 tpy, and VOC by 27,000 tpy.

---

<sup>5</sup> <http://www.epa.gov/ttn/atw/rice/fr03mr10.pdf>

### Category 3 Marine Diesel Engine Standards<sup>6</sup>

This new standard, effective in June 2010, promulgates more stringent exhaust emission standards for new large marine diesel engines with per-cylinder displacement at or above 30 liters (commonly referred to as Category 3 compression-ignition marine engines) as part of a coordinated strategy to address emissions from all ships that affect U.S. air quality. These emission standards are equivalent to those adopted in the amendments to Annex VI to the International Convention for the Prevention of Pollution from Ships (MARPOL Annex VI). The emission standards apply in two stages—near-term standards for newly built engines will apply beginning in 2011; long-term standards requiring an 80% reduction in NO<sub>x</sub> emissions will begin in 2016.

U.S. EPA is adopting changes to the diesel fuel program to allow for the production and sale of diesel fuel with up to 1,000 ppm sulfur for use in Category 3 marine vessels. The regulations generally forbid production and sale of fuels with more than 1,000 ppm sulfur for use in most U.S. waters, unless operators achieve equivalent emission reductions in other ways.

U.S. EPA is also adopting provisions to apply some emission and fuel standards to foreign-flagged and in-use vessels that are covered by MARPOL Annex VI.

When this strategy is fully implemented in 2030, U.S. EPA estimates that NO<sub>x</sub> and PM<sub>2.5</sub> emissions in the U.S. will be reduced by approximately 1.2 million tpy and 143,000 tpy, respectively.

### Clean Air Interstate Rule (CAIR)

On May 12, 2005, U.S. EPA promulgated the “Rule to Reduce Interstate Transport of Fine Particulate Matter and Ozone (Clean Air Interstate Rule); Revisions to Acid Rain Program; Revisions to the NO<sub>x</sub> SIP Call”; Final Rule (70 FR 25162). This rule established the requirement for states to adopt rules limiting the emissions of NO<sub>x</sub> and SO<sub>2</sub> and provided a model rule for the states to use in developing their rules to meet federal requirements. The purpose of CAIR was to reduce interstate transport of precursors to fine particles and ozone.

CAIR applied to: (1) any stationary, fossil-fuel-fired boiler or stationary, fossil-fuel-fired combustion turbine, a generator with a nameplate capacity of more than 25 megawatt electrical (MWe) producing electricity for sale; and (2) a unit that qualifies as a cogeneration unit during the 12-month period starting on the date that the unit first produces electricity and continues to qualify as a cogeneration unit, a cogeneration unit serving at any time a generator with a nameplate capacity of more than 25 MWe and supplying in any calendar year more than one-third of the unit’s potential electric output capacity or 219,000 megawatt hours (MWh), whichever is greater, to any utility power distribution system for sale.

---

<sup>6</sup> <http://www.regulations.gov/search/Regs/home.html#documentDetail?R=0900006480ae43a6>

This rule provides annual state caps for NO<sub>x</sub> and SO<sub>2</sub> in two phases, with the Phase I caps for NO<sub>x</sub> and SO<sub>2</sub> starting in 2009 and 2010, respectively. Phase II caps become effective in 2015. U.S. EPA is allowing the caps to be met through a cap and trade program if a state chooses to participate in the program.

In response to U.S. EPA's rulemaking, IDEM adopted a state rule in 2006 based on the model federal rule. IDEM's rule includes annual and seasonal NO<sub>x</sub> trading programs, and an annual SO<sub>2</sub> trading program. This rule requires compliance effective January 1, 2009.

SO<sub>2</sub> emissions from power plants in the 28 eastern states and the District of Columbia covered by CAIR will be cut by 4.3 million tons by 2009 and reduced by an additional 5.4 million tons in 2015. NO<sub>x</sub> emissions will be cut by 1.7 million tons by 2009 and reduced by an additional 1.3 million tons in 2015. The D.C. Circuit court's vacatur of CAIR in July of 2008 and subsequent remand without vacatur of CAIR in December of 2008 directs U.S. EPA to revise or replace CAIR in order to properly address the deficiencies outlined by the court.

Since the court's opinion made it clear that CAIR is deficient and must be revised or replaced, the program cannot be defined as permanent and enforceable for SIP purposes. On July 6, 2010, U.S. EPA proposed the Clean Air Transport Rule to replace CAIR. The Clean Air Transport

Rule will result in even further benefits above and beyond CAIR than what is assumed within the emission inventories and modeling.

Together, these rules will substantially reduce local and regional sources of fine particle precursors. The modeling analyses discussed in Section 7.0 include these rules and show the reductions in annual fine particle concentrations expected to result from the implementation of these rules.

#### 6.5 Controls to Remain in Effect

IDEM commits to maintain the control measures listed above after redesignation, or submit to U.S. EPA, as a SIP revision, any changes to its rules or emission limits applicable to NO<sub>x</sub>, SO<sub>2</sub>, or direct PM<sub>2.5</sub> sources as required for maintenance of the annual standard for fine particles in Lake and Porter counties, Indiana.

Indiana, through IDEM's Office of Air Quality and its Compliance and Enforcement Branch, has the legal authority and necessary resources to actively enforce any violations of its rules or permit provisions. After redesignation, it intends to continue enforcing all rules that relate to the emission of fine particles and fine particle precursors in Lake and Porter counties, Indiana.

#### 6.6 New Source Review Provisions

Indiana has a long standing and fully implemented New Source Review (NSR) program that is outlined in rule 326 IAC 2. The rule includes provisions for the Prevention of Significant Deterioration (PSD) permitting program in 326 IAC 2-2 and the emissions offset rule in 326 IAC

2-3. Indiana's PSD program was conditionally approved on March 3, 2003 (68 FR 9892) and received final approval on May 20, 2004 (69 FR 29071) by U.S. EPA as part of the SIP.

Any emission unit that is not listed in the 2005 emission inventory, or for which credit through the shutdown or curtailment of operations was taken in demonstrating attainment, will not be allowed to construct, reopen, modify, or reconstruct without meeting all applicable permit rule requirements. The review process will be identical to that used for new sources. Once the area is redesignated, OAQ will implement NSR through the PSD program, which requires an air quality analysis to evaluate whether the new source will threaten the NAAQS.

## **7.0 MODELING AND METEOROLOGY**

Although U.S. EPA's Redesignation Guidance does not require modeling for fine particle nonattainment areas seeking redesignation, extensive modeling has been performed for the Chicago Area to determine the effect of national emission control strategies on fine particle levels. These modeling analyses determined that the Chicago Area, including Lake and Porter counties in Northwest Indiana, are significantly impacted by regional transport of fine particles and its precursors, and that regional SO<sub>2</sub> and NO<sub>x</sub> reductions are an effective way to attain the annual standard for fine particles in this area. Future year modeled annual fine particle concentrations are expected to be reduced by 2% to 9% from baseline design values. Examples of these modeling analyses are described in this section and can also be found in Appendix H.

### **7.1 Summary of Modeling Results to Support Rulemakings**

#### **U.S. EPA Modeling for Clean Air Transport Rule 2010<sup>7</sup>**

U.S. EPA performed modeling to support the emission reductions associated with the proposed Clean Air Transport Rule. U.S. EPA used the Comprehensive Air Quality Model with Extension (CAMx version 5), applied to the 2005 meteorology, as processed by the Mesoscale Model (MM5), version 3.7.4. Emissions input into CAMx included SO<sub>2</sub>, NO<sub>x</sub>, VOCs, NH<sub>3</sub>, and direct PM<sub>2.5</sub> for 2005. The modeling was based on the annual fine particle design values calculated from 2003 through 2005, 2004 through 2006, and 2005 through 2007. Future year modeling was conducted, which included the Chicago Area, and the future year design values for 2012 and 2014 were evaluated for attainment of the annual NAAQS for fine particles of 15 µg/m<sup>3</sup>, as shown in Table 7.1. Fine particle concentrations are accounted for by modeling both the base future year emissions and then the emission reductions associated with the Clean Air Transport Rule. U.S. EPA found model performance met suggested benchmark performance goals within or close to the ranges found in other comparable modeling applications.

---

<sup>7</sup> [http://www.epa.gov/airquality/transport/pdfs/TR\\_AQModeling\\_TSD.pdf](http://www.epa.gov/airquality/transport/pdfs/TR_AQModeling_TSD.pdf)

**Table 7.1**  
**Clean Air Transport Rule Modeling Results from U.S. EPA – 2010**


<b>SITE ID</b>	<b>SITE NAME</b>	<b>COUNTY</b>	<b>Design Value 2003 – 2007 (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>Future Design Value 2012 Base (<math>\mu\text{g}/\text{m}^3</math>)</b>	<b>Future Design Value 2014 Base (<math>\mu\text{g}/\text{m}^3</math>)</b>
18-089-0006	East Chicago - Franklin School	Lake	14.27	13.83	13.35
18-089-0027	Griffith	Lake	13.83	13.44	12.97
18-089-1003	Gary – Ivanhoe School	Lake	14.02	13.61	13.15
18-089-2004	Hammond - Purdue	Lake	14.05	13.66	13.19
18-089-2010	Hammond – Clark High School	Lake	13.89	13.47	12.99
18-127-0020	Ogden Dunes	Porter	12.66	12.18	11.73
18-127-0024	Dunes National Lakeshore	Porter	13.21	12.73	12.26
17-031-0022	Chicago – Washington High School	Cook - IL	15.21	14.70	14.22
17-031-0050	Chicago – Southeast Police Station	Cook - IL	14.81	14.32	13.82
17-031-0052	Chicago – Mayfair Pump Station	Cook - IL	15.75	15.16	14.64
17-031-0057	Chicago – Springfield Pump Station	Cook - IL	15.03	14.50	13.98
17-031-0076	Chicago – Com Ed Maintenance Building	Cook - IL	14.89	14.41	13.87
17-031-2001	Blue Island	Cook - IL	14.77	14.28	13.8
17-031-3301	Summit	Cook - IL	15.24	14.73	14.2
17-031-4007	Des Plaines	Cook - IL	12.78	12.45	11.92
17-031-4201	Northbrook	Cook - IL	12.76	12.44	11.91
17-031-6005	Cicero	Cook - IL	15.48	14.92	14.40
17-043-4002	Naperville	Du Page - IL	13.82	13.51	12.95
17-089-0003	Elgin	Kane - IL	13.32	12.94	12.36
17-089-0007	Aurora	Kane - IL	14.34	13.95	13.39
17-097-1007	Zion	Lake - IL	11.81	11.57	11.06
17-111-0001	Cary	McHenry - IL	12.40	12.08	11.53
17-197-1002	Joliet	Will - IL	13.63	13.33	12.79
17-197-1011	Braidwood	Will - IL	11.52	11.26	10.77

Modeling results show that the base future year modeling with emission reductions from the Clean Air Transport Rule will account for a 0.4 to 0.5  $\mu\text{g}/\text{m}^3$  decrease in concentrations for 2012, as well as a 0.9 to 1.0  $\mu\text{g}/\text{m}^3$  decrease in concentrations for 2014 in the Chicago Area.

Only one  $\text{PM}_{2.5}$  monitor in the Chicago Area had a modeled design value above the annual standard for fine particles for 2012. The Chicago - Mayfair Pump Station monitor (ID 17-031-0052) had a 2003 through 2007 design value of 15.75  $\mu\text{g}/\text{m}^3$ . Review of this monitor's most current 2006 through 2010 design value shows that the design value has decreased by 2.42  $\mu\text{g}/\text{m}^3$  to 13.33  $\mu\text{g}/\text{m}^3$ . Therefore, modeling results should fall by a similar value and be well below the annual standard for fine particles. Design values from 2003 through 2007 to 2006 through 2010 have fallen by an average of 2.1  $\mu\text{g}/\text{m}^3$  throughout the Chicago Area. Graph 7.1 shows the downward trend of the design values from 2000 through 2010 from the  $\text{PM}_{2.5}$  monitors located in Lake and Porter counties, Indiana. In fact, all design values are below the annual NAAQS for fine particles.



**Graph 7.1**  
**PM<sub>2.5</sub> Design Value Trends for Lake and Porter Counties, Indiana, 2000 to 2010**



Results of the U.S. EPA's Clean Air Transport Rule modeling show that the Chicago Area will attain the annual fine particle NAAQS in 2012 with modeled impacts reduced by 3% to 4%, and will remain below 15  $\mu\text{g}/\text{m}^3$ . With further reductions projected in the Clean Air Transport Rule for 2014, all design values decrease by 6% to 7% and the area will continue to attain the annual NAAQS for fine particles.

#### LADCO Modeling for Clean Air Interstate Rule (CAIR)


LADCO conducted modeling to determine the impact of CAIR in the Midwest. LADCO's modeling used the CAMx model applied to the year 2005 meteorology, as processed by the MM5. Emissions input into CAMx included SO<sub>2</sub>, NO<sub>x</sub>, VOCs, NH<sub>3</sub>, and direct PM<sub>2.5</sub> for 2005. The modeling was based on 2003 through 2007 design values. Future year modeling for 2012 and 2018 was conducted and the future year design values were determined without the emission reductions associated with CAIR (Round 6), as shown in Table 7.2. The Clean Air Transport Rule is expected to provide reductions above and beyond CAIR.

**Table 7.2**  
**LADCO Round 6 Modeling Results**  
**for Annual PM<sub>2.5</sub> – Without CAIR Emission Reductions**  
**(Using 2003 – 2007 Design Values)**

SITE ID	SITE NAME	COUNTY	Design Value 2003-2007 (µg/m <sup>3</sup> )	Base-case 2012 (µg/m <sup>3</sup> )	Base-case 2018 (µg/m <sup>3</sup> )
18-089-0006	East Chicago – Franklin School	Lake	14.3	13.7	13.6
18-089-0027	Griffith	Lake	13.8	13.0	12.8
18-089-1003	Gary – Ivanhoe School	Lake	14.0	13.1	12.9
18-089-2004	Hammond – Purdue	Lake	14.1	13.4	13.3
18-089-2010	Hammond – Clark High School	Lake	13.9	13.3	13.2
18-127-0020	Ogden Dunes	Porter	12.7	11.9	11.7
18-127-0024	Dunes National Lakeshore	Porter	13.2	12.4	12.1
17-031-0022	Chicago – Washington High School	Cook - IL	15.2	14.6	14.6
17-031-0050	Chicago – Southeast Police Station	Cook - IL	14.8	14.2	14.1
17-031-0052	Chicago – Mayfair Pump Station	Cook - IL	15.8	14.9	14.7
17-031-0057	Chicago – Springfield Pump Station	Cook - IL	15.0	14.4	14.3
17-031-0076	Chicago – Com Ed Maintenance Building	Cook - IL	14.9	14.3	14.2
17-031-2001	Blue Island	Cook - IL	14.8	14.2	14.1
17-031-3301	Summit	Cook - IL	15.3	14.7	14.6
17-031-4007	Des Plaines	Cook - IL	12.8	12.1	11.8
17-031-4201	Northbrook	Cook - IL	12.8	12.1	11.8
17-031-6005	Cicero	Cook - IL	15.5	14.9	14.8
17-043-4002	Naperville	Du Page - IL	13.8	13.0	12.7
17-089-0003	Elgin	Kane - IL	13.3	12.5	12.1
17-089-0007	Aurora	Kane - IL	14.8	13.9	13.5
17-097-1007	Zion	Lake - IL	11.8	11.2	10.9
17-111-0001	Cary	McHenry - IL	12.4	11.5	11.2
17-197-1002	Joliet	Will - IL	13.7	12.9	12.5
17-197-1011	Braidwood	Will - IL	11.5	10.7	10.5

Results of the LADCO Round 6 modeling show that the Chicago Area would attain the annual NAAQS for fine particles by 2012. As shown in Table 7.2, future year modeled annual fine particle concentrations for 2012 will be 4% to 6% lower than baseline annual fine particle design values and 5% to 8% lower in 2018 and will continue to decrease thereafter. A graphical representation of the results for the fine particle monitors located in Lake and Porter counties, Indiana is shown in Graph 7.2.

**Graph 7.2**  
**LADCO Round 6 Modeling Results for Lake and Porter Counties, Indiana**  
**for 2012 and 2018**



It should be noted that PM<sub>2.5</sub> monitors exist at Gary – IITRI and Gary – Burr Street, however these are source oriented monitors used to compare to the daily standard only and are not used to determine annual fine particle attainment status.

## 7.2 LADCO Round 5 Speciated Modeled Attainment Test Results

The Speciated Modeled Attainment Test (SMAT) is the attainment test for annual fine particles. To determine the future year annual fine particle concentrations, speciated data is calculated. The different species that were modeled and are associated with fine particles include sulfates, nitrates, organic carbon, elemental carbon, ammonium, particle bound water, “other” primary inorganic fine particles, and passively collected mass. The SMAT results from LADCO’s Round 5 modeling are listed below. Round 5 modeled the emission reductions associated with CAIR so the results reflect the impacts from CAIR. Percent ranges of the model results from the seven active fine particle monitors in Lake and Porter counties, Indiana were broken down into the speciated constituents of fine particle emissions. The percent change from the observed speciated data in 2005 to the future year modeled results for 2009 are listed in Table 7.3. While these modeling results are outdated, this weight of evidence demonstration shows the reduction potential from national emission control measures for the PM<sub>2.5</sub> species in Northwest Indiana. An updated analysis will be conducted at a later time, using more current emissions and meteorological data and for a relevant future year projection.

**Table 7.3**  
**LADCO Round 5 SMAT Modeling Results for Lake and Porter Counties, Indiana**  
**(Percent decrease from observed to modeled concentrations)**

<b>Species of PM<sub>2.5</sub></b>	<b>2009</b>
Sulfates	9% - 10%
Nitrates	5% - 6%
Organic Carbon	1% - 2%
Elemental Carbon	10% - 20%
Ammonium	6% - 7%
Particle Bound Water	9%



The results demonstrate that elemental carbon, sulfate, particle bound water, ammonium, and nitrates concentration decreases are projected to be at least 6% in the future year 2009. Lesser organic carbon reductions are projected to occur, up to 2%. LADCO modeling shows good performance for sulfates and elemental carbon predicted baseline concentrations, slight over-prediction for nitrate concentrations, and under-predictions of organic carbon concentrations. Overall, model performance is adequate for SIP planning and gives a good idea of the effects of emission reductions from national emission control measures on Lake and Porter counties, Indiana.

### 7.3 LADCO Round 5 Particulate Source Apportionment Results


Particulate Source Apportionment (PSAT) modeling was conducted by LADCO. The results of the PSAT Round 5 modeling show the regional contributions by emission sectors on each monitor that was modeled. Chart 7.1 displays the PSAT modeling results for the East Chicago – Franklin School, Griffith, Gary – Ivanhoe School, Gary – Clark High School, and Hammond – Purdue fine particle monitors in Lake County, Indiana and the Dunes National Lakeshore and Ogden Dunes fine particle monitors in Porter County, Indiana. Emission contributions from Illinois’ portion of the nonattainment area was the biggest regional contributor to the East Chicago – Franklin School, Griffith, Gary – Clark High School, and Hammond – Purdue fine particle monitors. Emission contributions from both the Indiana and Illinois portions of the nonattainment area were the biggest regional contributors to the Dunes National Lakeshore, Ogden Dunes, and Gary – Ivanhoe High School fine particle monitors. Indiana and Illinois were the biggest state contributors to all of the monitors depicted in Chart 7.1.

The PSAT Round 5 modeling results indicate the majority of Indiana’s emission sector contributions to fine particle concentrations come from EGUs, ammonium emission sources, mobile, and off-road (including marine, aircraft, and railroad), and non-EGU sources. These results are considered to be representative of Lake and Porter counties, Indiana as EGU, ammonium, and non-EGU emissions impact the entire area.


## Chart 7.1 Regional/Emission Sector Particulate Source Apportionment Technology (PSAT) Results




### Gary – Ivanhoe School PM<sub>2.5</sub> Monitor, Lake County, Indiana




### Hammond – Purdue PM<sub>2.5</sub> Monitor, Lake County, Indiana




### Gary – Clark High School PM<sub>2.5</sub> Monitor, Lake County, Indiana



### Dunes National Lakeshore PM<sub>2.5</sub> Monitor, Porter County, Indiana






The following pie charts depict the specie contributions to fine particle concentrations at the Lake and Porter counties, Indiana monitors. The pie charts include both the observed 2005 contributions and 2009 modeled contributions for each monitor. Since the monitors are in close proximity to each other, results are fairly similar in the distribution of specie concentrations among the monitors. Charts 7.2 through 7.7 cover the fine particle monitors in Lake County while Charts 7.8 and 7.9 cover the monitors in Porter County with the highest monitored concentrations that are used to determine compliance with the annual NAAQS for fine particles.


The speciation listed in the pie charts include SO<sub>4</sub> – sulfate mass, NO<sub>3</sub> – nitrate mass, OC – organic carbon mass, EC – elemental carbon mass, Soil – crustal material mass, NH<sub>4</sub> – ammonium mass, PBW – particles bound water mass, and BLAN – passively collected mass.



**Chart 7.2**  
**Modeled Contributions by Specie at the East Chicago –**  
**Franklin School PM<sub>2.5</sub> Monitor**  
**(Observed Concentrations = 13.8 µg/m<sup>3</sup>) (Modeled Concentrations =13.2 µg/m<sup>3</sup>)**




**Chart 7.3**  
**Modeled Contributions by Specie at the Griffith PM<sub>2.5</sub> Monitor**  
**(Observed Concentrations = 13.3 µg/m<sup>3</sup>) (Modeled Concentrations =12.4 µg/m<sup>3</sup>)**




**Chart 7.4**

**Modeled Contributions by Specie at the Gary - Madison Street PM<sub>2.5</sub> Monitor**  
 (Observed Concentrations = 13.5 µg/m<sup>3</sup>) (Modeled Concentrations =12.3 µg/m<sup>3</sup>)




**Chart 7.5**

**Modeled Contributions by Specie at the Gary - Ivanhoe School PM<sub>2.5</sub> Monitor**  
 (Observed Concentrations = 15.2 µg/m<sup>3</sup>) (Modeled Concentrations =12.9 µg/m<sup>3</sup>)




**Chart 7.6**

**Modeled Contributions by Specie at the Hammond - Purdue PM<sub>2.5</sub> Monitor**  
**(Observed Concentrations = 13.6 µg/m<sup>3</sup>) (Modeled Concentrations =12.7 µg/m<sup>3</sup>)**




**Chart 7.7**


**Modeled Contributions by Specie at the Hammond – Clark High School PM<sub>2.5</sub> Monitor**  
**(Observed Concentrations = 13.4 µg/m<sup>3</sup>) (Modeled Concentrations =12.5 µg/m<sup>3</sup>)**



**Chart 7.8**  
**Modeled Contributions by Specie at the Dunes National Lakeshore PM<sub>2.5</sub> Monitor**  
**(Observed Concentrations = 12.1 µg/m<sup>3</sup>) (Modeled Concentrations =11.2 µg/m<sup>3</sup>)**



**Chart 7.9**  
**Modeled Contributions by Specie at the Ogden Dunes PM<sub>2.5</sub> Monitor**  
**(Observed Concentrations = 12.6 µg/m<sup>3</sup>) (Modeled Concentrations =11.5 µg/m<sup>3</sup>)**



Results of the Round 5 PSAT modeling for the Lake and Porter counties, Indiana fine particle monitors show the highest pollutant contributors to base-case and future year fine particle concentrations are sulfate, organic carbon, ammonium, and nitrate. Future year modeling shows decreases in sulfates, nitrates, and ammonium. The future year modeling did show slight increases in organic carbon and soils from the base-case modeled concentrations.

#### 7.4 Summary of Existing Modeling Results

U.S. EPA and LADCO modeling for future year design values have consistently shown that existing national emission control measures will bring Lake and Porter counties, Indiana into attainment of the annual NAAQS for fine particles. Emission control measures to be implemented in the next several years will provide even greater assurance that air quality will continue to meet the standard into the future. Modeling support for CAIR has shown that future year design values for Lake and Porter counties, Indiana will continue to attain the annual standard for fine particles with modeled future year design values below  $15 \mu\text{g}/\text{m}^3$ . U.S. EPA future year modeling of national emission control strategies will ensure that Lake and Porter counties, Indiana will maintain lower fine particle concentrations with an increasing margin of safety.

#### 7.5 Meteorological Analysis for Lake and Porter Counties, Indiana

Meteorological conditions are one of the most important factors that influence development and transport of fine particles. Stagnant surface conditions during any time of the year and upper air ridging provides conducive conditions for development and transport of fine particles. Ultimately, passage of surface cold fronts with a clean air mass change will lower fine particle readings in Lake and Porter counties, Indiana.

#### 7.6 Surface Air Conditions Present During High Fine Particle Concentration Days

Higher annual concentrations of fine particles tend to correlate with warmer temperatures and lighter wind speeds, although high fine particle episodes can occur in the summer or winter. It should be noted that higher annual fine particle concentrations are driven by individual days with higher fine particle concentrations throughout the monitored year. Therefore, it is difficult to attribute higher fine particle concentrations to annualized meteorological rankings. Review of several of the higher fine particle concentration episodes over the past few years reveal that conditions were hot in the summer with temperatures in the middle 80's Fahrenheit or higher and average wind speeds were fairly light. Fall and winter days with higher fine particle concentrations had near normal temperatures but wind speeds were very light, and humidity was higher.

#### 7.7 Upper Air Conditions Present During High Fine Particle Concentration Days

Upper air ridges and more stagnant surface wind conditions predominately affect development and build up of fine particles. Slow moving upper air ridges can effectively suppress mixing within the many levels of the atmosphere and cause pollutants to build up over time. Inversions or increases in temperature with a rise in altitude will prevent mixing with air from the upper atmosphere. These conditions can occur at any time of the year and are evident in elevated fine particle episodes in spring, summer, fall, and winter months. Review of surface and upper air features of higher fine particle concentration days showed stagnant surface conditions and upper air ridges existed on those days and helped in the buildup of fine particle concentrations.

## 7.8 Analyses of Atmospheric Conditions During High Fine Particle Concentration Days

Analyses have been conducted to determine the atmospheric conditions that are most prevalent during higher fine particle concentration days in Indiana. LADCO applied a Classification and Regression Tree (CART) analysis to data from Indiana that correlated different levels of fine particle concentrations to meteorological conditions from 1999 to 2004 (Donna Kenski, 2005). This type of analysis evaluates meteorological conditions, such as temperature, pressure, wind speed, wind direction, relative humidity, and dew point temperatures at the surface, as well as morning and evening mixing heights in the upper atmosphere which were present when higher concentrations of fine particles were monitored. Results of this CART analysis indicated factors that played a larger role in higher fine particle concentrations in Indiana were warm-weather conditions with high dew points, southwest winds, and high evening mixing heights. Previous day's concentrations of fine particles play a key role in higher impacts as well.

Fine particles are made up of several constituents, including direct PM<sub>2.5</sub>, sulfates, nitrates, ammonium, organic carbon, and elemental carbon. Depending on the time of the year, concentrations of fine particle constituents vary, with nitrates being more prevalent in the winter and sulfates more prevalent in the summer. Sulfate and nitrates emission reductions have the biggest impact on lower future year fine particle concentrations.

## 7.9 Summary of Air Quality Index Days in Lake and Porter Counties, Indiana

An analysis was conducted to review the daily fine particle concentrations over a year to determine the Air Quality Index (AQI) trends. Chart 7.10 shows by year (2000 through 2010), the percentage of days during the calendar year which fine particle concentrations reached the AQI ranges for “Good” (0 to 15.3  $\mu\text{g}/\text{m}^3$ ), “Moderate” (15.4  $\mu\text{g}/\text{m}^3$  to 40.4  $\mu\text{g}/\text{m}^3$ ), and “Unhealthy for Sensitive Groups (USG)” (40.5  $\mu\text{g}/\text{m}^3$  to 65.4  $\mu\text{g}/\text{m}^3$ ). There were no days that fine particle levels reached the “Unhealthy” level of 65.5  $\mu\text{g}/\text{m}^3$  to 150.4  $\mu\text{g}/\text{m}^3$ .

**Chart 7.10**  
**Distribution of PM<sub>2.5</sub> Concentration Days on the AQI Levels of Health Concern,**  
**Lake and Porter Counties, Indiana**




Table 7.4 shows how the years ranked for the three AQI ranges. The year 2009 had the most “Good” concentration days during the 11-year period analyzed (2000 through 2010). The year 2000 had the most “Moderate” concentration days and the years 2001, 2003, and 2005 had the most “USG” concentration days with no days recorded in 2006, 2008, and 2009. As can be seen, weather plays a large role in fine particle concentration development and transport as 2001, 2002, 2005, 2007, and 2010 were warmer than normal summers which translated to moderate and unhealthy for sensitive group levels of air quality.

**Table 7.4**  
**Ranking and Percentage of Highest Number of Days**  
**at AQI Levels of Health Concern**

<b>Ranking</b>	<b>Good</b>	<b>Moderate</b>	<b>Unhealthy for Sensitive Group</b>
<b>1<sup>st</sup></b>	<b>2009 – 78%</b>	<b>2000 – 46%</b>	<b>2001 – 3%</b>
<b>2<sup>nd</sup></b>	<b>2008 – 72%</b>	<b>2001 – 42%</b>	<b>2005 – 3%</b>
<b>3<sup>rd</sup></b>	<b>2004 – 69%</b>	<b>2005 – 40%</b>	<b>2003 – 3%</b>
<b>4<sup>th</sup></b>	<b>2007 – 69%</b>	<b>2002 – 39%</b>	<b>2000 – 2%</b>
<b>5<sup>th</sup></b>	<b>2006 – 66%</b>	<b>2003 – 36%</b>	<b>2010 – 2%</b>
<b>6<sup>th</sup></b>	<b>2010 – 64%</b>	<b>2006 – 34%</b>	<b>2002 – 2%</b>
<b>7<sup>th</sup></b>	<b>2003 – 63%</b>	<b>2010 – 34%</b>	<b>2007 – 2%</b>
<b>8<sup>th</sup></b>	<b>2002 – 60%</b>	<b>2007 – 30%</b>	<b>2004 – 1%</b>
<b>9<sup>th</sup></b>	<b>2005 – 59%</b>	<b>2004 – 30%</b>	
<b>10<sup>th</sup></b>	<b>2001 – 56%</b>	<b>2008 – 28%</b>	
<b>11<sup>th</sup></b>	<b>2000 – 52%</b>	<b>2009 – 22%</b>	

#### 7.10 Summary of Meteorological Analysis for Lake and Porter Counties, Indiana

Annual fine particle concentrations in Lake and Porter counties, Indiana are driven by higher fine particle concentration days that can occur during any time of the year. Conditions that are most prevalent during higher fine particle concentration days are lighter winds, higher relative humidity, and above average temperatures in the summer, and near normal temperatures in the fall, winter, or spring. Approximately 70% of the days when PM<sub>2.5</sub> concentrations were in the USG range occurred in the summer months with maximum high temperatures of 80°F or above. Weather plays a large role in fine particle concentration development and transport as 2001, 2002, 2005, and 2006 were warmer than normal summers which translated to more days of moderate and unhealthy for sensitive group levels of air quality. Upper air weather patterns generally include ridging over the area with stagnant conditions at the surface caused by lower mixing heights and stable conditions for summer episodes and ridging or troughs over the area in the fall, winter, or spring episodes. Surface winds from any direction can transport pollutants from surrounding areas into Lake and Porter counties, Indiana. Nitrates are bigger contributors to fine particle concentrations in the winter and sulfates are bigger contributors to fine particle concentrations in the summer.

## 8.0 CORRECTIVE ACTIONS

### 8.1 Commitment to Revise Plan

As noted in Section 4.6, IDEM commits to review and revise, as appropriate, its Maintenance Plan eight years after redesignation, as required by Section 175A of the CAA.



## 8.2 Commitment for Contingency Measures

IDEM will monitor fine particle concentrations to determine whether trends indicate higher values or whether emissions appear to be increasing. If it is determined that fine particle levels and emissions are increasing and action is necessary to reverse that trend, IDEM will take action to reverse the noted trend, prior to a violation of the standard occurring.

IDEM commits to adopt and expeditiously implement necessary corrective action in accordance with an Action Level Response described below.

### Action Level Response

An Action Level Response shall be prompted whenever a violation of the standard (three year average annual arithmetic mean value of  $15.1 \mu\text{g}/\text{m}^3$  or greater) occurs. In the event that the Action Level is triggered and is not found to be due to an atypical unfavorable meteorological condition, exceptional event, malfunction, or noncompliance with a permit condition or rule requirement, IDEM will determine additional control measures needed to assure future attainment of the annual NAAQS for fine particles. In this case, measures that can be implemented in a short time will be selected in order to be in place within eighteen months from the end of the year that prompted the Action Level Response.

### Control Measure Selection and Implementation

Adoption of any additional control measures is subject to the necessary administrative and legal processes. This process will include publication of notices, an opportunity for public hearing, and other measures required by Indiana law for rulemaking by state environmental boards.

If a new measure or control is already promulgated and scheduled to be implemented at the federal or state level, and that measure or control is determined to be sufficient to address the upward trend in air quality, additional local measures may be unnecessary. Furthermore, IDEM will submit to U.S. EPA an analysis to demonstrate the proposed measures are adequate to return the area to attainment.

## 8.3 Contingency Measures

Contingency measures to be considered will be selected from a comprehensive list of measures deemed appropriate and effective at the time the selection is made. Listed below are example measures that may be considered. The selection of measures will be based upon cost-effectiveness, emission reduction potential, economic and social considerations, or other factors that IDEM deems appropriate. IDEM will solicit input from interested and affected persons in the maintenance area prior to selecting appropriate contingency measures. All of the listed contingency measures are potentially effective or proven methods of obtaining significant reductions of fine particle precursor emissions. Because it is not possible at this time to determine what control measure will be appropriate at an unspecified time in the future, the list of contingency measures outlined below is not comprehensive. IDEM anticipates that if

contingency measures should ever be necessary, it is unlikely that a significant number (i.e., all those listed below) will be required.

- 1) Vehicle inspection and maintenance program enhancements (increase weight limit, addition of diesel vehicles, etc.).
- 2) Alternative fuel and diesel retrofit programs for fleet vehicle operations.
- 3) Require NO<sub>x</sub> or SO<sub>2</sub> emission offsets for new and modified major sources.
- 4) Require NO<sub>x</sub> or SO<sub>2</sub> emission offsets for new and modified minor sources.
- 5) Increase the ratio of emission offsets required for new sources.
- 6) Require NO<sub>x</sub> or SO<sub>2</sub> controls on new minor sources (less than 100 tons).
- 7) Wood stove change out program.
- 8) Require increased recovery efficiency at sulfur recovery plants.
- 9) Various emission reduction measures or dust suppressant for unpaved roads and/or parking lots.
- 10) Idling Restrictions.
- 11) Broader geographic applicability of existing measures.
- 12) One or more transportation control measures sufficient to achieve at least a 0.5% reduction in actual area-wide precursor emissions. Transportation measures will be selected from the following, based upon the factors listed above, after consultation with affected local governments:
  - a) Trip reduction programs, including, but not limited to, employer-based transportation management plans, area wide rideshare programs, work schedule changes, and telecommuting.
  - b) Transit improvements.
  - c) Traffic flow improvements.
  - d) Other new or innovative transportation measures not yet in widespread use that affected state and local governments deem appropriate.

No contingency measure shall be implemented without providing the opportunity for full public participation during which the relative costs and benefits of individual measures, at the time they are under consideration, can be fully evaluated.

## **9.0 PUBLIC PARTICIPATION**

Indiana published notification for a public hearing and solicitation for public comment concerning the draft Redesignation Petition and Maintenance Plan on the IDEM Web site on April 14, 2011, with publication in the following newspapers on the following dates:

- 1) The Indianapolis Star, Indianapolis, Indiana (April 15, 2011).
- 2) The Times, Munster, Indiana (April 14, 2011).
- 3) The Post Tribune, Merrillville, Indiana (April 14, 2011).

A public hearing to receive comments concerning the redesignation request was conducted on May 18, 2011, at the Lake County Public Library-Highland Branch in Highland, Indiana. The public comment period closed on May 20, 2011. One comment letter was received during the public comment period. Appendix K includes a copy of the public notice, public hearing script, certifications of newspaper publication of the public notice, the official transcript from the public hearing, and a detailed summary of and response to substantive comments.

## **10.0 CONCLUSIONS**

Lake and Porter counties, Indiana, have attained the annual NAAQS for fine particles. This petition demonstrates that Lake and Porter counties, Indiana, have complied with the applicable provisions of the CAA regarding redesignation of nonattainment areas for fine particles. IDEM has prepared a State Implementation and Maintenance Plan that meets the requirement of Section 110(a)(1) of the CAA.

Indiana has performed an analysis that shows the air quality improvements are due to permanent and enforceable measures and that additional significant regional NO<sub>x</sub> and SO<sub>2</sub> reductions following implementation of the Phase II NO<sub>x</sub> SIP Call rule, and CAIR or its replacement rule will ensure continued compliance (maintenance) with the standard. Furthermore, emission projections indicate that NO<sub>x</sub> and SO<sub>2</sub> emissions will continue to decline, ensuring that the area continues to maintain compliance with the standard and provide for an increased margin of safety. Based on this presentation, Lake and Porter counties, Indiana, meet the requirements for redesignation under the CAA (Section 107(d)(3)) and U.S. EPA guidance for fine particles.

Consistent with the authority granted to U.S. EPA, the State of Indiana requests that Lake and Porter counties, Indiana, be redesignated to attainment for the annual fine particles standard simultaneously with U.S. EPA approval of this Indiana State Implementation and Maintenance Plan.

# **APPENDIX A**

**Air Quality System (AQS) and Indiana  
Department of Environmental Management  
(IDEM) Fine Particle Monitor Data Values for the  
Chicago Area (2000 – 2010)**

**This page left intentionally blank.**

## Monitoring Data for Indiana's Portion of the Chicago Area

SITE ID	COUNTY	SITE NAME	YEAR	Annual Average $\mu\text{g}/\text{m}^3$	2008-2010 Average $\mu\text{g}/\text{m}^3$
18-089-0006	Lake	East Chicago - Franklin School	2008	11.95	<b>11.9</b>
			2009	11.34	
			2010	12.48	
18-089-0027	Lake	Griffith	2008	11.69	<b>11.7</b>
			2009	11.00	
			2010	12.39	
18-089-2004	Lake	Hammond - Purdue	2008	11.66	<b>N/A*</b>
			2009	N/A*	
			2010	12.30	
18-089-2010	Lake	Hammond - Clark High School	2008	12.42	<b>11.7</b>
			2009	10.80	
			2010	11.90	
18-089-0031	Lake	Gary - Madison Street	2008	12.27	<b>12.4</b>
			2009	12.12	
			2010	12.90	
18-127-0024	Porter	Ogden Dunes	2008	10.89	<b>11.2</b>
			2009	11.29	
			2008	11.56	

\*The Hammond - Purdue monitoring site was physically removed to accommodate heating and cooling building repairs on the roof which were conducted from February 12, 2009 through November 18, 2009. As a result, the 2009 monitoring data and the 2008 through 2010 three year design value are considered incomplete.

Site ID	County	Site Name	Yearly Annual Means										
			2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
18-089-0006	Lake	East Chicago - Franklin School	15.76	16.11	14.92	14.60	13.18	15.76	13.18	14.44	11.95	11.34	12.48
18-089-0022	Lake	Gary - IITRI	17.38	17.99	16.43	16.64	16.11	18.25	13.57	15.06	12.39	12.48	13.61
18-089-0026	Lake	Gary - Burr Street	17.24	18.19	17.67	17.38	16.53	18.72	14.71	15.87 <sup>E</sup>	13.82	13.37	14.10
18-089-0027	Lake	Griffith	14.04	15.18	14.60	14.10	12.82	15.46	12.29	13.17	11.69	11.00	12.39
18-089-1003	Lake	Gary - Ivanhoe School	15.33	14.98	15.22	14.14	12.92	15.71	12.57	14.01			
18-089-1016	Lake	Gary - Federal Building	16.03	16.26	15.92								
18-089-2004	Lake	Hammond - Purdue	14.96	15.38	14.70	14.55	13.26	15.40	12.67	13.80	11.66	N/A*	12.30
18-089-2010	Lake	Hammond - Clark High School	14.34	15.55	14.88	14.26	12.47	15.59	12.79	13.68	12.42	10.80	11.90
18-089-0031	Lake	Gary - Madison Street						16.61	13.30	14.55	12.27	12.12	12.90
18-127-0020	Porter	Dunes National Lakeshore	13.53	13.62	13.24	13.19	11.84	14.00	11.02	13.04			
18-127-0024	Porter	Ogden Dunes	14.55	14.18	14.20	12.94	12.38	14.59	11.81	13.79	10.89	11.29	11.56

Source-Oriented Site NOT used for comparison to the annual standard for fine particles.

\*The Hammond - Purdue monitoring site was physically removed to accommodate heating and cooling building repairs on the roof which were conducted from February 12, 2009 through November 18, 2009. As a result, the 2009 monitoring data is considered incomplete.

<sup>E</sup> Exceptional event data removed from calculations.

The Gary - Ivanhoe School monitor was discontinued on December 31, 2007.

The Gary - Federal Building monitor was discontinued in September 2002.

The Gary - Madison Street monitor began operation on July 1, 2005.

The Dunes National Lakeshore monitor was discontinued on December 31, 2007.

Site ID	County	Site Name	Three Year Design Values								
			00-02	01-03	02-04	03-05	04-06	05-07	06-08	07-09	08-10
18-089-0006	Lake	East Chicago - Franklin School	15.6	15.2	14.2	14.5	14.0	14.5	13.2	12.6	11.9
18-089-0022	Lake	Gary - IITRI	17.3	17.0	16.4	17.0	16.0	15.6	13.7	13.3	12.8
18-089-0026	Lake	Gary - Burr Street	17.7	17.7	17.2	17.5	16.7	16.4	14.8	14.4	13.8
18-089-0027	Lake	Griffith	14.6	14.6	13.8	14.1	13.5	13.6	12.4	12.0	11.7
18-089-1003	Lake	Gary - Ivanhoe School	15.2	14.8	14.1	14.3	13.7	14.1			
18-089-1016	Lake	Gary - Federal Building	16.1								
18-089-2004	Lake	Hammond - Purdue	15.0	14.9	14.2	14.4	13.8	14.0	12.7	N/A*	N/A*
18-089-2010	Lake	Hammond - Clark High School	14.9	14.9	13.9	14.1	13.6	14.0	13.0	12.3	11.7
18-089-0031	Lake	Gary - Madison Street				16.6	15.0	14.8	13.4	13.0	12.4
18-127-0020	Porter	Dunes National Lakeshore	13.5	13.4	12.8	13.0	12.3	12.7			
18-127-0024	Porter	Ogden Dunes	14.3	13.8	13.2	13.3	12.9	13.4	12.2	12.0	11.2
Value above the annual standard for fine particles											

Source-Oriented Site NOT used for comparison to the annual standard for fine particles.

Red Text Indicates Incomplete Data

Blue Text Indicates Design Value Based on One Year of Data

Text Indicates Design Value Based on Two Years of Data

\*The Hammond - Purdue monitoring site was physically removed to accommodate heating and cooling building repairs on the roof which were conducted from February 12, 2009 through November 18, 2009. As a result, the 2007 through 2009 and 2008 through 2010 three year design values are considered incomplete.


The Gary - Ivanhoe School monitor was discontinued on December 31, 2007.

The Gary - Federal Building monitor was discontinued in September 2002.

The Gary - Madison Street monitor began operation on July 1, 2005.

The Dunes National Lakeshore monitor was discontinued on December 31, 2007.


### Annual Fine Particle Trends for Indiana's Portion of the Chicago Area, 2006 through 2010



Red line represents the annual standard for fine particles of 15.0  $\mu\text{g}/\text{m}^3$

The Hammond - Purdue monitoring site was physically removed to accommodate heating and cooling building repairs on the roof which were conducted from February 12, 2009 through November 18, 2009. As a result, the 2009 monitoring data is considered incomplete.

### Fine Particle Design Values for Indiana's Portion of the Chicago Area, 2006 through 2010



Red line represents the annual standard for fine particles of 15  $\mu\text{g}/\text{m}^3$

The Hammond - Purdue monitoring site was physically removed to accommodate heating and cooling building repairs on the roof which were conducted from February 12, 2009 through November 18, 2009. As a result, the 2007 through 2009 and 2008 through 2010 three year design values are considered incomplete.



**This page left intentionally blank.**

## Monitoring Data for Illinois' Portion of the Chicago Area

SITE ID	COUNTY	SITE NAME	YEAR	Annual Average $\mu\text{g}/\text{m}^3$	2008-2010 Average $\mu\text{g}/\text{m}^3$
17-031-0022	Cook	Chicago - Washington High School	2008	12.54	12.7
			2009	11.62	
			2010	14.04	
17-031-0050	Cook	Chicago - Southeast Police Station	2008	11.80	11.8
			2009	10.99	
			2010	12.47	
17-031-0052	Cook	Chicago - Mayfair Pump Station	2008	12.18	12.5
			2009	12.69	
			2010	12.57	
17-031-0057	Cook	Chicago - Springfield Pump Station	2008	12.03	11.8
			2009	11.33	
			2010	12.03*	
17-031-0076	Cook	Chicago - Com Ed Maintenance Building	2008	11.89	11.8
			2009	11.12	
			2010	12.25	
17-031-2001	Cook	Blue Island	2008	12.50	11.9
			2009	11.68	
			2010	11.59	
17-031-3301	Cook	Summit	2008	12.03	12.0
			2009	11.62	
			2010	12.23	
17-031-4007	Cook	Des Plaines	2008	11.35	11.0
			2009	11.02	
			2010	10.60	
17-031-4201	Cook	Northbrook	2008	10.09	9.6
			2009	9.33	
			2010	9.34	
17-031-6005	Cook	Cicero	2008	13.25*	12.4
			2009	11.98*	
			2010	11.90	
17-043-4002	DuPage	Naperville	2008	11.28	10.9
			2009	9.83	
			2010	11.67	
17-089-0003	Kane	Elgin	2008	10.79	10.6
			2009	9.61	
			2010	11.29	
17-089-0007	Kane	Aurora	2008	10.34	10.6
			2009	10.01	
			2010	11.44	
17-097-1007	Lake	Zion	2008	9.34	9.3
			2009	8.83	
			2010	9.66	
17-111-0001	McHenry	Cary	2008	10.10	10.0
			2009	9.65	
			2010	10.24	
17-197-1002	Will	Joliet	2008	11.66	11.3
			2009	10.52	
			2010	11.83	
17-197-1011	Will	Braidwood	2008	10.31	9.7
			2009	8.73	
			2010	10.02	

\*Indicates that the mean does not satisfy summary criteria

Red Text Indicates Incomplete Data

Site ID	County	Site Name	Yearly Annual Means										
			2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010
17-031-0022	Cook	Chicago - Washington High School	17.87	17.11	15.31	15.58	14.16	16.94	13.23	15.73	12.54	11.62	14.04
17-031-0050	Cook	Chicago - Southeast Police Station	16.46*	18.12*	15.47	15.36	13.76	16.56	13.33	14.14	11.80	10.99	12.47
17-031-0052	Cook	Chicago - Mayfair Pump Station	18.33	19.39	16.51	15.85	15.27	16.97	14.50	15.49	12.18	12.69	12.57
17-031-0057	Cook	Chicago - Springfield Pump Station	17.31	16.24	15.21	15.61	13.75	16.66	13.51	15.18	12.03	11.33	12.03*
17-031-0076	Cook	Chicago - Com Ed Maintenance Building	16.61	15.53*	15.66	14.84	14.15	16.56	13.48	14.30	11.89	11.12	12.25
17-031-1016	Cook	McCook	20.20	20.85	17.71	16.69	16.73	18.30	15.58	15.63	12.88	12.60	12.58
17-031-2001	Cook	Blue Island	16.82	17.11	15.18*	14.92	14.12	16.39	13.18	14.32	12.50	11.68	11.59
17-031-3103	Cook	Schiller Park					16.00	17.54	14.84	15.35	13.59*	12.91	12.64
17-031-3301	Cook	Summit	16.90	16.51	16.12	15.60	14.24	16.94	13.78	14.77	12.03	11.62	12.23
17-031-4007	Cook	Des Plaines		14.82	14.44	13.19	12.37	13.90	11.36	12.72	11.35	11.02	10.60
17-031-4201	Cook	Northbrook	14.28	14.70	13.17	12.14	11.15	14.52	11.84	13.21	10.09	9.33	9.34
17-031-6005	Cook	Cicero	16.54*	17.34	15.99	16.77*	15.19	16.25	14.34	14.79	13.25*	11.98*	11.90
17-043-4002	DuPage	Naperville	15.25	15.54	14.68	13.11	12.69	15.62	12.66	13.77	11.28	9.83	11.67
17-089-0003	Kane	Elgin	14.52*	15.04	14.24	13.33	11.44	15.64	11.80	13.19	10.79	9.61	11.29
17-089-0007	Kane	Aurora						15.89	12.61	14.53	10.34	10.01	11.44
17-097-1007	Lake	Zion	12.16	13.81*	13.44	11.26	10.25	13.78	10.69	11.86	9.34	8.83	9.66
17-111-0001	McHenry	Cary	14.74*	13.70	12.26	12.21	11.31	13.90	11.76	11.59	10.10	9.65	10.24
17-197-1002	Will	Joliet	16.02	16.06*	14.33	13.76	11.90*	15.43	12.17	14.58	11.66	10.52	11.83
17-197-1011	Will	Braidwood	14.21	12.92	13.47	11.88	10.26	13.21	9.82	12.07*	10.31	8.73	10.02

Source-Oriented Site NOT used for comparison to the Annual PM<sub>2.5</sub> Standard.

\*Indicates that the mean does not satisfy summary criteria

The Schiller Park monitor began operation in 2004.

The Des Plaines monitor began operation in 2001.

The Aurora monitor began operation in 2005.

Site ID	County	Site Name	Three Year Design Values								
			00-02	01-03	02-04	03-05	04-06	05-07	06-08	07-09	08-10
17-031-0022	Cook	Chicago - Washington High School	16.76	16.00	15.02	15.56	14.78	15.30	13.83	13.30	12.73
17-031-0050	Cook	Chicago - Southeast Police Station	16.68	16.31	14.86	15.23	14.55	14.68	13.09	12.31	11.75
17-031-0052	Cook	Chicago - Mayfair Pump Station	18.08	17.25	15.88	16.03	15.58	15.65	14.06	13.45	12.48
17-031-0057	Cook	Chicago - Springfield Pump Station	16.25	15.69	14.86	15.34	14.64	15.12	13.57	12.85	11.80
17-031-0076	Cook	Chicago - Com Ed Maintenance Building	15.93	15.34	14.88	15.18	14.73	14.78	13.22	12.44	11.80
17-031-1016	Cook	McCook	19.59	18.42	17.04	17.24	16.87	16.50	14.70	13.70	12.69
17-031-2001	Cook	Blue Island	16.37	15.73	14.74	15.14	14.56	14.63	13.33	12.83	11.92
17-031-3103	Cook	Schiller Park			16.00	16.77	16.13	15.91	14.59	13.95	13.05
17-031-3301	Cook	Summit	16.51	16.08	15.32	15.59	14.99	15.16	13.53	12.81	11.96
17-031-4007	Cook	Des Plaines	14.63	14.15	13.33	13.15	12.54	12.66	11.81	11.70	10.99
17-031-4201	Cook	Northbrook	14.05	13.34	12.15	12.60	12.50	13.19	11.71	10.88	9.59
17-031-6005	Cook	Cicero	16.62	16.70	15.98	16.07	15.26	15.13	14.12	13.34	12.38
17-043-4002	DuPage	Naperville	15.16	14.44	13.49	13.81	13.66	14.02	12.57	11.63	10.93
17-089-0003	Kane	Elgin	14.60	14.20	13.00	13.47	12.96	13.54	11.93	11.20	10.56
17-089-0007	Kane	Aurora				15.89	14.25	14.34	12.49	11.63	10.60
17-097-1007	Lake	Zion	13.14	12.84	11.65	11.76	11.57	12.11	10.63	10.01	9.28
17-111-0001	McHenry	Cary	13.56	12.72	11.93	12.47	12.32	12.42	11.15	10.45	10.00
17-197-1002	Will	Joliet	15.47	14.72	13.33	13.69	13.16	14.06	12.80	12.25	11.34
17-197-1011	Will	Braidwood	13.53	12.76	11.87	11.78	11.10	11.70	10.73	10.37	9.69
Value above the annual standard for fine particles.											

Source-Oriented Site NOT used for comparison to the annual standard for fine particles.

Red Text Indicates Incomplete Data

Blue Text Indicates Design Value Based on One Year of Data


Green Text Indicates Design Value Based on Two Years of Data

The Schiller Park monitor began operation in 2004.

The Des Plaines monitor began operation in 2001.


The Aurora monitor began operation in 2005.

# **Annual Fine Particle Trends for Illinois' Portion of the Chicago Area, 2006 through 2010**



Red line represents the annual standard for fine particles of  $15 \mu\text{g}/\text{m}^3$

# **Fine Particle Design Values for Illinois' Portion of the Chicago Area, 2006 through 2010**



Red line represents the annual standard for fine particles of  $15 \mu\text{g}/\text{m}^3$

**This page left intentionally blank.**

# **APPENDIX B**


**Nitrogen Oxides (NO<sub>x</sub>), Sulfur Dioxide (SO<sub>2</sub>), and  
Direct Fine Particle (PM<sub>2.5</sub>) Point Source  
Emissions (2002, 2005, and 2008), Chicago Area**




**This page left intentionally blank.**

Lake and Porter Counties, IN Point Source Totals (Tons per Year)			
Year	NO <sub>x</sub>	SO <sub>2</sub>	Direct PM <sub>2.5</sub>
2002	60,808.11	59,263.34	7,313.70
2005	41,948.57	48,139.53	6,451.40
2008	39,945.76	54,916.02	6,676.32


**NO<sub>x</sub> Point Source Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008**



**SO<sub>2</sub> Point Source Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008**




**Direct PM<sub>2.5</sub> Point Source Emission Trends, Lake and Porter Counties, Indiana,  
2002, 2005, and 2008**




**This page left intentionally blank.**

Entire Chicago Area Point Source Totals (Tons per Year)			
Year	NO <sub>x</sub>	SO <sub>2</sub>	Direct PM <sub>2.5</sub>
2002	114,858.11	180,861.34	10,070.70
2005	86,943.07	154,291.53	9,759.40
2008	75,884.76	145,622.02	10,535.32


### NO<sub>x</sub> Point Source Emission Trends, Entire Chicago Area, 2002, 2005, and 2008



### SO<sub>2</sub> Point Source Emission Trends, Entire Chicago Area, 2002, 2005 and 2008



## Direct PM<sub>2.5</sub> Point Source Emission Trends, Entire Chicago Area, 2002, 2005 and 2008



**This page left intentionally blank.**

# **APPENDIX C**

**Nitrogen Oxides (NO<sub>x</sub>), Sulfur Dioxide (SO<sub>2</sub>), and  
Direct Fine Particle (PM<sub>2.5</sub>) Emissions, All Sources  
(2002, 2005, and 2008), Chicago Area**



**This page left intentionally blank.**

<b>Lake and Porter Counties, IN Totals (Tons per Year)</b>			
<b>Year</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
2002	106,180.29	64,999.42	12,966.68
2005	66,426.65	49,876.77	7,152.31
2008	59,581.74	56,198.68	7,250.93

<b>2002-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,626.91	12,347.30	30,397.97	60,808.11	<b>106,180.29</b>
SO <sub>2</sub>	4,364.85	1,106.59	264.64	59,263.34	<b>64,999.42</b>
Direct PM <sub>2.5</sub>	4,404.91	685.43	562.64	7,313.70	<b>12,966.68</b>

<b>2002-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	2,010.98	3,318.89	2,322.39
LAKE	INDIANA	NONROAD	9,099.92	805.61	475.02
LAKE	INDIANA	POINT	35,255.39	35,957.52	5,025.35
PORTER	INDIANA	AREA	615.93	1,045.96	2,082.52
PORTER	INDIANA	NONROAD	3,247.38	300.98	210.41
PORTER	INDIANA	POINT	25,552.72	23,305.82	2,288.35
LAKE/PORTER	INDIANA	ONROAD	30,397.97	264.64	562.64


<b>2005-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,236.89	8,145.64	14,095.55	41,948.57	<b>66,426.65</b>
SO <sub>2</sub>	697.87	892.93	146.44	48,139.53	<b>49,876.77</b>
Direct PM <sub>2.5</sub>	23.65	447.87	229.39	6,451.40	<b>7,152.31</b>

<b>2005-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	1,687.72	525.81	18.48
LAKE	INDIANA	NONROAD	4,957.96	551.71	271.08
LAKE	INDIANA	POINT	22,394.78	25,354.10	5,086.98
PORTER	INDIANA	AREA	549.17	172.06	5.17
PORTER	INDIANA	NONROAD	3,187.68	341.22	176.79
PORTER	INDIANA	POINT	19,553.79	22,785.43	1,364.42
LAKE/PORTER	INDIANA	ONROAD	14,095.55	146.44	229.39


<b>2008-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,264.46	6,667.71	10,703.81	39,945.76	<b>59,581.74</b>
SO <sub>2</sub>	703.25	476.33	103.08	54,916.02	<b>56,198.68</b>
Direct PM <sub>2.5</sub>	23.60	363.91	187.10	6,676.32	<b>7,250.93</b>

<b>2008-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	1,707.89	529.04	18.43
LAKE	INDIANA	NONROAD	4,061.76	286.08	220.14
LAKE	INDIANA	POINT	23,273.37	28,744.32	5,249.09
PORTER	INDIANA	AREA	556.57	174.21	5.17
PORTER	INDIANA	NONROAD	2,605.95	190.25	143.77
PORTER	INDIANA	POINT	16,672.39	26,171.70	1,427.23
LAKE/PORTER	INDIANA	ONROAD	10,703.81	103.08	187.10


#### **NO<sub>x</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008**



### SO<sub>2</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008



### Direct PM<sub>2.5</sub> Emission Trends, Lake and Porter Counties, Indiana, 2002, 2005, and 2008



**This page left intentionally blank.**


Chicago Area Totals (Tons per Year)			
Year	NO <sub>x</sub>	SO <sub>2</sub>	Direct PM <sub>2.5</sub>
2002	447,601.29	197,480.42	32,069.68
2005	346,671.15	164,171.77	25,962.31
2008	278,649.74	152,367.68	25,767.93

2002-Chicago Area Totals (Tons per Year)					
	AREA	NONROAD	ONROAD	POINT	GRAND TOTAL
NO <sub>x</sub>	34,951.91	99,773.30	198,017.97	114,858.11	<b>447,601.29</b>
SO <sub>2</sub>	7,654.85	4,849.59	4,114.64	180,861.34	<b>197,480.42</b>
Direct PM <sub>2.5</sub>	12,845.91	5,519.43	3,633.64	10,070.70	<b>32,069.68</b>


2005-Chicago Area Totals (Tons per Year)					
	AREA	NONROAD	ONROAD	POINT	GRAND TOTAL
NO <sub>x</sub>	34,558.39	77,450.64	147,719.05	86,943.07	<b>346,671.15</b>
SO <sub>2</sub>	4,367.37	3,153.93	2,358.94	154,291.53	<b>164,171.77</b>
Direct PM <sub>2.5</sub>	8,838.65	4,691.37	2,672.89	9,759.40	<b>25,962.31</b>

2008-Chicago Area Totals (Tons per Year)					
	AREA	NONROAD	ONROAD	POINT	GRAND TOTAL
NO <sub>x</sub>	34,582.46	57,851.71	110,330.81	75,884.76	<b>278,649.74</b>
SO <sub>2</sub>	4,812.25	1,255.33	678.08	145,622.02	<b>152,367.68</b>
Direct PM <sub>2.5</sub>	9,212.60	4,016.91	2,003.10	10,535.32	<b>25,767.93</b>


#### NO<sub>x</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008



**SO<sub>2</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008**



**Direct PM<sub>2.5</sub> Emission Trends, Entire Chicago Area, 2002, 2005, and 2008**



# **APPENDIX D**

**Nitrogen Oxides (NO<sub>x</sub>) and Sulfur Dioxide (SO<sub>2</sub>)  
Emission Trends from Electric Generating Units  
(EGUs), Lake and Porter Counties, Indiana (1999  
to 2009)**




**This page left intentionally blank.**

### Indiana Statewide NO<sub>x</sub> Emissions from EGUs, 1999 to 2009

Year	Total NO <sub>x</sub> Emissions, tons/year
1999	347,216.50
2000	334,522.10
2001	315,419.70
2002	281,146.10
2003	260,980.00
2004	224,311.30
2005	207,981.60
2006	202,728.00
2007	196,553.10
2008	196,134.50
2009	110,968.90
Budget 2009-2014	108,935
Budget 2015 and later	90,779


### Indiana Statewide NO<sub>x</sub> Emissions from EGUs, 1999 to 2009



### NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana, 1999 to 2009

Year	Total NO <sub>x</sub> Emissions (tons/year)
1999	36,680.40
2000	33,554.90
2001	29,444.80
2002	23,645.40
2003	21,726.90
2004	22,289.60
2005	19,228.20
2006	17,688.90
2007	18,390.30
2008	18,916.30
2009	8,530.20

### NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana, 1999 to 2009



<b>NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 1999</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	25,168.20
Dean H. Mitchell	3,397.10
State Line	8,115.10
<b>Total</b>	<b>36,680.40</b>

<b>NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2000</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	20,824.40
Dean H. Mitchell	3,144.40
State Line	9,586.10
<b>Total</b>	<b>33,554.90</b>

<b>NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2001</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	18,611.50
Dean H. Mitchell	2,966.90
State Line	7,866.40
<b>Total</b>	<b>29,444.80</b>

<b>NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2002</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	16,380.00
Dean H. Mitchell	50.40
State Line	7,140.70
Whiting Clean Energy	74.30
<b>Total</b>	<b>23,645.40</b>

<b>NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2003</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	13,332.90
State Line	8,369.70
Whiting Clean Energy	24.30
<b>Total</b>	<b>21,726.90</b>

<b>NOx Emissions from EGUs, Lake and Porter Counties, Indiana 2004</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	13,900.70
State Line	8,347.90
Whiting Clean Energy	41.00
<b>Total</b>	<b>22,289.60</b>

<b>NOx Emissions from EGUs, Lake and Porter Counties, Indiana 2005</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	12,329.30
State Line	6,847.90
Whiting Clean Energy	51.00
<b>Total</b>	<b>19,228.20</b>

<b>NOx Emissions from EGUs, Lake and Porter Counties, Indiana 2006</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	10,354.00
State Line	7,288.10
Whiting Clean Energy	46.80
<b>Total</b>	<b>17,688.90</b>

<b>NOx Emissions from EGUs, Lake and Porter Counties, Indiana 2007</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	10,118.30
State Line	8,205.40
Whiting Clean Energy	66.60
<b>Total</b>	<b>18,390.30</b>

<b>NOx Emissions from EGUs, Lake and Porter Counties, Indiana 2008</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	9,741.60
State Line	9,101.60
Whiting Clean Energy	73.10
<b>Total</b>	<b>18,916.30</b>


<b>NO<sub>x</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2009</b>	
<b>Facility</b>	<b>NO<sub>x</sub> Emissions (tons/year)</b>
Bailly	2,459.30
State Line	5,990.90
Whiting Clean Energy	80.00
<b>Total</b>	<b>8,530.20</b>

**This page left intentionally blank.**

### Indiana Statewide SO<sub>2</sub> Emissions from EGUs, 1999 to 2009

Year	Total SO <sub>2</sub> Emissions (tons/year)
1999	941,852.40
2000	874,617.20
2001	795,505.60
2002	778,868.00
2003	804,828.60
2004	862,876.40
2005	870,811.80
2006	820,993.40
2007	714,529.20
2008	593,154.00
2009	413,726.40
Budget 2009-2014	254,599.0
Budget 2015 and later	178,219.0

### Indiana Statewide SO<sub>2</sub> Emissions from EGUs, 1999 to 2009






### SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana, 1999 to 2009

Year	Total SO <sub>2</sub> Emissions (tons/year)
1999	19,728.30
2000	20,502.80
2001	20,733.50
2002	13,789.50
2003	12,556.20
2004	13,312.20
2005	12,672.40
2006	10,659.60
2007	13,651.20
2008	20,331.00
2009	14,684.70

### SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana, 1999 to 2009



<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 1999</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	3,813.30
Dean H. Mitchell	8,830.00
State Line	7,085.00
<b>Total</b>	<b>19,728.30</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2000</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	5,144.80
Dean H. Mitchell	6,364.00
State Line	8,994.00
<b>Total</b>	<b>20,502.80</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2001</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	6,076.20
Dean H. Mitchell	6,797.20
State Line	7,860.10
<b>Total</b>	<b>20,733.50</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2002</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	5,217.60
Dean H. Mitchell	127.30
State Line	8,442.90
Whiting Clean Energy	1.70
<b>Total</b>	<b>13,789.50</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2003</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	4,544.00
State Line	8,011.00
Whiting Clean Energy	1.20
<b>Total</b>	<b>12,556.20</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2004</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	3,608.70
State Line	9,701.50
Whiting Clean Energy	2.00
<b>Total</b>	<b>13,312.20</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2005</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	4,720.20
State Line	7,949.40
Whiting Clean Energy	2.80
<b>Total</b>	<b>12,672.40</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2006</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	3,308.70
State Line	7,348.00
Whiting Clean Energy	2.90
<b>Total</b>	<b>10,659.60</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2007</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	4,313.60
State Line	9,333.20
Whiting Clean Energy	4.40
<b>Total</b>	<b>13,651.20</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2008</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	8,700.90
State Line	11,625.00
Whiting Clean Energy	5.10
<b>Total</b>	<b>20,331.00</b>

<b>SO<sub>2</sub> Emissions from EGUs, Lake and Porter Counties, Indiana 2009</b>	
<b>Facility</b>	<b>SO<sub>2</sub> Emissions, tons / year</b>
Bailly	4,903.40
State Line	9,776.00
Whiting Clean Energy	5.30
<b>Total</b>	<b>14,684.70</b>

**This page left intentionally blank.**

# **APPENDIX E**

**2005 and 2008 Base Year Emission Inventories  
and 2015, 2020, and 2025 Projected Emission  
Inventories for Nitrogen Oxides (NO<sub>x</sub>), Sulfur  
Dioxide (SO<sub>2</sub>), and Direct Fine Particles (PM<sub>2.5</sub>),  
Chicago Area**

**This page left intentionally blank.**

<b>Lake and Porter Counties, IN Totals (Tons per Year)</b>			
<b>Year</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
2002	106,180.29	64,999.42	12,966.68
2005	66,426.65	49,876.77	7,152.31
2008	59,581.74	56,198.68	7,250.93
2015	41,795.31	56,235.30	7,057.65
2020	37,082.26	39,445.89	6,888.26
2025	35,013.79	36,787.24	6,762.10

<b>2002-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,626.91	12,347.30	30,397.97	60,808.11	<b>106,180.29</b>
SO <sub>2</sub>	4,364.85	1,106.59	264.64	59,263.34	<b>64,999.42</b>
Direct PM <sub>2.5</sub>	4,404.91	685.43	562.64	7,313.70	<b>12,966.68</b>

<b>2002-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	2,010.98	3,318.89	2,322.39
LAKE	INDIANA	NONROAD	9,099.92	805.61	475.02
LAKE	INDIANA	POINT	35,255.39	35,957.52	5,025.35
PORTER	INDIANA	AREA	615.93	1,045.96	2,082.52
PORTER	INDIANA	NONROAD	3,247.38	300.98	210.41
PORTER	INDIANA	POINT	25,552.72	23,305.82	2,288.35
LAKE/PORTER	INDIANA	ONROAD	30,397.97	264.64	562.64

<b>2005-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,236.89	8,145.64	14,095.55	41,948.57	<b>66,426.65</b>
SO <sub>2</sub>	697.87	892.93	146.44	48,139.53	<b>49,876.77</b>
Direct PM <sub>2.5</sub>	23.65	447.87	229.39	6,451.40	<b>7,152.31</b>

<b>2005-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	1,687.72	525.81	18.48
LAKE	INDIANA	NONROAD	4,957.96	551.71	271.08
LAKE	INDIANA	POINT	22,394.78	25,354.10	5,086.98
PORTER	INDIANA	AREA	549.17	172.06	5.17
PORTER	INDIANA	NONROAD	3,187.68	341.22	176.79
PORTER	INDIANA	POINT	19,553.79	22,785.43	1,364.42
LAKE/PORTER	INDIANA	ONROAD	14,095.55	146.44	229.39



<b>2008-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,264.46	6,667.71	10,703.81	39,945.76	<b>59,581.74</b>
SO <sub>2</sub>	703.25	476.33	103.08	54,916.02	<b>56,198.68</b>
Direct PM <sub>2.5</sub>	23.60	363.91	187.10	6,676.32	<b>7,250.93</b>

<b>2008-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	1,707.89	529.04	18.43
LAKE	INDIANA	NONROAD	4,061.76	286.08	220.14
LAKE	INDIANA	POINT	23,273.37	28,744.32	5,249.09
PORTER	INDIANA	AREA	556.57	174.21	5.17
PORTER	INDIANA	NONROAD	2,605.95	190.25	143.77
PORTER	INDIANA	POINT	16,672.39	26,171.70	1,427.23
LAKE/PORTER	INDIANA	ONROAD	10,703.81	103.08	187.10

<b>2015-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,226.21	4,962.17	5,723.67	28,883.26	<b>41,795.31</b>
SO <sub>2</sub>	682.86	267.22	66.23	42,394.24	<b>43,410.55</b>
Direct PM <sub>2.5</sub>	22.70	248.01	136.61	6,650.33	<b>7,057.65</b>

<b>2015-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	1,678.82	513.12	17.70
LAKE	INDIANA	NONROAD	2,958.96	151.94	148.95
LAKE	INDIANA	POINT	20,585.05	25,544.26	5,214.61
PORTER	INDIANA	AREA	547.39	169.74	5.00
PORTER	INDIANA	NONROAD	2,003.21	115.28	99.06
PORTER	INDIANA	POINT	8,298.21	16,849.98	1,435.72
LAKE/PORTER	INDIANA	ONROAD	5,723.67	66.23	136.61

<b>2020-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,187.09	4,057.84	3,004.68	27,832.65	<b>37,082.26</b>
SO <sub>2</sub>	664.67	215.27	72.76	38,493.19	<b>39,445.89</b>
Direct PM <sub>2.5</sub>	21.97	185.11	114.32	6,566.86	<b>6,888.26</b>

<b>2020-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	1,649.30	499.19	17.12
LAKE	INDIANA	NONROAD	2,365.01	118.63	110.29
LAKE	INDIANA	POINT	20,498.81	25,099.75	5,141.47
PORTER	INDIANA	AREA	537.79	165.48	4.85
PORTER	INDIANA	NONROAD	1,692.83	96.64	74.82
PORTER	INDIANA	POINT	7,333.84	10,788.52	1,425.39
LAKE/PORTER	INDIANA	ONROAD	3,004.68	72.76	114.32


<b>2025-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	2,148.80	3,349.95	2,534.95	26,980.09	<b>35,013.79</b>
SO <sub>2</sub>	647.07	175.39	76.51	35,888.27	<b>36,787.24</b>
Direct PM <sub>2.5</sub>	21.29	140.67	115.39	6,484.75	<b>6,762.10</b>

<b>2025-Lake and Porter Counties, IN Totals (Tons per Year)</b>					
<b>COUNTY</b>	<b>STATE</b>	<b>SECTOR</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
LAKE	INDIANA	AREA	1,620.41	485.72	16.57
LAKE	INDIANA	NONROAD	1,910.65	93.83	83.22
LAKE	INDIANA	POINT	20,413.24	25,099.75	5,069.59
PORTER	INDIANA	AREA	528.39	161.35	4.72
PORTER	INDIANA	NONROAD	1,439.30	81.56	57.45
PORTER	INDIANA	POINT	6,566.85	10,788.52	1,415.16
LAKE/PORTER	INDIANA	ONROAD	2,534.95	76.51	115.39


**Comparison of 2008 Estimated and 2025 Projected Emission Estimates,  
Lake and Porter Counties, Indiana (Tons per Year)**

	<b>2008</b>	<b>2025</b>	<b>Change</b>	<b>% Change</b>
<b>NO<sub>x</sub></b>	59,581.74	35,013.79	-24,567.95	41.23% decrease
<b>SO<sub>2</sub></b>	56,198.68	36,787.24	-19,411.44	35.54% decrease
<b>Direct PM<sub>2.5</sub></b>	7,250.93	6,762.10	-488.83	6.74% decrease


**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected  
NO<sub>x</sub> Emissions, Lake and Porter Counties, Indiana**




**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected  
SO<sub>2</sub> Emissions, Lake and Porter Counties, Indiana**



**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected Direct PM<sub>2.5</sub> Emissions, Lake and Porter Counties, Indiana**



**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO<sub>x</sub>, SO<sub>2</sub>, and Direct PM<sub>2.5</sub> Emissions, Lake and Porter Counties, Indiana**



**This page left intentionally blank.**

<b>Chicago Area Totals (Tons per Year)</b>			
<b>Year</b>	<b>NO<sub>x</sub></b>	<b>SO<sub>2</sub></b>	<b>Direct PM<sub>2.5</sub></b>
2002	447,601.29	197,480.42	32,069.68
2005	346,671.15	164,171.77	25,962.31
2008	278,649.74	152,367.68	25,767.93
2015	187,557.31	107,285.55	25,128.65
2020	156,231.26	98,829.89	24,729.26
2025	149,198.79	99,453.24	25,074.10

<b>2002-Chicago Area Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	34,951.91	99,773.30	198,017.97	114,858.11	<b>447,601.29</b>
SO <sub>2</sub>	7,654.85	4,849.59	4,114.64	180,861.34	<b>197,480.42</b>
Direct PM <sub>2.5</sub>	12,845.91	5,519.43	3,633.64	10,070.70	<b>32,069.68</b>

<b>2005-Chicago Area Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	34,558.39	77,450.64	147,719.05	86,943.07	<b>346,671.15</b>
SO <sub>2</sub>	4,367.37	3,153.93	2,358.94	154,291.53	<b>164,171.77</b>
Direct PM <sub>2.5</sub>	8,838.65	4,691.37	2,672.89	9,759.40	<b>25,962.31</b>

<b>2008-Chicago Area Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	34,582.46	57,851.71	110,330.81	75,884.76	<b>278,649.74</b>
SO <sub>2</sub>	4,812.25	1,255.33	678.08	145,622.02	<b>152,367.68</b>
Direct PM <sub>2.5</sub>	9,212.60	4,016.91	2,003.10	10,535.32	<b>25,767.93</b>

<b>2015-Chicago Area Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	35,223.21	40,899.17	55,469.67	55,965.26	<b>187,557.31</b>
SO <sub>2</sub>	4,948.86	1,133.22	717.23	100,486.24	<b>107,285.55</b>
Direct PM <sub>2.5</sub>	9,698.70	3,243.01	1,367.61	10,819.33	<b>25,128.65</b>


<b>2020-Chicago Area Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	35,464.09	32,328.84	32,106.68	56,331.65	<b>156,231.26</b>
SO <sub>2</sub>	4,996.67	1,134.27	753.76	91,945.19	<b>98,829.89</b>
Direct PM <sub>2.5</sub>	10,030.97	2,583.11	1,157.32	10,957.86	<b>24,729.26</b>

<b>2025-Chicago Area Totals (Tons per Year)</b>					
	<b>AREA</b>	<b>NONROAD</b>	<b>ONROAD</b>	<b>POINT</b>	<b>GRAND TOTAL</b>
NO <sub>x</sub>	35,835.80	30,522.95	26,221.95	56,618.09	<b>149,198.79</b>
SO <sub>2</sub>	5,054.07	1,390.39	810.51	92,198.27	<b>99,453.24</b>
Direct PM <sub>2.5</sub>	10,398.29	2,407.67	1,179.39	11,088.75	<b>25,074.10</b>


**Comparison of 2008 Estimated and 2025 Projected NO<sub>x</sub>, SO<sub>2</sub>, and Direct PM<sub>2.5</sub> Emissions, Entire Chicago Area (Tons per Year)**

	<b>2008</b>	<b>2025</b>	<b>Change</b>	<b>% Change</b>
<b>NO<sub>x</sub></b>	278,649.74	149,198.79	-129,450.95	46.46% decrease
<b>SO<sub>2</sub></b>	152,367.68	99,453.24	-52,914.44	34.73% decrease
<b>Direct PM<sub>2.5</sub></b>	25,767.93	25,074.10	-693.83	2.69% decrease


**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO<sub>x</sub> Emissions, Entire Chicago Area**



**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected SO<sub>2</sub> Emissions, Entire Chicago Area**




**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected Direct PM<sub>2.5</sub> Emissions, Entire Chicago Area**





**Comparison of 2005 and 2008 Estimated and 2015, 2020, and 2025 Projected NO<sub>x</sub>, SO<sub>2</sub>, and Direct PM<sub>2.5</sub> Emissions, Entire Chicago Area**



# **APPENDIX F**

## **Example Mobile Source Input and Output Calculation Files for Lake and Porter Counties, Indiana**

**This page left intentionally blank**

MOBILE6 INPUT FILE

POLLUTANTS : HC NOX  
DATABASE OUTPUT :  
WITH FIELDNAMES :  
DATABASE EMISSIONS : 2222 2221  
DATABASE FACILITIES: ARTERIAL FREEWAY LOCAL RAMP

RUN DATA

MIN/MAX TEMPERATURE: 60. 82.  
FUEL RVP : 9.0  
FUEL PROGRAM : 2 N  
NO REFUELING :  
EXPAND EXHAUST :  
EXPAND EVAPORATIVE :  
ANTI-TAMP PROG :  
90 76 95 22222 21111111 1 12 095. 12111112  
REG DIST : iregdata.d

\* The following describes the I/M programs within Lake/Porter Counties:

\* First I/M Program

I/M PROGRAM : 1 1997 2050 2 T/O IDLE  
I/M MODEL YEARS : 1 1976 1980  
I/M VEHICLES : 1 22222 21111111 1  
I/M STRINGENCY : 1 20.0  
I/M COMPLIANCE : 1 95.0  
I/M WAIVER RATES : 1 3.0 3.0

\* Second I/M Program (Cutpoints for LDGV, LDGT2, LDGT4 and HDGV2B)

I/M PROGRAM : 2 1997 2050 2 T/O IM240  
I/M MODEL YEARS : 2 1981 1995  
I/M VEHICLES : 2 21212 21111111 1  
I/M STRINGENCY : 2 20.0  
I/M COMPLIANCE : 2 95.0  
I/M WAIVER RATES : 2 3.0 3.0  
I/M CUTPOINTS : 2 IM2002A.d  
I/M GRACE PERIOD : 2 4

\* Third I/M Program (Cutpoints for LDGT1 and LDGT3)

I/M PROGRAM : 3 1997 2050 2 T/O IM240  
I/M MODEL YEARS : 3 1981 1995  
I/M VEHICLES : 3 12121 11111111 1  
I/M STRINGENCY : 3 20.0  
I/M COMPLIANCE : 3 95.0  
I/M WAIVER RATES : 3 3.0 3.0  
I/M CUTPOINTS : 3 IM2002B.d  
I/M GRACE PERIOD : 3 4

\* Fourth I/M Program

I/M PROGRAM : 4 1997 2050 2 T/O GC  
I/M MODEL YEARS : 4 1976 1995  
I/M VEHICLES : 4 22222 21111111 1

\* Fifth I/M Program

I/M PROGRAM : 5 2002 2050 2 T/O OBD I/M

I/M MODEL YEARS : 5 1996 2050  
 I/M VEHICLES : 5 22222 21111111 1  
 I/M STRINGENCY : 5 20.0  
 I/M COMPLIANCE : 5 95.0  
 I/M WAIVER RATES : 5 3.0 3.0  
 I/M GRACE PERIOD : 5 4  
 \* Sixth I/M Program  
 I/M PROGRAM : 6 1997 2050 2 T/O EVAP OBD & GC  
 I/M MODEL YEARS : 6 1996 2050  
 I/M VEHICLES : 6 22222 11111111 1

# SCENARIO RECORD

CALENDAR YEAR : 2010  
 EVALUATION MONTH : 7  
 VMT FRACTIONS :  
 0.364243579 0.091574782 0.304735387 0.093938666 0.035952411  
 0.033384381 0.003252837 0.002739231  
 0.002054423 0.007447285 0.008816901 0.009587310 0.034154790  
 0.001712020 0.000856010 0.005549987  
 VMT BY FACILITY : 2010nvmt.d  
 SPEED VMT : svmt10.d

END OF RUN

MOBILE6 NITROGEN OXIDES (NO<sub>x</sub>) OUTPUT FILE FOR LAKE AND PORTER COUNTIES

January										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.732	0.891	1.346	1.002	2.898	0.498	0.719	8.845	2.28	1.544
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.141	0.172	0.23	0.186		0.019	0.02		0.624	0.152305
NOx Running:	0.591	0.72	1.116	0.816		0.479	0.7		1.659	1.391394
NOx Total Exhaust:	0.732	0.891	1.346	1.002	2.898	0.498	0.719	8.845	2.28	1.544

February										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.713	0.865	1.306	0.973	2.872	0.498	0.719	8.845	2.22	1.52
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.137	0.167	0.223	0.18		0.019	0.02		0.609	0.147622
NOx Running:	0.575	0.699	1.083	0.792		0.479	0.7		1.613	1.37194
NOx Total Exhaust:	0.713	0.865	1.306	0.973	2.872	0.498	0.719	8.845	2.22	1.52

March										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.664	0.801	1.209	0.901	2.81	0.498	0.719	8.845	2.07	1.462
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.128	0.154	0.207	0.167		0.019	0.02		0.572	0.137327
NOx Running:	0.536	0.647	1.002	0.733		0.479	0.7		1.497	1.324279
NOx Total Exhaust:	0.664	0.801	1.209	0.901	2.81	0.498	0.719	8.845	2.07	1.462

April										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.595	0.716	1.07	0.802	2.566	0.48	0.69	8.217	1.87	1.331
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.116	0.14	0.187	0.151		0.018	0.02		0.518	0.124274
NOx Running:	0.479	0.576	0.884	0.651		0.462	0.67		1.354	1.206133
NOx Total Exhaust:	0.595	0.716	1.07	0.802	2.566	0.48	0.69	8.217	1.87	1.331

May										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.528	0.632	0.945	0.709	2.518	0.48	0.69	8.217	1.66	1.254
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.103	0.123	0.164	0.133		0.018	0.02		0.463	0.109804
NOx Running:	0.425	0.509	0.781	0.575		0.462	0.67		1.195	1.144293
NOx Total Exhaust:	0.528	0.632	0.945	0.709	2.518	0.48	0.69	8.217	1.66	1.254

June										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.484	0.569	0.85	0.637	2.486	0.48	0.69	8.217	1.44	1.199
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.091	0.109	0.145	0.117		0.018	0.02		0.406	0.096734
NOx Running:	0.394	0.46	0.705	0.52		0.462	0.67		1.036	1.102322
NOx Total Exhaust:	0.484	0.569	0.85	0.637	2.486	0.48	0.69	8.217	1.44	1.199



July										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.465	0.54	0.807	0.605	2.479	0.48	0.69	8.217	1.33	1.174
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.085	0.102	0.136	0.11		0.018	0.02		0.375	0.09071
NOx Running:	0.38	0.438	0.672	0.495		0.462	0.67		0.952	1.083443
NOx Total Exhaust:	0.465	0.54	0.807	0.605	2.479	0.48	0.69	8.217	1.33	1.174

August										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.47	0.55	0.822	0.616	2.481	0.48	0.69	8.217	1.38	1.182
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.087	0.104	0.139	0.113		0.018	0.02		0.389	0.093088
NOx Running:	0.383	0.445	0.683	0.503		0.462	0.67		0.989	1.088994
NOx Total Exhaust:	0.47	0.55	0.822	0.616	2.481	0.48	0.69	8.217	1.38	1.182

September										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.502	0.596	0.891	0.668	2.495	0.48	0.69	8.217	1.55	1.222
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.096	0.115	0.154	0.125		0.018	0.02		0.436	0.102913
NOx Running:	0.405	0.48	0.737	0.543		0.462	0.67		1.118	1.11911
NOx Total Exhaust:	0.502	0.596	0.891	0.668	2.495	0.48	0.69	8.217	1.55	1.222

October										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.517	0.606	0.934	0.686	2.393	0.424	0.61	7.606	1.79	1.191
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.1	0.118	0.162	0.129		0.016	0.017		0.501	0.106817
NOx Running:	0.418	0.488	0.772	0.558		0.408	0.593		1.293	1.084536
NOx Total Exhaust:	0.517	0.606	0.934	0.686	2.393	0.424	0.61	7.606	1.79	1.191

November										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.58	0.683	1.053	0.774	2.468	0.424	0.61	7.606	2.02	1.263
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.112	0.133	0.183	0.145		0.016	0.017		0.559	0.119892
NOx Running:	0.468	0.55	0.87	0.628		0.408	0.593		1.46	1.142646
NOx Total Exhaust:	0.58	0.683	1.053	0.774	2.468	0.424	0.61	7.606	2.02	1.263

December										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128	0.5243	0.0306	0.0003	0.0019	0.0734	0.0055	1
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Composite NOX :	0.634	0.751	1.158	0.85	2.531	0.424	0.61	7.606	2.21	1.326
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Exhaust emissions (g/mi ):										
NOx Start:	0.122	0.146	0.201	0.16		0.016	0.017		0.607	0.131659
NOx Running:	0.511	0.604	0.956	0.69		0.408	0.593		1.602	1.193509
NOx Total Exhaust:	0.634	0.751	1.158	0.85	2.531	0.424	0.61	7.606	2.21	1.326

MOBILE6 FINE PARTICULATE MATTER (PM<sub>2.5</sub>) OUTPUT FILE FOR LAKE AND PORTER COUNTIES

January										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Factors (g/mi):										
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0038	0.0037	0.0038	0.0037	0.0347	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0401	0.0178	0.1078	-----	0.008
OCARBON:	-----	-----	-----	-----	-----	0.0113	0.0257	0.0547	-----	0.0041
SO4:	0.0002	0.0004	0.0004	0.0004	0.0016	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0364	0.0516	0.0438	0.1634	0.0143	0.0169
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0115	0.0439	0.059	0.0511	0.1752	0.0206	0.0246
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

February										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Factors (g/mi):										
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0038	0.0037	0.0038	0.0037	0.0347	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0401	0.0178	0.1078	-----	0.008

OCARBON:	-----	-----	-----	-----	-----	0.0113	0.0257	0.0547	-----	0.0041
SO4:	0.0002	0.0004	0.0004	0.0004	0.0016	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0364	0.0516	0.0438	0.1634	0.0143	0.0169
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0115	0.0439	0.059	0.0511	0.1752	0.0206	0.0246
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

March										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0038	0.0037	0.0038	0.0037	0.0347	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0401	0.0178	0.1078	-----	0.008
OCARBON:	-----	-----	-----	-----	-----	0.0113	0.0257	0.0547	-----	0.0041
SO4:	0.0002	0.0004	0.0004	0.0004	0.0016	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0364	0.0516	0.0438	0.1634	0.0143	0.0169
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0115	0.0439	0.059	0.0511	0.1752	0.0206	0.0246
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

April										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0037	0.0037	0.0326	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0385	0.0169	0.1006	-----	0.0074
OCARBON:	-----	-----	-----	-----	-----	0.0109	0.0243	0.0511	-----	0.0038
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0343	0.0496	0.0415	0.1526	0.0143	0.016
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0114	0.0418	0.0569	0.0488	0.1644	0.0206	0.0237
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

May										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0037	0.0037	0.0326	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0385	0.0169	0.1006	-----	0.0074
OCARBON:	-----	-----	-----	-----	-----	0.0109	0.0243	0.0511	-----	0.0038
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004

Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0343	0.0496	0.0415	0.1526	0.0143	0.016
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0114	0.0418	0.0569	0.0488	0.1644	0.0206	0.0237
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

June										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Factors (g/mi):										
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0037	0.0037	0.0326	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0385	0.0169	0.1006	-----	0.0074
OCARBON:	-----	-----	-----	-----	-----	0.0109	0.0243	0.0511	-----	0.0038
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0343	0.0496	0.0415	0.1526	0.0143	0.016
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0114	0.0418	0.0569	0.0488	0.1644	0.0206	0.0237
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

July										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0037	0.0037	0.0326	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0385	0.0169	0.1006	-----	0.0074
OCARBON:	-----	-----	-----	-----	-----	0.0109	0.0243	0.0511	-----	0.0038
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0343	0.0496	0.0415	0.1526	0.0143	0.016
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0114	0.0418	0.0569	0.0488	0.1644	0.0206	0.0237
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

August										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0037	0.0037	0.0326	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0385	0.0169	0.1006	-----	0.0074
OCARBON:	-----	-----	-----	-----	-----	0.0109	0.0243	0.0511	-----	0.0038
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004



Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0343	0.0496	0.0415	0.1526	0.0143	0.016
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0114	0.0418	0.0569	0.0488	0.1644	0.0206	0.0237
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

September										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Factors (g/mi):										
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0037	0.0037	0.0326	-----	-----	-----	0.0142	0.0044
ECARBON:	-----	-----	-----	-----	-----	0.0385	0.0169	0.1006	-----	0.0074
OCARBON:	-----	-----	-----	-----	-----	0.0109	0.0243	0.0511	-----	0.0038
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.0041	0.0042	0.0041	0.0343	0.0496	0.0415	0.1526	0.0143	0.016
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0115	0.0114	0.0418	0.0569	0.0488	0.1644	0.0206	0.0237
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.0029	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1016	0.1013	0.1016	0.0451	0.0068	0.0068	0.027	0.0113	0.0937

October										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0036	0.0036	0.029	-----	-----	-----	0.0142	0.0042
ECARBON:	-----	-----	-----	-----	-----	0.0334	0.0146	0.0936	-----	0.0069
OCARBON:	-----	-----	-----	-----	-----	0.0094	0.021	0.0475	-----	0.0035
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.004	0.0041	0.0041	0.0307	0.043	0.036	0.1421	0.0143	0.0151
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0114	0.0114	0.0382	0.0503	0.0433	0.1539	0.0206	0.0228
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.003	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1017	0.1017	0.1017	0.0451	0.0068	0.0068	0.027	0.0113	0.0938

## November

Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Fac	tors (g/mi ):									
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0036	0.0036	0.029	-----	-----	-----	0.0142	0.0042
ECARBON:	-----	-----	-----	-----	-----	0.0334	0.0146	0.0936	-----	0.0069
OCARBON:	-----	-----	-----	-----	-----	0.0094	0.021	0.0475	-----	0.0035
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004

Total Exhaust PM:	0.004	0.004	0.0041	0.0041	0.0307	0.043	0.036	0.1421	0.0143	0.0151
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0114	0.0114	0.0382	0.0503	0.0433	0.1539	0.0206	0.0228
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.003	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1017	0.1017	0.1017	0.0451	0.0068	0.0068	0.027	0.0113	0.0938

December										
Vehicle Type:	LDGV	LDGT12	LDGT34	LDGT	HDGV	LDDV	LDDT	HDDV	MC	All Veh
GVWR:		<6000	>6000	(All)						
	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
VMT Distribution:	0.3639	0.3963	0.128		0.0306	0.0003	0.0019	0.0734	0.0055	1
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----
Composite Emission Factors (g/mi):										
Lead:	0	0	0	0	0	-----	-----	-----	0	0
GASPM:	0.0037	0.0036	0.0036	0.0036	0.029	-----	-----	-----	0.0142	0.0042
ECARBON:	-----	-----	-----	-----	-----	0.0334	0.0146	0.0936	-----	0.0069
OCARBON:	-----	-----	-----	-----	-----	0.0094	0.021	0.0475	-----	0.0035
SO4:	0.0002	0.0004	0.0004	0.0004	0.0017	0.0002	0.0003	0.0009	0.0001	0.0004
Total Exhaust PM:	0.004	0.004	0.0041	0.0041	0.0307	0.043	0.036	0.1421	0.0143	0.0151
Brake:	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053	0.0053
Tire:	0.002	0.002	0.002	0.002	0.0022	0.002	0.002	0.0065	0.001	0.0023
Total PM:	0.0113	0.0114	0.0114	0.0114	0.0382	0.0503	0.0433	0.1539	0.0206	0.0228
SO2:	0.0068	0.0088	0.0115	0.0095	0.0166	0.003	0.0056	0.0132	0.0033	0.0089
NH3:	0.1017	0.1017	0.1017	0.1017	0.0451	0.0068	0.0068	0.027	0.0113	0.0938

MONTHLY VEHICLE MILES TRAVELLED FOR LAKE AND PORTER COUNTIES

month	iimvmt	iemvmt	eimvmt	eemvmt	MVMT
1	257856354	53480480	49596650	119671310	480,604,795
2	238912901	49551529	45953025	110879640	445,297,096
3	271719599	56355775	52263137	126105251	506,443,763
4	269394152	55873468	51815855	125026010	502,109,484
5	281146606	58310975	54076348	130480331	524,014,260
6	281736912	58433407	54189888	130754293	525,114,500
7	285582844	59231070	54929623	132539193	532,282,730
8	286691904	59461093	55142942	133053908	534,349,847
9	271004077	56207373	52125512	125773177	505,110,138
10	277264897	57505893	53329732	128678828	516,779,350
11	266710943	55316959	51299761	123780731	497,108,394
12	277264897	57505893	53329732	128678828	516,779,350

**This page left intentionally blank.**

# **APPENDIX G**

## **Indiana Department of Environmental Management (IDEM) – Area Source Inventory Standard Operating Procedure**

**This page left intentionally blank.**



**Area Source Inventory**  
S-006-OAQ-R-MO-08-S-R1  
**Standard Operating Procedure**

**Office:** Office of Air Quality  
**Branch:** Air Programs Branch  
**Section:** Technical Support and Modeling Section

**Revised:** 02/27/2008 **Revision Cycle:** 2 years  
**Effective date:** 02/15/07

**Scope of operations**

This SOP is to identify source categories and develop emissions not calculated in point source inventories. This data is compiled every three years as mandated by EPA.

**Scope of applicability**

This SOP is for the Senior Environmental Manager and the Environmental Manager in the Emissions Group.

**Authorized Signatures**

I approve and authorize this Standard Operating Procedure:

**Branch Chief**

Scott Deloney  
Typed/Printed

  
Signature

3/12/08  
Date

**Section Chief**


Ken Ritter  
Typed/Printed

  
Signature

3/10/08  
Date

**Section QA Contact**

Michele Boner  
Typed/Printed

  
Signature

3/10/08  
Date

**Branch QA Coordinator**

Chris Pedersen  
Typed/Printed

  
Signature

3-10-08  
Date

**Author**

Michele Boner  
Typed/Printed

  
Signature

3/10/08  
Date

This Standard Operating Procedure is consistent with agency requirements.

  
Indiana Department of Environmental Management  
Quality Assurance Program  
Planning and Assessment

3-17-08  
Date



## Table of Contents

Scope of operations.....	1
Scope of applicability .....	1
Authorized Signatures .....	1
1. Overview work flow chart .....	3
2. Definitions.....	3
3. Roles.....	4
Responsibilities: .....	4
4. Description of equipment, forms, and/or software to be used.....	4
5. Procedure .....	4
5.1     Procedural Flowchart .....	4
5.2     Procedure .....	5
6. Standards and checklists.....	24
7. Records Management.....	24
8. Quality Assurance / Quality Control .....	25
9. Continuous Improvement Cycle .....	25
10. References .....	25
11. History of Revisions .....	25
12. Appendices.....	25

## 1. Overview work flow chart

The process described is not part of a larger system and does not need an Overview work flow chart.

## 2. Definitions

**AP-42** – Compilation of Air Pollutant Emission Factors AP-42, Fifth Edition, Volume I: Stationary Point and Area Sources (January 1995) plus Supplements A – F (Updates 2001 – 2004). AP-42 can be obtained at [www.epa.gov/ttn/chief/ap42/](http://www.epa.gov/ttn/chief/ap42/).

**Area Sources** - A collection of similar emission units within a geographic area that collectively represent individual sources that are small and numerous and have not been inventoried as a specific point, mobile, or biogenic source.

**Authorized** - Established by official authority and usage; as with a policy, standard operating procedure (SOP), or quality assurance project plan (QAPP) that is signed and dated.

**EIIP (Emission Inventory Improvement Program)** -The EIIP is an EPA program established in 1993 to promote the development and use of standard procedures for collecting, calculating, storing, reporting, and sharing air emissions data.

**Emission Factors** - An emission factor is the estimate of the quantity of pollutant released to the atmosphere (because of some operation or activity such as combustion or industrial production) divided by the level of that activity.

**Process** - The term “process” used when describing area sources is used to name an operation or activity that produces emissions.

**NEI** - National Emission Inventory Air Pollutant Emission Trends, U.S. EPA.

**Standard Industrial Classification (SIC) Code** - A Standard Industrial Classification code from the series of codes devised by the United States Office of Management and Budget (OMB) to classify establishments according to the type of economic activity in which they engage.

**Source Classification Code (SCC)** - Source Classification Code is a process-level code that describes the equipment or operation emitting pollutants.

### 3. Roles

Title	# of Staff	Experience	Qualifications	Location
Senior Environmental Manager	1	N/A	MS ACCESS, Emission Inventories and familiarity with the EIIP	Air Programs Branch
Environmental Manager	1	N/A	MS ACCESS, Emission Inventories and familiarity with the EIIP	Air Programs Branch

#### Responsibilities:

##### Senior Environmental Manager

Oversees work of the Environmental Manager and ensures that all goals are met. The Senior Environmental Manager also does the final upload to the NEI.

##### Environmental Manager

The Environmental Manager calculates the Area Source Emissions using the EIIP or other EPA guidance as provided. The Environmental Manager is also responsible for updating the SOP for the Emissions Group.


### 4. Description of equipment, forms, and/or software to be used

Equipment, Form, &/or Software	Who uses it?	Location
AP42	Senior Environmental Manager and Environmental Manager	EPA's website: <a href="http://www.epa.gov/ttn/chief/ap42/index.html">http://www.epa.gov/ttn/chief/ap42/index.html</a>
Emission Inventory Improvement Program (EIIP)	Senior Environmental Manager and Environmental Manager	EPA's website: <a href="http://www.epa.gov/ttn/chief/eiip/techreport/">http://www.epa.gov/ttn/chief/eiip/techreport/</a>
National Emission Inventory (NEI) Air Pollutant Emission Trends, U.S. EPA	Senior Environmental Manager and Environmental Manager	EPA's website <a href="http://www.epa.gov/ttn/chief/trends/">http://www.epa.gov/ttn/chief/trends/</a>

### 5. Procedure

#### 5.1 Procedural Flowchart

The procedural flowchart below titled "Area Source Inventory" is used to calculate non-point source inventories. This data is compiled every three years as mandated by EPA. The guidance followed is located in the EIIP. Emissions from area sources are calculated at the county level and consist of individual sources that are small, numerous and that have not been inventoried as specific point, mobile, or biogenic sources according to the EIIP.



## 5.2 Procedure

### Category 1: Stationary Fuel Combustion

#### Sub-Category 1.1: Industrial Fuel Combustion

SCC: 2102002000, 210200400, 2102005000, 2102006000, 2102007000

Follow these steps when calculating emissions from industrial fuel combustion:

1. Obtain statewide fuel consumption for “Other Industrial” for the following fuels: coal, distillate oil, natural gas, and liquefied petroleum gas (LPG). Use the Energy Information Administration’s website at <http://www.eia.doe.gov/> to find fuel consumption.

Note: As of the date of this SOP, the following steps will lead to data for fuel consumption.

- a. Go to <http://www.eia.doe.gov/>
  - b. Click on link for the various types of fuel consumption
  - c. Click on consumption tab for state totals
2. To avoid double calculating the various fuel combustions, subtract reported source totals from the total statewide fuel consumption by querying the total process rates for the various SCC codes using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb. The remaining number is the area source fuel consumption for the state.
  3. To distribute the remaining fuel to the county level, calculate the ratio of county to state employment for the manufacturing sector by dividing the number of Manufacturing Employees for each county by the number of manufacturing employees statewide. Use the County Business Patterns website at <http://www.census.gov/> to find the number of manufacturing employees for each county.

Note: As of the date of this SOP, the following steps will lead to data for Economic Census.

- a. Go to <http://www.census.gov/>
  - b. Click on Economic Census
  - c. Under 2002 Reports by State, use the down arrow key to select Indiana
  - d. Now, select each of the counties to find the county manufacturing employees
  - e. Use the total of employees for manufacturing under the paid employees’ column
4. Multiply the ratio calculated above in step 3 by the area source fuel consumption to distribute the fuel to the county level. The remaining number is the process rate for each county. Multiply the process rate by the appropriate EPA emission factors for the various fuels for industrial manufacturing found in AP-42, Fifth Edition, Volume 1, Chapter 1, External Combustion Sources at <http://www.epa.gov/ttn/chief/ap42/ch01/>.

#### Sub-Category 1.2: Commercial/Institutional Fuel Combustion

SCC: 2103004000, 2103005000, 2103006000, 2103007000

Follow these steps when calculating emissions from commercial/institutional fuel combustion:

1. Obtain statewide fuel consumption for “Commercial” for the following fuels: distillate fuel oil, liquefied petroleum gas (LPG), natural gas, and residual fuel oil. Use the Energy Information Administration’s website at <http://www.eia.doe.gov/> to find fuel consumption.

Note: Use the steps in sub-category 1.1-1 to navigate through the Energy Information Administration’s website.

2. To avoid double calculating the various fuel combustions, subtract reported source totals from the total statewide fuel consumption by querying the total process rates for the various fuels using the SIC codes greater than 4999 using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb. These are the SIC codes that identify all the commercial/institutional area sources.
3. To distribute the remaining fuel to the county level, calculate the ratio of county to state employment for the commercial/institutional sector by dividing the number of commercial/institutional employees for each county by the number of commercial/institutional employees statewide. Use the County Business Patterns website at <http://www.census.gov/> to find the number of commercial/institutional employees for each county.

Note: Use the steps in sub-category 1.1-3 to navigate through the U.S. Census Bureau's website.

4. Multiply the ratio calculated above in step 3 by the area source fuel consumption to distribute the fuel to the county level. The remaining number is the process rate for each county. Multiply the process rate by the appropriate EPA emission factors for the various fuels for commercial/institutional found in AP-42, Fifth Edition, Volume 1, Chapter 1, External Combustion Sources at <http://www.epa.gov/ttn/chief/ap42/ch01/>.

### **Sub-Category 1.3: Residential Fuel Combustion**

SCC: 2104002000, 2104004000, 2104006000, 2104007000

Follow these steps when calculating emissions from residential fuel combustion:

1. Obtain statewide fuel consumption for "Residential" for the following fuels: coal, distillate oil, natural gas, and liquid petroleum gas. Use the Energy Information Administration's website at <http://www.eia.doe.gov/> to find fuel consumption.

Note: Use the steps in sub-category 1.1-1 to navigate through the Energy Information Administration's website.

2. To distribute residential fuel to the county level, calculate the ratio of county fuel usage to statewide fuel usage using the breakdown of fuels by household per county divided by the breakdown of fuels by household per state using the U.S. Census Bureau's website at <http://www.census.gov/>.

Note: As of the date of this SOP, the following steps will lead to data for breakdown of fuels by household.

- a. Go to <http://www.census.gov/>
  - b. On the left hand side click on "American Fact Finder"
  - c. Using the drop down menu, click on Indiana
  - d. Scroll to "Housing Characteristics" and select "show more"
  - e. On the left hand side, select "change geography (state, county, place...)"
  - f. Using the drop down menu, select county, state, and each county name to obtain housing information
3. Multiply the ratio calculated above in step 3 by the area residential fuel use by state to distribute the fuel to the county level. The remaining number is the process rate for each county for the various fuels. Multiply the process rate by the appropriate EPA emission factors for the various fuels for residential found in AP-42, Fifth Edition, Volume 1, Chapter 1 External Combustion Sources at <http://www.epa.gov/ttn/chief/ap42/ch01/>.

#### **Sub-Category 1.4: Residential Heating Using Wood**

SCC: 2104008001, 2104008002, 2104008003, 2104008004, 2104008010, 2104008030, 2104008050

Follow these steps when calculating emissions from residential heating using wood:

1. Obtain statewide wood consumption for “Residential” using the Energy Information Administration’s website at <http://www.eia.doe.gov/>. To convert the statewide wood consumption from cords of wood consumed to tons, multiply the total cords consumed by 1.25.

Note: As of the date of this SOP, the following steps will lead to data for wood consumption.

- a. Go to <http://www.eia.doe.gov/>
  - b. Click on Households, Buildings & Industry
  - c. Under Consumption Summaries, click on “Annual”
  - d. Now, over to the right click on “State Energy”
  - e. Using the drop down menu at the bottom, select “Indiana”
  - f. Under “Consumption” click on the “Residential” document
2. Using the ratio estimates provided by EPA found in the “Documentation For The Final 2002 NONPOINT SECTOR (FEB 06 version) NATIONAL EMISSIONS INVENTORY FOR CRITERIA AND HAZARDOUS AIR POLLUTANTS” at <http://www.epa.gov/ttn/chief/net/2002inventory.html#documentaiton> the number calculated above in step 1 is broken out into three categories (fireplace without inserts, fireplaces with inserts and woodstoves).
  3. To distribute to the county level for the three categories above, calculate a ratio of county to state using the statewide total of households and the county total of households that burn wood found at the U.S. Census Bureau website <http://www.census.gov/>. The remaining number is the process rate for each county. Multiply the process rate by the appropriate EPA emission factors for each of the categories using the EIIP, Volume 3, Chapter 2, Residential Wood Combustion at [http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii02\\_apr2001.pdf](http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii02_apr2001.pdf).

Note: Use the steps in sub-category 1.3-2 to navigate through the Energy Information Administration’s website.

### **Category 2: Industrial Processes**

#### **Sub-Category 2.1: Bakeries**

SCC: 2302050000

Follow these steps when calculating emissions from bakeries:

1. Calculate a per capita consumption factor using the reported weight of yeast–raised product reported under the Bread, Cake, and Frozen Bakery Products from the Economic Census Bureau at <http://www.census.gov/econ/census02/> and the U.S. population at the U.S. Census Bureau at <http://census.gov/>.

Note: As of the date of this SOP, the following steps will lead to data for yeast-raised product.

- a. Go to <http://www.census.gov>
- b. Under Business & Industry open “Economic Census”
- c. Now open “Subject Series”
- d. Under Manufacturing, open the table “Product Summary”
- e. Use the yeast – raised product under Commercial Bakeries (NAICS code 311812) and Frozen cakes, pies, and other pastries manufacturing (NAICS code 311813)

2. Multiply the per capita consumption factor calculated above in step 1 by the Indiana population found at the U.S. Census Bureau at <http://www.census.gov>.  
Note: As of the date of this SOP, the following steps will lead to Indiana population data.
  - a. Go to <http://www.census.gov>
  - b. Under Population Finder, use the drop down menu to select Indiana
3. To avoid double calculating the amount consumed for the state, subtract the reported process rate for both the straight-dough and sponge-dough by querying the total process rates for the SCC 30203202 (straight-dough) and SCC 30203201 (sponge-dough) using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb.
4. Multiply the remaining process rate by the straight-dough emission factor of .5 lbs VOC/1,000 pounds baked found in the EIIP, Volume 3, Area Source Method Abstracts: Baked Goods at Commercial/Retail Bakeries at <http://www.epa.gov/ttn/chiep/eiip/techreport/volume03/index.html>.
5. Calculate a per capita factor by dividing the Indiana population found in step 2 by the remaining process rate. Now multiply the per capita factor by each of the county populations to calculate the VOC emissions for each county.

Note: As of the date of this SOP, the following steps will lead to county population data.

- a. Go to <http://www.census.gov>
- b. Under Population Finder, use the drop down menu to select Indiana
- c. Under "View more results", select the county table

### Category 3: Solvent Utilization

#### Sub-Category 3.1: Architectural Coatings

SCC: 2401001000

Follow these steps when calculating emissions from architectural coatings:

1. Calculate an emission factor for architectural coating area sources first by adding all the solvent-based paints together and all the water based paints together using the U.S. Census Bureau's website <http://www.census.gov>. Use Table 1 to select all solvent-based paints and Table 2 to select all water based paints.

Table 1  
National Solvent Coating Sales

Solvent Type	1,000 gallons
Exterior Solvent Type	XX
Interior Solvent Type	XX
Architectural Lacquers	XX
Architectural Coating N.S.K.	XX
<b>Total Solvents</b>	XX

Table 2  
National Water Based Coating Sales

Water Type	1,000 gallons
Exterior Water Type	XX
Interior Water Type	XX
<b>Total Water Type</b>	XX

Note: As of the date of this SOP, the following steps will lead to architectural coating data.

- a. Go to <http://www.census.gov>
  - b. Under Business & Industry, select more
  - c. Now select Current Industrial Reports (CIR)
  - d. Select CIRs by Subject
  - e. Tab down to find the report "Paints and Allied Products"
2. Now multiply the total national number for solvent-based paints by the average solvent-based coating content number (3.87 lbs VOC/gallon) and the total national number for water-based paints by the average water-based coating content number (0.74 lbs VOC/gal) found in the EIIP, Volume 3, Chapter 3: Architectural Surface Coating at <http://www.epa.gov/ttn/chief/eiip/techreport/volume03/archsfc.pdf>.
  3. Add the total solvent-based coatings and the water-based paints together for a total national VOC emission factor from architectural surface coating. Then divide this number by the total national population using the U.S. Census Bureau's website <http://www.census.gov>.
  4. Multiply the number calculated above in step 3 by each of the county populations to calculate the total emissions per county.

Note: Use the steps in sub-category 2.1-5 to navigate through the Census Bureau's website.

### **Sub-Category 3.2:      Automobile Refinishing**

SCC: 2401005000

Follow these steps when calculating emissions from automobile refinishing:

1. To avoid double calculating, first query the employees from the reported sources using the SIC 7532- Body Repair and Paint Shops using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb. Subtract this number from the county employment for the same SIC using the U.S. Census Bureau's website <http://www.census.gov>.

Note: As of the date of this SOP, the following steps will lead to county employment data.

- a. Go to <http://www.census.gov>
  - b. Under Business & Industry, select more
  - c. Now select the County Business Patterns report for county
  - d. Select Indiana
  - e. Select each of the counties to find the number of employees for the corresponding SIC or NAICS code
2. Multiply the emission factor 3,519 lbs VOC/employee found in the EIIP, Volume 3, Chapter 13 Auto Body Refinishing at <http://www.epa.gov/ttn/chief/eiip/techreport/volume03/archsfc.pdf> and the county employment found above in step 1 to calculate the VOC emissions for each county.

### **Sub-Category 3.3:      Traffic Markings**

SCC: 2401008000

Follow these steps when calculating for traffic markings:

1. First calculate the national emissions by finding the amount of sales for traffic marking paints from the U.S. Census Bureau's website <http://www.census.gov> and multiply 3.36 lb VOC/gallon the national average VOC content for water and solvent-based paints from the EIIP, Volume 3, Chapter 14, Traffic Markings at <http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii14.pdf>.



Note: As of the date of this SOP, the following steps will lead to traffic marking paints.

- a. Go to <http://www.census.gov>
  - b. Under Business & Industry, select more
  - c. Now select Current Industrial Reports (CIR)
  - d. Select CIRs by Subject
  - e. Tab down to find the report "Paints and Allied Products"
  - f. Use the quantity amount in 1000/gallons under "Traffic marking paints (all types: shelf goods and highway department)"
2. Allocate the national emissions calculated above in step 1 to the state level by dividing the amount of money spent in Indiana by the money spent nationally on highway maintenance using the category "Total Disbursements" at the Federal Highway Administration's website <http://www.fhwa.dot.gov/policy/ohim/hs04/htm/sf2.htm>.
  3. Calculate the emission factor for Indiana by dividing the state level emissions by the total number of roadway miles in Indiana, given by contacting the Program Development Division, Highway Statistics, Indiana Department of Transportation or the Office of Air Quality, Technical Support and Modeling Section's mobile inventory preparer.
  4. Multiply the emission factor by the total number of roadway miles in each county using the information supplied from above in step 3.

**Sub-Category 3.4: Industrial Surface Coating (employment based emission factor)**

SCC: 2401015000, 2401020000, 2401030000, 2401040000, 2401045000, 2401055000, 2401060000, 2401065000, 2401070000, 2401075000, 2401080000

Follow these steps when calculating for industrial surface coating using the employment based emission factor:

1. Calculate an employee based emission factor for the following SIC's in the table below running a query to find the point source employment for each of the SIC's and the reported VOC emissions for each using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb.

SCC	Description	SIC's
2401015000	Factory Finished Wood	2426-2429, 243-245, 2492, 2499
2401020000	Wood Furniture	25
2401030000	Paper Coating	26
2401040000	Metal Cans *	341
2401045000	Metal Coils *	3479
2401055000	Machinery and Equipment	35
2401060000	Appliances *	363
2401065000	Electronic and Other Electrical	3612, 3357
2401070000	New Motor Vehicles **	3711
2401075000	Other Transportation	37 (not 3711, 373)
2401080000	Marine Coatings	373

\* Use the National default emission factor because the reporting sources are low.  
\*\* Emissions reported in point source

2. Divide the reported VOC emissions for each of the SIC's by the reported employment for each SIC. Use this number for the emission factor.
3. Subtract the number of reported employees found in step 1 from each of the SIC county totals using the U.S. Census Bureau's website <http://www.census.gov>. Use the remaining number for the process rate for each of the counties.

Note: Use the steps in sub-category 3.2-1 to navigate through the County Business Patterns.

4. Multiply the process rates above found for each of the SIC's in step 4 by the emission factors found in step 3 to allocate the emissions to each of the counties.

### Sub-Category 3.5: Industrial Surface Coating (default emission factor)

SCC: 2401090000, 2401100000, 2401200000

Follow these steps when calculating emissions from industrial surface coating using the default emission factor:

1. Calculate industrial surface coating emissions using the default emission factor in the EIIP, Volume 3, Chapter 8, Industrial Surface Coating at <http://www.epa.gov/ttn/chiep/techreport/volume03/iii08.pdf> and multiply by the county populations found at the U.S. Census Bureau's website <http://www.census.gov>.

Note: Use the steps in 2.1-5 to navigate through U.S. Census Bureau's website.

SCC's	Description	Default Emission Factor
24-01-090-000	Miscellaneous Manufacturing	0.600 lbs VOC/person
24-01-100-000	Industrial Maintenance Coatings	0.800 lbs VOC/person
24-01-200-000	Other Special Purpose Coatings	0.800 lbs VOC/person

### Sub-Category 3.6: Degreasing

SCC: 2415230000, 2415245000, 2415345000, 2415360000

Follow these steps when calculating emissions from degreasing activities:

1. Use the U.S. Census Bureau to find employment numbers for each of the counties for the categories in Table 1 below at <http://www.census.gov>.

Note: Use the steps in 2.1-5 to navigate through U.S. Census Bureau's website.

Source Classification Codes and Industries Associated with Degreasing		
SCC	SIC	Description
2415230000	36	Electronic and other electronic equipment
	25	Furniture and fixtures
	33	Primary metal industries
	34	Fabricated metal products
	35	Industrial machinery and equipment
	37	Transportation equipment
	38	Instruments and related products

2415245000	39	Miscellaneous manufacturing industries
	417	Bus Terminal and Service Facilities
	423	Trucking terminal facilities
	551	New and used car dealers
	552	Used car dealers
	554	Gasoline service stations
	555	Boat dealers
	556	Recreational vehicle dealers
	753	Automotive repair shops
2415345000	25	Furniture and fixtures
	33	Primary metal industries
	34	Fabricated metal products
	35	Industrial machinery and equipment
	36	Electronic and other electronic equipment
	37	Transportation equipment
	38	Instruments and related products
	39	Miscellaneous manufacturing industries
2415345000 cont.		
2415360000	417	Bus Terminal and Service Facilities
	423	Trucking terminal facilities
	551	New and used car dealers
	552	Used car dealers
	554	Gasoline service stations
	555	Boat dealers
	556	Recreational vehicle dealers
	753	Automotive repair shops

- Run a query to find reported employment numbers for each of the categories in the table above using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb.
- Subtract the reported employment from the U.S Census Bureau's numbers to find the process rates for each of the counties.
- Calculate the VOC emissions by multiplying the default emission factor in the EIIP, Volume 3, Chapter 6, Solvent Cleaning at <http://www.epa.gov/ttn/chiep/techreport/volume03/iii06fin.pdf> and the process rate for each of the counties found in step 3.

### Sub-Category 3.7: Dry Cleaners

SCC: 2420010370

Follow these steps when calculating emissions from dry cleaners:

- Calculate an emission factor by finding the number of employees state wide and county wide for SIC 7216(Laundry and Garment Services) at the U.S. Census Bureau's website <http://www.census.gov>.

Note: Use the steps in 2.1-5 to navigate through U.S. Census Bureau's website

- Take the sum of the employment from the counties, multiply by 2000, and divide by the statewide total found in step 1. Use this number for the emission factor.
- Calculate the process rate by running a query to find the number of reported employees for SIC 7216 using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb and subtract this number from the county total.
- Multiply the process rate for each of the counties above by the emission factor to calculate for VOC emissions.

### **Sub-Category 3.8: Graphic Arts**

SCC: 2425000000

Follow these steps when calculating emissions from graphic arts activities:

1. Multiply the per capita factor found in the EIIP, Volume 3, Chapter 7, Graphic Arts at <http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii07.pdf> by the state population from the Census Bureau <http://www.census.gov> to find the total emissions for the state.

Note: Use the steps in 2.1-2 to navigate through the U.S. Census Bureau's website.

2. Develop an emission factor by subtracting point source emissions from the total emissions and dividing by the state population found in step 1.
3. Distribute to the counties by multiplying the emission factor by the population for each county.

Note: Use the steps in 2.1-5 to navigate through the U.S. Census Bureau's website.

### **Sub-Category 3.9: Rubber and Plastics**

SCC: 2430000000

Follow these steps when calculating emissions from rubber and plastics activities:

1. Run a query to find the total of reported emissions and number of reported employees for all SIC's beginning with 30 using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb.
2. Calculate the emission factor by dividing the point source emissions by the reported employees.
3. Subtract the reported employment for SIC's beginning with 30 from total employment for each of the counties.

Note: Use step 3.2-1 to navigate through the County Business Patterns.

4. Multiply the remaining number from above with the emission factor calculated in step 2.

### **Sub-Category 3.10: Miscellaneous Industrial Adhesives**

SCC: 2440020000

Follow these steps when calculating emissions from industrial adhesives activities:

1. Using the guidance in the Air Pollutant Emission Trends at <http://www.epa.gov/ttn/chief/trends>, calculate an emission factor by finding the total National Emissions from Industrial Adhesives and divide by the National Manufacturing Employment from the U.S. Census Bureau's website <http://www.census.gov>.

Note: As of the date of this SOP, the following steps will lead to emission trends data for industrial adhesives.

- a. Go to <http://www.epa.gov/air/airtrends/aqtrnd03/>
- b. Select "Appendix A –Data Tables"
- c. Search for industrial adhesives

Note: As of the date of this SOP, the following steps will lead to National Manufacturing Employment.

- a. Go to <http://www.census.gov>

- b. Select Economic Census
  - c. Now select "Businesses with paid employees"
  - d. Use the manufacturing number under "paid employees"
2. To avoid double calculating, run a query collecting sources reporting adhesives using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb. Subtract the reported employment from the total amount of manufacturing employment. The remaining number is the process rate.

### Sub-Category 3.11: Commercial/Consumer Solvents

SCC: 2460100000, 2460200000, 2460400000, 2460500000, 2460600000, 2460800000, 2460900000

Follow these steps when calculating emissions from commercial/consumer solvent usage:

1. Using the EIIP, Volume 3, Chapter 5, Consumer, and Commercial Solvent Use at <http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii05.pdf>, multiply the per capita factors for each of SCC codes by the population for each county from the U.S. Census Bureau's website <http://www.census.gov>.

Note: Use the steps in 2.1-5 to navigate through the U.S. Census Bureau's website.

Emission Factors for Commercial/Consumer Solvents

Source Classification Codes	Product Category	Per Capita Emission Factor (lb VOC/person)
2460100000	Personal Care Products	2.32
2460200000	Household Products	0.79
2460400000	Automotive Aftermarket Products	1.36
2460500000	Coatings and Related Products	0.95
2460600000	Adhesives and Sealants	0.57
2460800000	FIFRA-Regulated Products	1.78
2460900000	Miscellaneous Products	0.07

### Sub-Category 3.12: Asphalt Emulsions

SCC: 2461022000

Follow these steps when calculating emissions from asphalt emulsions:

1. To calculate the process rate, find the number of barrels of asphalt used for the state found at the State Energy Data website at [http://www.eia.doe.gov/emeu/states/seds\\_updates.html](http://www.eia.doe.gov/emeu/states/seds_updates.html).
2. Obtain the amount of roadway miles for the state and county from the Indiana Department of Transportation's, Division of Roadway Management Section.
3. Divide the county roadway miles by the state roadway miles and multiply by the total asphalt usage for the state found above in step 1.
4. Multiply the process rate by the default emission factor in the EIIP, Volume 3, Chapter 17, Asphalt Paving [http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii17\\_apr2001.pdf](http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii17_apr2001.pdf).

### **Sub-Category 3.13: Pesticide Usage**

SCC: 2461800000

Follow these steps when calculating emissions from pesticide usage:

1. Calculate pesticide usage by using a state specific emission factor. Develop the factor using a methodology that includes the retrieval of information of pesticides used, an emission factor for each pesticide used, a calculation about the inert ingredients in each pesticide, and an estimate of the amount of crop oil concentrate (an adjuvant used for the application of herbicides) used in the state of Indiana.
2. Find the amount of active ingredients for herbicides and insecticides applied to Indiana fields at the Indiana Agricultural Statistics Service at <http://www.usda.gov/nass/pubs/agr02/acro02.htm>.
3. Insert the numbers for both corn and soybeans to the Excel pesticide table found at K:\OAQ\_INV\Inv\pesticide.
4. Calculate the emission factor by adding the emissions from crop oil concentrates obtained in the pesticide Excel table, pesticides, and solvent carriers and then divide by the total number of acres of corn and soybeans in Indiana found at the National Agricultural Statistics Services, United States Department of Agriculture <http://www.nass.usda.gov/QuickStats/>.
5. Multiply the emission factor by the county-specific acreage for both corn and soybeans found at the National Agricultural Statistics Services, United States Department of Agriculture <http://www.nass.usda.gov/QuickStats/>.

### **Category 4: Petroleum Marketing**

Follow these steps when calculating emissions for bulk terminals:

#### **Sub-Category 4.1: Bulk Terminals**

SCC: 2501050120

1. Find the amount of gasoline sold in Indiana at the Federal Highway Administration, U.S. Department of Transportation <http://www.fhwa.dot.gov/policy/ohim/hs04/htm/mf21.htm>.
2. Find the amount of gasoline sold statewide and by county using, the NAICS code 447-Gasoline Service Station from the U.S. Census Bureau's, Economic Census at [http://www.census.gov/econ/census02/data/in/IN000\\_44.HTM#N447](http://www.census.gov/econ/census02/data/in/IN000_44.HTM#N447).
3. Run a query to find the amount of point source reported gasoline using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb and subtract from the amount sold statewide. Use this to allocate to each county.
4. Allocate the amount gasoline sold to each of the counties by dividing the amount of sales in each county by statewide sales and multiplying by the number of gallons sold statewide found above in step 1.
5. EPA guidance suggests that only 25% of all gasoline consumed goes through bulk plants. To calculate process rate, multiply each county by 25% to estimate the amount of fuel transferred through bulk terminals.
6. Multiply process rate by the emission factors in the table below:

Emission Factors	
Source	Emission Factor (lb VOC/1000) gal
Storage Tanks Breathing Loss	5.0
Storage Tank Working Loss - Filling	9.6
Storage Tank Working Loss - Emptying	3.8
Gasoline Loading Racks (Vapor balance controlled)	11.9 (0.3)
Total	30.3

7. Bulk terminals also have controls set forth in the Indiana rule (326 IAC 8-4). This rule says that any source of this type that is new after January 1, 1980 is required to make sure that any transfer between a tank and transport uses a submerged pipe vapor balance system. Using EPA's default rule effectiveness, multiply the number in step 2 by the Control Efficiency (CE) 38%, a Rule Effectiveness (RE) of 80%, and a Rule Penetration (RP) of 13%, i.e. process rate X emission factor X (1-(CE x RE x RP)) X 1 ton/2000 lb = VOC tons.

#### Sub-Category 4.2: Portable Fuel Containers

SCC: 2501011011, 2501011012, 2501011016, 2501012011, 2501012012, 2501012016

Follow these steps when calculating emissions for portable fuel containers:

1. Calculate the emissions for Commercial and Residential gas cans by using the method developed by the California Environmental Protection Agency's document Public Meeting to Consider Approval of California's Portable Gasoline-Container Emissions Inventory. Use the excel spreadsheet found at K:\OAQ\_INV\Inv\Area Source\Gasoline.zip to calculate the emissions for permeation, diurnal, and transport. Both the Spillage and Vapor losses are estimated in the nonroad emissions inventory by EPA models.
2. Using the survey results below in Table 1, estimate the number of fuel containers in the state for residential categories. The calculations are set up in an excel spreadsheet at K:\OAQ\_INV\Inv\Area Source\ Gasoline.zip\250101\GasCans.xls, insert the number of occupied housing, from the U.S. Census Bureau's website at <http://www.census.gov/>, in the space marked "households".

Note: As of the data of this SOP, the following steps will lead to number of households in Indiana.

- a. Go to <http://www.census.gov/>
- b. On the left hand side select American Fact finder
- c. Now select housing
- d. Under "Occupancy Status", select occupies housing units
- e. Now use the drop down menu and select Indiana

**Table 1**

Residential Survey Results	
Percentage of households with at least one gas can	46%
Number of gas cans per household	1.8
Percentage of plastic cans/metal cans	76% / 24%
Weighted average gas can capacity (gal)	2.34
Percentage of gas cans stored with fuel	70%
Weighted average stored fuel volume (% of capacity)	49%

Percentage of all gas cans that are plastic and stored open/closed	23% / 53%
Percentage of all gas cans that are metal and stored open/closed	11% / 13%
Percent of all cans stored open/closed	34% / 66%

- Using the survey results below in Table 2, estimate the number of fuel containers for commercial categories for the state. Do this by using the commercial population based on the number of identified businesses in Table 3 and insert into the excel spreadsheet at K:\OAQ\_INV\Inv\Area Source\ Gasoline.zip\250101\GasCans.xls.

**Table 2**

<b>Commercial Survey Results</b>	
Percentage of businesses with at least one gas can	80%
Number of gas cans per business	6.9
Percentage of plastic cans/metal cans	72% / 28%
Weighted average gas can capacity (gal)	3.43
Weighted average stored fuel volume (% of capacity)	49%
Percentage of all gas cans that are plastic and stored open/closed	39% / 33%
Percentage of all gas cans that are metal and stored open/closed	10% / 18%
Percent of all cans stored open/closed	49% / 51%

**Table 3**

<b>Category</b>	<b>NAICS</b>
Agricultural	115
Automotive Club and Towing Services	48841
Service Stations	8111
Lawn and Garden Maintenance Services	81141
General Contractors	23
Construction and Rental Yards	5324
Landscaping Services	561730

- Calculate permeable emissions separately for both residential and commercial by using the emission rates given in the California document. Use 1.57g/gal/day for plastic containers and 0.6g/gal/day for metal containers. Insert the numbers for both residential and commercial into the excel spreadsheet at K:\OAQ\_INV\Inv\Area Source\ Gasoline.zip\250101\GasCans.xls.
- Calculate diurnal emissions by inserting the numbers for both residential and commercial into the excel spreadsheet at K:\OAQ\_INV\Inv\Area Source\ Gasoline.zip\250101\GasCans.xls.
- Calculate transport spillage emissions by inserting the numbers for both residential and commercial into the excel spreadsheet at K:\OAQ\_INV\Inv\Area Source\ Gasoline.zip\250101\GasCans.xls

**Sub-Category 4.3: Service Station Tank Loading or Tank Truck Unloading (Stage 1)**

SCC: 2501060052 (uncontrolled), 2501060053 (controlled)

Follow these steps when calculating emissions from tank loading and unloading

- Find the amount of gasoline sold in Indiana at the Federal Highway Administration, U.S. Department of Transportation <http://www.fhwa.dot.gov/policy/ohim/hs04/htm/mf21.htm>.
- Find the amount of gasoline sold statewide and county wide by using the NAICS code 447-Gasoline Service Station from the U.S. Census Bureau's, Economic Census at [http://www.census.gov/econ/census02/data/in/IN000\\_44.HTM#N447](http://www.census.gov/econ/census02/data/in/IN000_44.HTM#N447).



3. Run a query to find the amount of point source reported gasoline using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb and subtract from the amount sold statewide. Use this to allocate to each county.
4. Allocate the amount sold to each of the counties by dividing the amount of sales in each county by statewide sales and multiplying by the number of gallons sold statewide found above in step 1.
5. Find the amount of gasoline tanks from the Underground Storage Tank data files from the Office of Land Quality, Indiana Department of Environmental Management  
<http://www.in.gov/idem/programs/land/ust/ust.html>.
6. Now copy the data into an Excel spreadsheet. Filter finding the tanks that have only gasoline. Also filter out the tanks that are “permanently out of service”, “suspended per inspection”, and “unregulated”.
7. Using the Petroleum Sources Applicability Rule 326 IAC 8-4-1, filter out the tanks that are located in Clark, Boone, Dearborn, Elkhart, Floyd, Hamilton, Hancock, Harrison, Hendricks, Johnson, Lake, Marion, Morgan, Porter, Saint Joseph, and Shelby counties.
8. To find the amount of balanced tanks in Indiana, use the total of gasoline tanks found in step 7 and divide by the number of tanks that constructed after 1985 through current year. Use the spreadsheet created in step 7 and filter out the tanks that constructed prior to 1985.
9. Now apply the percentage found in step 8 to the amount of gasoline found in each county.
10. Apply the controlled emission factor to only those counties identified in 326 IAC 8-4, i.e. Boone, Clark, Dearborn, Elkhart, Hamilton, Hancock, Harrison, Hendricks, Johnson, Lake, Marion, Morgan, Porter, Saint Joseph, and Shelby. Use the emission factors for stage 1 controlled and uncontrolled in the EIIP, Volume 3, Chapter 11, Gasoline Marketing (Stage 1 and Stage 2)  
[http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii11\\_apr2001.pdf](http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii11_apr2001.pdf).

#### **Sub-Category 4.4: Vehicle Fueling (Stage II) – Vapor Displacement**

SCC: 2501060101 (uncontrolled), 2501060102 (controlled)

Follow these steps when calculating emissions from vehicle fueling – Vapor Displacement:

1. Find the amount of gasoline sold in Indiana at the Federal Highway Administration, U.S. Department of Transportation <http://www.fhwa.dot.gov/policy/ohim/hs04/htm/mf21.htm>.
2. Find the amount of gasoline sold statewide and by county using the NAICS code 447-Gasoline Service Station from the U.S. Census Bureau's, Economic Census at [http://www.census.gov/econ/census02/data/in/IN000\\_44.HTM#N447](http://www.census.gov/econ/census02/data/in/IN000_44.HTM#N447).
3. Allocate the amount sold to each of the counties by dividing the amount of sales in each county by statewide sales and multiplying by the number of gallons sold statewide found above in step 1.
4. Calculate an emission factor using the input files supplied from the mobile model. Table 1 and Table 2 show examples of how the emission factors for January and July for the Southern Counties were calculated. By using these two months, the other months are distributed. Use the average of all months for the emission factor for the Southern counties. Use the same methodology for the Northern counties, Central Counties, Clark/Floyd, and Lake/Porter.

Table 1

January Run for Southern Counties

VTYPE	GM_MILE	MILES	MPG	VMT	G/GAL	Month	Factor
1	0.0628	29.4642	23.89	0.463793	0.322719	1	1.01
2	0.1058	35.2923	18.77	0.070491	0.009868	2	1.14
3	0.1058	35.2923	18.77	0.234672	0.109364	3	1.28
4	0.1486	34.0851	14.31	0.071379	0.010834	4	1.41
5	0.1486	34.0851	14.31	0.032825	0.002291	5	1.55
6	0.2152	35.8919	9.88	0.028896	0.001775	6	1.69
7	0.2342	32.3617	9.08	0.001027	2.24E-06	7	1.82
8	0.2465	19.9098	8.63	0.000522	5.8E-07	8	1.69
9	0.2719	27.6093	7.82	0.001164	2.88E-06	9	1.55
10	0.2733	27.4686	7.78	0.002489	1.32E-05	10	1.41
11	0.2972	24.3758	7.15	0.001132	2.72E-06	11	1.28
12	0.3169	23.6257	6.71	0.000004	3.4E-11	12	1.14
25	0.3421	27.2301	6.22	0.000496	5.23E-07	Sum	16.97
					0.456873	g/gal	Average
					1.007222	lb/E3gal	1.41

Table 2  
July Run for Southern Counties

VTYPE	GM_MILE	MILES	MPG	VMT	G/GAL
1	0.1144	29.1752	23.9	0.456768	0.570447
2	0.1955	34.8826	18.75	0.071404	0.018689
3	0.1955	34.8826	18.75	0.237712	0.207133
4	0.2882	33.944	14.3	0.072838	0.021865
5	0.2882	33.944	14.3	0.033496	0.004624
6	0.4164	35.8288	9.9	0.029201	0.003515
7	0.4529	32.4716	9.1	0.001038	4.44E-06
8	0.4763	19.6757	8.66	0.000509	1.07E-06
9	0.5264	27.4602	7.83	0.00116	5.55E-06
10	0.5283	27.3328	7.8	0.002482	2.54E-05
11	0.5749	24.2458	7.17	0.001122	5.19E-06
12	0.6128	23.3718	6.73	0.000004	6.6E-11
25	0.6629	27.2301	6.22	0.000485	9.7E-07
					0.826316 g/gal
					1.821697 lb/E3gal

5. Multiply the process rate in step 4 by the emission factor found in the mobile model.

#### Sub-Category 4.5: Vehicle Fueling (Stage II) – Spillage

SCC: 2501060103

Follow these steps when calculating emissions from vehicle fueling – Spillage:

1. Find the amount of gasoline sold in Indiana at the Federal Highway Administration, U.S. Department of Transportation <http://www.fhwa.dot.gov/policy/ohim/hs04/htm/mf21.htm>.
2. Find the amount of gasoline sold statewide and by county using the NAICS code 447-Gasoline Service Station from the U.S. Census Bureau's, Economic Census at [http://www.census.gov/econ/census02/data/in/IN000\\_44.HTM#N447](http://www.census.gov/econ/census02/data/in/IN000_44.HTM#N447).
3. Allocate the amount sold to each of the counties by dividing the amount of sales in each county by statewide sales and multiplying by the number of gallons sold statewide found above in step 1.
4. Apply the emission factor 0.7 lb VOC/1000 gallons in AP-42, Fifth Edition, Volume 1, Chapter 5, Petroleum Industry, Transportation, and Marketing of Petroleum Liquids <http://www.epa.gov/ttn/chieff/ap42/ch05/final/c05s02.pdf> to the process rate found in step 4.

#### Sub-Category 4.6: Underground Tank Breathing

SCC: 2501060200

Follow these steps when calculating emissions from underground tank breathing:

1. Find the amount of gasoline sold in Indiana at the Federal Highway Administration, U.S. Department of Transportation <http://www.fhwa.dot.gov/policy/ohim/hs04/htm/mf21.htm>.
2. Find the amount of gasoline sold statewide and by county using the NAICS code 447-Gasoline Service Station from the U.S. Census Bureau's, Economic Census at [http://www.census.gov/econ/census02/data/in/IN000\\_44.HTM#N447](http://www.census.gov/econ/census02/data/in/IN000_44.HTM#N447).

3. Allocate the amount sold to each of the counties by dividing the amount of sales in each county by statewide sales and multiplying by the number of gallons sold statewide found above in step 1.
4. Apply the emission factor 1.0 lb VOC/1000 gallons in AP-42, Fifth Edition, Volume 1, Chapter 5, Petroleum Industry, Transportation, and Marketing of Petroleum Liquids <http://www.epa.gov/ttn/chief/ap42/ch05/final/c05s02.pdf> to the process rate found in step 4.

#### **Sub-Category 4.7: Tank Trucks in Transit**

SCC: 2505030120

Follow these steps when calculating emissions from tank trucks in transit:

1. Find the amount of gasoline sold in Indiana at the Federal Highway Administration, U.S. Department of Transportation <http://www.fhwa.dot.gov/policy/ohim/hs04/htm/mf21.htm>.
2. Find the amount of gasoline sold statewide and by county using the NAICS code 447-Gasoline Service Station from the U.S. Census Bureau's, Economic Census at [http://www.census.gov/econ/census02/data/in/IN000\\_44.HTM#N447](http://www.census.gov/econ/census02/data/in/IN000_44.HTM#N447).
3. Allocate the amount sold to each of the counties by dividing the amount of sales in each county by statewide sales and multiplying by the number of gallons sold statewide found above in step 1.
4. Using the guidance in the EIIP, Volume 3, Chapter 11, Gasoline Marketing (Stage I and State II) at [http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii11\\_apr2001.pdf](http://www.epa.gov/ttn/chief/eiip/techreport/volume03/iii11_apr2001.pdf), multiply the activity rate 1.25 by the amount sold per county found in step 4.
5. Now multiply the process rate found in step 5 by the emission factor .06 lb VOC/gallon transported using the EIIP guidance above.

### **Category 5: Waste Management Practices**

#### **Sub-Category 5.1: Solid Waste Incineration**

##### **5.1.1: Industrial Solid Waste Incineration**

SCC: 2601010000

Follow these steps when calculating emissions from industrial solid waste incineration:

1. Find the number of manufacturing employees, NAICS code 31, for each county using the County Business Patterns at the U.S. Census Bureau's website <http://censtats.census.gov/cgi-bin/cbpnaic/cbpsel.pl>.

Note: Use the steps in 3.2-1 to navigate through the county business patterns.

2. Multiply the county manufacturing employment by the default fuel-loading factor 420 tons / 1,000 manufacturing employees.
3. Multiply the process rate in step 2 by AP-42, Fifth Edition, Volume 1, Chapter 2-1.12, Solid Waste Disposal at <http://www.epa.gov/ttn/chief/ap42/ch02/index.html>.

### 5.1.2: Commercial Solid Waste Incineration

SCC: 2601020000

Follow these steps when calculating emissions from commercial solid waste incineration:

1. Find the population for each county at the U.S. Census Bureau's website <http://www.census.gov/>.  
  
Note: Use steps 2.1-5 to navigate through the U.S. Census Bureau's website.
2. Next find the default factor of .65lb/person/day from U.S. EPA Municipal Solid Waste Report <http://www.epa.gov/epaoswer/non-hw/muncpl/msw99.htm>.
3. Find the percent of commercial solid waste from the U.S. EPA Municipal Solid Waste Report above.
4. Now, calculate the process rate for commercial solid waste incineration by multiplying population by the default factor of .65lb/person/day by the percent of commercial solid waste and number of days in a year.
5. Multiply the process rate in step 4 by AP-42, Fifth Edition, Volume 1, Chapter 2-1.12, Solid Waste Disposal at <http://www.epa.gov/ttn/chief/ap42/ch02/index.html>.

### 5.1.3: Residential Solid Waste Incineration

SCC: 2601030000

Follow these steps when calculating emissions from residential solid waste incineration:

1. Find the population for each county at the U.S. Census Bureau's website <http://www.census.gov/>.  
  
Note: Use step 2.1-5 to navigate through the U.S. Census Bureau's website.
2. Next find the default factor of .65lb/person/day from U.S. EPA Municipal Solid Waste Report <http://www.epa.gov/epaoswer/non-hw/muncpl/msw99.htm>.
3. Find the percent of residential solid waste from the U.S. EPA Municipal Solid Waste Report above.
4. Now, calculate the process rate for residential solid waste incineration by multiplying population by the default factor of .65lb/person/day by the percent of commercial solid waste and number of days in a year.
5. Multiply the process rate in step 4 by AP-42, Fifth Edition, Volume 1, Chapter 2-1.12, Solid Waste Disposal at <http://www.epa.gov/ttn/chief/ap42/ch02/index.html>.

## Sub-Category 5.2: Residential Open Burning

### 5.2.1: Leaf and Brush Burning

SCC: 2610000100 and 2610000400

Follow these steps when calculating emissions from leaf and brush burning:

1. Find a per capita factor for leaf burning and a per capita for brush burning by using the U.S. EPA's Solid Waste Report at <http://www.epa.gov/epaoswer/non-hw/muncpl/msw99.htm>.
2. Allocate the amount burned by adjusting the per capita factor for leaves at 25% and for brush at 25%. Of the total waste generated only 28% burns.

- Once all the percentages from above are calculated, multiply the adjusted per capita factor by the rural population for each county from the U.S. Census Bureau at <http://www.census.gov/>

Note: As of the data of this SOP, the following steps will lead to county rural population.

- Go to <http://www.census.gov/>
  - On the left hand side, select American Fact Finder
  - Select data sets
  - Detailed tables
  - County
  - Indiana
  - All counties
- Use the table below to adjust the amount of waste generated to account for the percentage of forest in each county. The percentages come from a document from the United States Department of Agriculture at [http://ncrs.fs.fed.us/pubs/rb/rb\\_nc253b.pdf](http://ncrs.fs.fed.us/pubs/rb/rb_nc253b.pdf).

Percent Forested Acres per County	Adjusted for Yard Waste Generated
< 10%	0% generated
>= 10%, and < 50%	50% generated
>= 50%	100% generated

- Now, multiply the amount of leaves and brush by the emission factors found in AP-42, Fifth Edition, Volume 1, Chapter 2, Solid Waste Disposal, Table 2.5-5, and Table 2.5-6 at <http://www.epa.gov/ttn/chief/ap42/ch02/final/c02s05.pdf>.

### 5.2.2: Residential Waste Incineration

SCC: 2610030000

Follow these steps when calculating emissions from for residential waste incineration:

- Find a per capita factor for residential waste incineration by using the U.S. EPA's Solid Waste Report at <http://www.epa.gov/epaoswer/non-hw/muncpl/pubs/mswchar05.pdf>.
- Using the Solid Waste Report above, subtract the percentage of recycled and composted material from the per capita factor above.
- Now, subtract the percentages of combustibles i.e. glass, metal, yard trimmings, and other waste.
- Using a document from EPA, it states that only 28% of waste generated by rural population burns and of that percent, 49% is actually combusted. Using this information multiply the per capita factor by 0.28 and then multiply that number by 0.49 actually burned in rural counties.
- Once all the percentages are calculated, multiply the adjusted per capita factor by the rural population for each county from the U.S. Census Bureau at <http://www.census.gov/>.

Note: Use steps 5.2.1-3 to find county rural population.

- Calculate the amount of residential waste by the emission factors in the EIIP, Volume 3, Chapter 16, Open Burning at <http://www.epa.gov/ttn/chief/eiip/techreport/volume03/index.html>.

### Sub-Category 5.3: Public Owned Treatment Works (POTW's)

SCC: 2630020000

Follow these steps when calculating emissions from POTW's:

1. To calculate the amount of annual flow for public owned treatment works, obtain the amount of monthly flow rate for each county. This is data is supplied by the Office of Water Quality. To calculate for annual flow multiply the monthly flow by the default of 0.16 that represents the amount of industrial flow.
2. Calculate the process rate above by the emission factors in FIRE 6.25 using the SCC code 2630020000.

#### **Sub-Category 5.4: Treatment, Storage, and Disposal Facilities**

SCC: 2640000004

Follow these steps when calculating emissions from treatment, storage, and disposal facilities:

1. Obtain a list of treatment facilities and the amount of ignitable waste from each facility from IDEM's Office of Land Quality.
2. Using the list of facilities from step 1, run a query using the ACCESS data tables at K:\OAQ\_INV\Steptool\Stptl\_02.mdb to obtain the amount of ignitable waste reported to IDEM's Office of Air Quality.
3. Compare the two lists obtained in step 1 and step 2, for each facility subtract any quantity reported to OAQ from the quantity reported to OLQ. Do this in order to avoid double counting quantities reported to both offices. Combine the quantities reported from facilities within the same counties. Use these quantities as the process rate for each county.
4. Multiply the process rate above with the combined emission factor in the table below:

<b>Emission Source</b>	<b>Emission Factor in AP-42 (lb VOC/Ton)</b>	<b>Emission Factor Used (lb VOC/Ton)</b>
Storage Tank Vent	0.004-0.09	0.09
Spillage (filling)	0.20	0.20
Loading (filling)	0.00024-1.42	1.42
Spillage (emptying)	0.20	0.20
Loading (emptying)	0.00024-1.42	1.42
<b>Combined Emission Factor</b>		<b>3.33</b>

### **Category 6: Submit Data to EPA**

Submit data in a format that is acceptable to EPA. At the present time the format is the National Emission Inventory (NEI).

#### **6. Standards and checklists**

The Emission Reporting program does not have any checklist for the Area Source Inventory at this time. The Emission Group does this electronically through an excel spreadsheet that is created when needed.

#### **7. Records Management**

The Area Source Inventory files are kept electronically at K:\OAQ\_INV\Inv\Area Source.

The Branch Contact for the Air Programs Branch and the Section contact for the Technical Support and Modeling Section will keep copies of the SOPs for the Technical Support and Modeling Section to be referenced as needed. An electronic copy will also be available on K:\OAQ\_INV\SOPs.

## 8. Quality Assurance / Quality Control

Comparisons are made against the emissions estimates made by The U.S. EPA in the NEI.

## 9. Continuous Improvement Cycle

A periodic review will be completed per updates and changes made to the EIIP.

## 10. References

The Area Source Inventory is a requirement of 40 CFR Part 51 Subpart A - Emission Inventory Reporting Requirements.

## 11. History of Revisions

<b>Date Month/day/year</b>	<b>Revision Number</b>	<b>Description</b>
02/27/2008	1	Revised using new SOP template.

## 12. Appendices

None



**This page left intentionally blank.**