


Crawford County Nutrient and Sediment Load




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 23,680,315 lbs.
Enough to fill 118 train cars.



x 118

Nitrogen Reduced 20,165 lbs.
Enough to fill 20 8' truck beds.



x 20

Phosphorus Reduced 10,076 lbs.
Enough to fill 10 8' truck beds.



x 10

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	64	64	14,985,765	6,851	13,707
2014	48	88	14,056,450	6,205	12,415
2015	86	154	23,569,755	10,405	20,818
2016	52	167	22,701,500	9,918	19,846
2017	67	208	27,190,395	11,845	23,703
2018	102	264	30,664,435	13,252	26,520
2019	43	257	23,680,315	10,076	20,165

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA