


Dearborn County Nutrient and Sediment Load




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding Percentage by Source



Comparison of Land Use Across County



Sediment Reduced 9,776,770 lbs.

Enough to fill 49 train cars.



x 49

Nitrogen Reduced 10,198 lbs.

Enough to fill 10 8' truck beds.



x 10

Phosphorus Reduced 5,107 lbs.

Enough to fill 5 8' truck beds.



x 5

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	131	131	6,344,520	3,537	7,070
2014	79	130	5,798,445	3,219	6,426
2015	54	137	5,430,770	3,013	6,017
2016	34	144	4,508,285	2,387	4,764
2017	49	184	6,909,550	3,612	7,211
2018	55	209	6,652,705	3,321	6,626
2019	106	311	9,776,770	5,107	10,198

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA