



Delaware County Nutrient and Sediment Load




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 15,846,715 lbs.
Enough to fill 79 train cars.



x 79

Nitrogen Reduced 20,656 lbs.
Enough to fill 21 8' truck beds.



x 21

Phosphorus Reduced 10,339 lbs.
Enough to fill 10 8' truck beds.



x 10

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	14	14	1,265,800	827	1,652
2014	20	25	1,607,630	1,050	2,098
2015	28	34	1,644,310	1,072	2,138
2016	19	30	956,385	594	1,183
2017	29	44	2,120,300	1,380	2,754
2018	53	70	5,886,850	3,580	7,150
2019	153	185	15,846,715	10,339	20,656

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA