


Dubois County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 111,589,910
Enough to fill 558 train cars.



x 558

Nitrogen Reduced 97,183 lbs.
Enough to fill 97 8' truck beds.



x 97

Phosphorus Reduced 48,574 lbs.
Enough to fill 49 8' truck beds.



x 49

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	159	159	57,044,795	24,309	48,650
2014	246	314	119,249,785	51,642	103,381
2015	286	393	133,473,935	57,698	115,421
2016	299	483	123,167,005	53,521	107,132
2017	171	458	88,895,620	39,067	78,167
2018	271	610	102,373,615	44,810	89,660
2019	237	728	111,589,910	48,574	97,183

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA