



# Huntington County Nutrient and Sediment Load




Accomplished By Private Landowners  
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding  
Percentage by Source




Comparison of Land  
Use Across County



**Sediment Reduced 19,002,295 lbs.**

Enough to fill 95 train cars.



x 95

**Nitrogen Reduced 25,819 lbs.**

Enough to fill 26 8' truck beds.



x 26

**Phosphorus Reduced 12,967 lbs.**

Enough to fill 13 8' truck beds.



x 13

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	112	112	8,469,665	5,425	10,842
2014	48	66	5,359,280	3,201	6,401
2015	184	213	11,849,255	7,886	15,702
2016	41	136	10,064,200	6,703	13,332
2017	63	187	13,612,130	9,373	18,649
2018	113	309	18,105,045	12,434	24,764
2019	79	291	19,002,295	12,967	25,819

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA