


Warren County Nutrient and Sediment Load Reductions




Accomplished By Private Landowners
and the Indiana Conservation Partnership




Practices Installed Over Time



Conservation Funding
Percentage by Source



Comparison of Land
Use Across County



Sediment Reduced 15,903,435 lbs.
Enough to fill 80 train cars.



x 80

Nitrogen Reduced 19,122 lbs.
Enough to fill 19 8' truck beds.



x 19

Phosphorus Reduced 9,610 lbs.
Enough to fill 10 8' truck beds.



x 10

Calendar Year	Practices Installed	Active Practices	Sediment Reduction (lbs)	Phosphorus Reduction (lbs)	Nitrogen Reduction (lbs)
2013	130	130	15,398,915	9,338	18,593
2014	83	115	11,691,045	7,107	14,124
2015	85	138	21,677,375	13,237	26,376
2016	86	144	18,087,100	11,127	22,158
2017	97	165	21,858,225	12,975	25,850
2018	163	246	27,344,610	16,503	32,902
2019	106	203	15,903,435	9,610	19,122

The "practices installed" column indicates the number of newly installed best management practices within a given calendar year, while the "active practices" column indicates the number of best management practices that are actively reducing sediment, nitrogen, and phosphorus loading regardless of the year of installation. Practices do not include the many unassisted practices designed and installed by private landowners without ICP assistance. Nutrient estimates only consider sediment bound N and P, not dissolved components. Load reductions are calculated using the EPA's Region 5 Load Reduction Model, and rounded to nearest integer for display.

Data provided by: Indiana State Department of Agriculture, Indiana Department of Natural Resources, Indiana Department of Environmental Management, Indiana Soil and Water Conservation Districts, and the USDA Natural Resource Conservation Service.

Last Updated: 5/15/20, Sam Stroebe ISDA