

**Leading Causes of Death for Women  
Knox County vs. Indiana, 1995-1999**

|                 |                   | Knox County | Indiana |
|-----------------|-------------------|-------------|---------|
| All Causes      | Deaths            | 1,266       | 136,704 |
|                 | Age-Adjusted Rate | 806.87      | 778.40  |
| Cardiovascular  | Deaths            | 641         | 59,221  |
|                 | Age-Adjusted Rate | 383.38 s    | 326.70  |
| Homicide        | Deaths            | 4           | 555     |
|                 | Age-Adjusted Rate | * u         | 3.65    |
| HIV/AIDS        | Deaths            | 0           | 109     |
|                 | Age-Adjusted Rate | * u         | 0.72    |
| Lung Cancer     | Deaths            | 55          | 7,517   |
|                 | Age-Adjusted Rate | 41.20       | 45.37   |
| Breast Cancer   | Deaths            | 37          | 4,778   |
|                 | Age-Adjusted Rate | 28.68       | 29.16   |
| Ovarian Cancer  | Deaths            | 15          | 1,564   |
|                 | Age-Adjusted Rate | * u         | 9.48    |
| Cervical Cancer | Deaths            | 5           | 522     |
|                 | Age-Adjusted Rate | * u         | 3.31    |

s = significantly different from the state rate  
u = unstable rate (numerator less than 20)

Source: Indiana State Department of Health,  
Epidemiology Resource Center, Data Analysis Team