

State Prevalence Rankings*: BRFSS, 2015

| State | Current Smokers | | Adults 18-64 without Health Care Coverage | | Obese Based on BMI | | Diabetes | | High Blood Pressure | | <150 Minutes of Physical Activity in Past Week | | High Cholesterol | |
|----------------------|-----------------|-----------|---|-----------|--------------------|-----------|-------------|-----------|---------------------|-----------|--|----------|------------------|----------|
| | % | Rank | % | Rank | % | Rank | % | Rank | % | Rank | % | Rank | % | Rank |
| Alabama | 21.4 | 10 | 16.4 | 15 | 35.6 | 2 | 13.5 | 3 | 40.4 | 3 | 55.4 | 4 | 42.0 | 1 |
| Alaska | 19.1 | 16 | 17.2 | 11 | 29.8 | 26 | 7.6 | 48 | 27.5 | 48 | 41.7 | 47 | 33.9 | 45 |
| Arizona | 14.0 | 45 | 17.6 | 8 | 28.4 | 34 | 10.1 | 24 | 30.8 | 27 | 46.2 | 32 | 37.3 | 18 |
| Arkansas | 24.9 | 3 | 15.2 | 17 | 34.5 | 6 | 12.6 | 7 | 39.3 | 4 | 54.9 | 5 | 41.3 | 2 |
| California | 11.7 | 50 | 13.5 | 24 | 24.2 | 47 | 10.0 | 25 | 28.5 | 46 | 42.7 | 43 | 34.2 | 43 |
| Colorado | 15.7 | 37 | 12.9 | 27 | 20.2 | 51 | 6.8 | 51 | 25.7 | 50 | 39.4 | 51 | 31.5 | 50 |
| Connecticut | 13.5 | 48 | 8.7 | 45 | 25.3 | 42 | 9.3 | 31 | 30.4 | 30 | 45.5 | 35 | 37.4 | 16 |
| Delaware | 17.4 | 27 | 10.3 | 35 | 29.7 | 28 | 11.5 | 10 | 34.5 | 12 | 51.5 | 15 | 39.1 | 6 |
| District of Columbia | 16.0 | 33 | 6.7 | 50 | 22.1 | 50 | 8.5 | 39 | 29.4 | 41 | 42.1 | 45 | 31.3 | 51 |
| Florida | 15.8 | 36 | 20.9 | 3 | 26.8 | 35 | 11.3 | 15 | 33.5 | 16 | 48.4 | 27 | 38.0 | 13 |
| Georgia | 17.7 | 23 | 20.8 | 4 | 30.7 | 19 | 11.3 | 15 | 36.2 | 9 | 52.0 | 12 | 36.3 | 26 |
| Hawaii | 14.1 | 44 | 8.8 | 43 | 22.7 | 49 | 8.5 | 39 | 32.0 | 23 | 43.4 | 41 | 36.4 | 25 |
| Idaho | 13.8 | 47 | 17.5 | 9 | 28.6 | 33 | 8.1 | 45 | 31.3 | 25 | 44.7 | 38 | 37.6 | 15 |
| Illinois | 15.1 | 41 | 11.0 | 32 | 30.8 | 18 | 9.9 | 26 | 30.8 | 27 | 50.2 | 18 | 36.0 | 32 |
| Indiana | 20.6 | 12 | 14.0 | 21 | 31.3 | 15 | 11.4 | 13 | 32.4 | 21 | 55.9 | 2 | 39.1 | 6 |
| Iowa | 18.1 | 21 | 7.8 | 47 | 32.1 | 12 | 8.8 | 36 | 30.6 | 29 | 51.2 | 16 | 36.2 | 27 |
| Kansas | 17.7 | 23 | 15.7 | 16 | 34.2 | 7 | 9.7 | 29 | 31.6 | 24 | 50.0 | 20 | 37.4 | 16 |
| Kentucky | 26.0 | 1 | 8.1 | 46 | 34.6 | 5 | 13.4 | 4 | 39.0 | 6 | 54.8 | 6 | 40.7 | 3 |
| Louisiana | 21.9 | 7 | 18.6 | 6 | 36.2 | 1 | 12.7 | 5 | 39.3 | 4 | 53.8 | 8 | 39.2 | 5 |
| Maine | 19.5 | 15 | 12.1 | 28 | 30.0 | 24 | 9.9 | 26 | 34.1 | 14 | 46.1 | 33 | 38.5 | 11 |
| Maryland | 15.1 | 41 | 10.4 | 34 | 28.9 | 31 | 10.3 | 23 | 32.5 | 19 | 47.1 | 30 | 35.9 | 33 |
| Massachusetts | 14.0 | 45 | 6.0 | 51 | 24.3 | 46 | 8.9 | 35 | 29.6 | 38 | 48.2 | 28 | 34.5 | 41 |
| Michigan | 20.7 | 11 | 12.0 | 30 | 31.2 | 16 | 10.7 | 18 | 33.1 | 18 | 47.9 | 29 | 38.2 | 12 |
| Minnesota | 16.2 | 32 | 7.3 | 48 | 26.1 | 39 | 7.6 | 48 | 26.3 | 49 | 45.1 | 37 | 32.3 | 48 |
| Mississippi | 22.6 | 4 | 21.6 | 2 | 35.6 | 2 | 14.7 | 1 | 42.4 | 2 | 62.0 | 1 | 38.7 | 10 |
| Missouri | 22.3 | 5 | 15.1 | 18 | 32.4 | 10 | 11.5 | 10 | 34.1 | 14 | 49.5 | 23 | 37.1 | 19 |
| Montana | 18.9 | 19 | 15.1 | 18 | 23.6 | 48 | 7.9 | 47 | 29.1 | 45 | 41.8 | 46 | 33.1 | 47 |
| Nebraska | 17.1 | 29 | 14.4 | 20 | 31.4 | 14 | 8.8 | 36 | 29.9 | 35 | 48.7 | 26 | 35.1 | 38 |
| Nevada | 17.6 | 25 | 17.7 | 7 | 26.7 | 36 | 9.7 | 29 | 28.3 | 47 | 45.5 | 35 | 36.7 | 21 |
| New Hampshire | 15.9 | 35 | 8.8 | 43 | 26.3 | 38 | 8.1 | 45 | 29.2 | 44 | 42.4 | 44 | 35.7 | 34 |
| New Jersey | 13.5 | 48 | 13.8 | 22 | 25.6 | 41 | 9.0 | 33 | 30.9 | 26 | 51.1 | 17 | 35.4 | 36 |
| New Mexico | 17.5 | 26 | 13.6 | 23 | 28.8 | 32 | 11.5 | 10 | 30.0 | 32 | 44.0 | 40 | 34.3 | 42 |
| New York | 15.2 | 39 | 12.1 | 28 | 25.0 | 44 | 9.8 | 28 | 29.3 | 43 | 52.9 | 11 | 36.6 | 24 |
| North Carolina | 19.0 | 18 | 19.1 | 5 | 30.1 | 22 | 10.7 | 18 | 35.2 | 11 | 51.9 | 14 | 37.0 | 20 |
| North Dakota | 18.7 | 20 | 9.7 | 39 | 31.0 | 17 | 8.7 | 38 | 30.4 | 30 | 53.0 | 10 | 35.0 | 39 |
| Ohio | 21.6 | 9 | 10.2 | 36 | 29.8 | 26 | 11.0 | 17 | 34.3 | 13 | 49.8 | 21 | 36.7 | 21 |
| Oklahoma | 22.2 | 6 | 16.7 | 13 | 33.9 | 8 | 11.7 | 9 | 36.2 | 9 | 53.4 | 9 | 38.0 | 13 |
| Oregon | 17.1 | 29 | 10.2 | 36 | 30.1 | 22 | 10.7 | 18 | 30.0 | 32 | 39.6 | 50 | 36.7 | 21 |
| Pennsylvania | 18.1 | 21 | 10.0 | 38 | 30.0 | 24 | 10.4 | 21 | 32.5 | 19 | 50.2 | 18 | 36.2 | 27 |
| Rhode Island | 15.5 | 38 | 9.4 | 41 | 26.0 | 40 | 9.0 | 33 | 32.4 | 21 | 49.6 | 22 | 35.2 | 37 |
| South Carolina | 19.7 | 14 | 17.5 | 9 | 31.7 | 13 | 11.8 | 8 | 37.8 | 8 | 49.5 | 23 | 38.9 | 9 |
| South Dakota | 20.1 | 13 | 10.6 | 33 | 30.4 | 21 | 9.3 | 31 | 30.0 | 32 | 46.4 | 31 | 33.3 | 46 |
| Tennessee | 21.9 | 7 | 16.6 | 14 | 33.8 | 9 | 12.7 | 5 | 38.5 | 7 | 54.6 | 7 | 40.0 | 4 |
| Texas | 15.2 | 39 | 27.5 | 1 | 32.4 | 10 | 11.4 | 13 | 29.5 | 40 | 55.7 | 3 | 36.1 | 30 |
| Utah | 9.1 | 51 | 13.3 | 25 | 24.5 | 45 | 7.0 | 50 | 23.6 | 51 | 44.7 | 38 | 32.2 | 49 |
| Vermont | 16.0 | 33 | 7.2 | 49 | 25.1 | 43 | 8.2 | 44 | 29.4 | 41 | 41.1 | 49 | 34.0 | 44 |
| Virginia | 16.5 | 31 | 13.0 | 26 | 29.2 | 29 | 10.4 | 21 | 33.2 | 17 | 49.0 | 25 | 35.6 | 35 |
| Washington | 15.0 | 43 | 12.0 | 30 | 26.4 | 37 | 8.4 | 41 | 29.7 | 37 | 41.7 | 47 | 36.2 | 27 |
| West Virginia | 25.7 | 2 | 9.6 | 40 | 35.6 | 2 | 14.5 | 2 | 42.7 | 1 | 52.0 | 12 | 39.0 | 8 |
| Wisconsin | 17.3 | 28 | 9.2 | 42 | 30.7 | 19 | 8.4 | 41 | 29.6 | 38 | 43.2 | 42 | 36.1 | 30 |
| Wyoming | 19.1 | 16 | 17.0 | 12 | 29.0 | 30 | 8.4 | 41 | 29.9 | 35 | 45.7 | 34 | 35.0 | 39 |
| Range | 9.1-26.0 | | 6.0-27.5 | | 20.2-36.2 | | 6.8-14.7 | | 23.6-42.7 | | 39.4-62.0 | | 31.3-42.0 | |
| National Median | 17.5 | | 13.0 | | 29.8 | | 9.9 | | 30.9 | | 48.7 | | 36.3 | |

NOTES: *Prevalence weighted to landline and cellphone households. BRFSS data from 2011-2015 are not directly comparable to previous years.
 Each percent has a different standard error and confidence interval.
Because of this, rankings are not truly meaningful.
 Unknown and refused responses are not included.
 A ranking between 1 and 25 (above the median value) implies poor status relative to other states.
 Data obtained from CDC BRFSS Prevalence and Trends Data accessed December 29, 2015